



264-601

01/11/2007

1/11

 <p>U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460</p>	EPA Registration Number: 264-601	Date of Issuance: JAN 11 2007
	Term of Issuance: Conditional	
	Name of Pesticide Product: Triticonazole SC Fungicide	
NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration <small>(under FIFRA, as amended)</small>		
Name and Address of Registrant (include ZIP Code): Bayer Crop Science LP P.O. Box 12014, 2 T.W. Alexander Drive Research Triangle Park, NC 27709		
<small>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</small>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3 (c) (7) (A) provided that you:</p> <ol style="list-style-type: none"> 1. Submit/cite all data required for the registration/ reregistration of your product when the Agency requires all registrants of similar products to submit data; and submit acceptable responses required for reregistration of your product under FIFRA section 4. 2. Make the following label changes <ol style="list-style-type: none"> a. Change the EPA Registration Number to 264-601 		
(Continued)		
Signature of Approving Official:  Mary L. Waller, Product Manager 21 Fungicide Branch Registration Division (7505P)	Date: 1/11/2007	

2. Label changes (continued)

b. Page 0 – Heading:

Change “For use by commercial applicators only.” to “Intended for use by commercial applicators only.”

c. Page 0 – Ingredient statement:

- Add a line below the inert ingredients line with “Total” directly under “inert ingredients” and 100.0% under the “30.0%” so that the decimal points fall along the same vertical line.
- Change “* Contains 1.69 lbs. active Triconazole per U.S. gallon.” to “* Contains 1.69 lbs. Triconazole per U.S. gallon.”
- The Agency suggests you replace “Inert Ingredients” with “Other Ingredients”

d. Page 1 – Chemigation statement:

Move the statement “Do not apply Triconazole 70 WDG Fungicide” through any type of irrigation system.” to the “Restrictions and Limitations” section on page 6

e. Page 2 – Turf – General Information:

- In the heading beginning Turf (Golf courses . . .) change “For use only by licensed, commercial applicators.” to “Intended for use only by licensed, commercial applicators.”
- In the second sentence of the text of this section, change “Only licensed, commercial application is approved.” to “Intended for use only by licensed, commercial applicators.”
- Change the sentence “Begin applications when conditions favor disease development per local environment unless otherwise noted.” to “Begin applications when local environmental conditions favor disease development unless otherwise noted.”

f. Page 3 – Change the table heading “Recommendations for Use” to “Application Directions”.

g. Page 6 – Restrictions and Limitations:

Add the following sentences to the third bullet “No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against mixing.”

h. Page 7 – Warranty Language:

At the end of the “Conditions” paragraph, change the sentence “All such risks shall be assumed by the user or buyer.” to “To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.”

3. Provide the following studies as conditions of registration:

- a. OPPTS number 830.6316 - storage stability - due May 1, 2007.

3/11

- b. OPPTS number 830.6320 - corrosion characteristics - due May 1, 2007
- c. OPPTD number 830.1800 – enforcement analytical method – due August 1, 2007.
- d. A biliary excretion study performed with triticonazole – due February 1, 2009.

4. Submit one (1) copy of the revised final printed label for the record before the product is released for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

In the Agency's risk assessment for 1,2,4-triazole and its metabolites triazole alanine, and triazole acetic acid, dated February 7, 2006 (see docket number 2005-0497 at <http://www.regulations.gov>), the Agency identified additional data that are needed for triazole-derivative active ingredients which have the 1,2,4,-triazole ring. The Agency plans to issue a Data Call-In for all triazole-derivative active ingredients which will identify all data needs for these active ingredients listed in the February 7, 2006 risk assessment. The Data Call-In will not address data needs that are related to parent-only data requirements.

A stamped copy of the label is enclosed for your records. If you have any questions, contact Lisa Jones of my staff at (703) 308-9424 or at jones.lisa@epa.gov.

Sincerely,



Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505P)

Enclosures: Stamped copy of label

4/11

Triticonazole SC Fungicide

A Fungicide For The Prevention And Control Of Certain Diseases Of Residential and Commercial Turfgrass, Golf Courses and Sod Farms. For Use by Commercial Applicators Only.

ACTIVE INGREDIENT:

Triticonazole*:	19.2%
INERT INGREDIENTS:	80.8%
	TOTAL 100.0%

* Contains 1.69 lbs. active Triticonazole per U.S. gallon.

EPA Reg. No. 264-ANR

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN

CAUTION

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

FIRST AID

IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • Do not give anything by mouth to an unconscious person.

For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

NOTE TO PHYSICIAN: No specific antidote is available. Treat symptomatically.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

JAN 11 2007

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.**

264-601

5/11

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, socks and shoes, and chemical-resistant gloves made of waterproof material. Wash hands and exposed skin thoroughly after handling the concentrate and after application.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove Personal Protective Equipment immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

DIRECTIONS FOR USE

**It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
Read entire label before using this product.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is coveralls, chemical-resistant gloves made out of any waterproof material, and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or green houses.

Keep unprotected persons out of treated areas untils prays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Store in original container. Keep container closed. Store in an area that is out of reach of children and animals, away from food, feedstuffs, fertilizers and seed. Protect from frost and freezing.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill or by other procedures approved by State and local authorities.

Do not apply Triticonazole Fungicide through any type of irrigation system.

6/11

TURF

GENERAL INFORMATION

TURF (Golf courses, Commercial turfgrass, Sod Farms and Residential Turf). For use only by licensed, commercial applicators.

Triticonazole SC Fungicide is a foliar applied fungicide, recommended for turfgrass disease control on golf courses, commercial turfgrass, sod farms and residential turf where fine turf is grown. Only licensed, commercial application is approved. When used in conjunction with good turf management practices, Triticonazole Fungicide is effective in controlling the following diseases:

Dollar Spot	Anthraxnose	Red Leafspot	Red Thread
Brown Patch	Take-All Patch	Gray Snow Mold	Large Brown Patch
Summer Patch	Rust	Pink Snow Mold	

Triticonazole SC Fungicide is also effective in controlling the following diseases when combined with other products:

- Pythium Blight
- Summer Stress Complex / Decline

Begin applications when conditions favor disease development per local environment unless otherwise noted. Apply with sufficient water to obtain adequate coverage (1-5 gallons of spray solution per 1000 sq. ft.).

Recommended Turf Grass Species

- Kentucky Bluegrass
- Poa Annua
- Tall Fescue
- Fine Fescue
- Bentgrass (Creeping, Colonial)
- Perennial Ryegrass
- Zoysia

Triticonazole SC Fungicide on Turf Treated with Plant Growth Regulators

Triticonazole SC Fungicide has been tested and shown no adverse effects when applied to recommended turf types with the following Plant Growth Regulators (PGR's):

PGR	Acceptable Turf Types
CHIPCO® Proxy™ (Ethephon)	Kentucky Bluegrass Perennial Rye Bentgrass Tall Fescue
Primo® (Trinexapac-ethyl)	Kentucky Bluegrass Perennial Rye Bentgrass Tall Fescue
Embark® (Mefluidide)	Bentgrass Tall Fescue

7/11

RECOMMENDATIONS FOR USE

<u>DISEASE</u>	USE DIRECTIONS	RATE		Notes
		FL. OZ. PRODUCT PER 1000 FT²		
		Preventative	Curative	
Dollar Spot (Lanzia spp. and Moellerodiscus spp.)	Repeat applications at 14-28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under light disease pressure use the lower curative rate and shorter interval. Under severe disease conditions use the higher rate and shorter interval.	0.5-0.75 fl.oz.	0.75-1.0 fl.oz.	Triticonazole SC is compatible with ProStar®, Compass®, and Heritage™. For control of Dollar Spot, combine Triticonazole SC at labeled rates with ProStar®, Compass® or Heritage® at labeled rates.
Brown Patch (Rhizoctonia solani)	Repeat applications at 14-28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under active disease conditions use the curative rate and shorter interval.	0.5-1.0 fl.oz.	1.0-2.0* fl.oz. *2.0- fl.oz.rate may be applied if needed in Transition areas of the South under heavy disease pressure.	In conditions favoring moderate to heavy disease pressure, Triticonazole SC can be combined with other fungicides labeled for Brown Patch including ProStar® or 26GT® at labeled rates.
Summer Patch (Magnaporthe poae)	Begin applications in spring (April-May) and repeat at 14-28 day intervals per the following situation: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals.	1.0-2.0 fl.oz.		
Anthracnose (Colletotrichum sp.)	Repeat applications at 14-28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under light disease pressure use the lower curative rate and shorter interval. Under severe disease conditions use the higher rate and shorter interval.	0.5-0.75 fl.oz.	0.75-1.0 fl.oz.	

<p>Take-All Patch (<i>Gaeumannomyces graminis</i>)</p>	<p>Make 1-2 fall applications (September-October) and 1-2 spring applications (April-May) depending on local conditions. Repeat applications at 14-28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under active disease conditions use the curative rate and shorter interval.</p>	<p>0.5-1.0 fl.oz.</p>	<p>1.0 fl.oz.</p>	
<p>Large Brown Patch (<i>Rhizoctonia solani</i>)</p>	<p>Repeat applications at 14-28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under active disease conditions use the curative rate and shorter interval.</p>	<p>0.5-1.0 fl.oz.</p>	<p>1.0 fl.oz.</p>	
<p>RED LEAFSPOT (<i>Drechslera erythrospila</i>)</p>	<p>Repeat applications at 14-28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under light disease pressure use the lower curative rate and shorter interval. Under severe disease conditions use the higher rate and shorter interval.</p>	<p>0.5-0.75 fl.oz.</p>	<p>0.75-1.0 fl.oz.</p>	
<p>ALGAE</p>	<p>For algae suppression repeat applications at 14-28 day intervals per the following situation: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals.</p>	<p>0.5-1.0 fl.oz.</p>		
<p>Rust (<i>Puccina spp.</i>)</p>	<p>Repeat applications at 14-28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under light disease pressure use the lower curative rate and shorter interval. Under severe disease conditions use the higher rate and shorter interval.</p>	<p>0.5-0.75 fl.oz.</p>	<p>0.75-1.0 fl.oz.</p>	

GRAY SNOW MOLD <i>(Typhula spp)</i>	<p>Make 1-2 applications late in the fall before snow cover occurs. Repeat applications at 14-28 day intervals per the following situation: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals.</p>	<p>0.5-1.0 fl.oz.</p>		<p>Triticonazole SC can be combined with the following fungicides for greater efficacy: ProStar, Chlorothalonil, PCNB, Chipco Signature or 26 GT. Use only at labeled rates.</p>
PINK SNOW MOLD <i>(Microdochium)</i>	<p>Make 1-2 applications late in the fall before snow cover occurs. Repeat applications at 14-28 day intervals per the following situation: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals.</p>	<p>0.5-1.0 fl.oz.</p>	<p>1.0 fl.oz.</p>	<p>Triticonazole SC can be combined with the following fungicides for greater efficacy: Chlorothalonil, PCNB, Chipco Signature or 26 GT. Use only at labeled rates.</p>
RED THREAD	<p>Repeat applications at 14-28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under light disease pressure use the lower curative rate and shorter interval. Under severe disease conditions use the higher rate and shorter interval.</p>	<p>0.5-0.75 fl.oz.</p>	<p>0.75-1.0 fl.oz.</p>	
<p>Do not apply more than 6 fluid ounces per 1000 sq. ft of Triticonazole SC Fungicide annually</p>				

Additional Tank Mixtures

DISEASE	USE DIRECTIONS
PYTHIUM BLIGHT	<p>Combine Triticonazole SC Fungicide with either CHIPCO® Signature™ or Banol® brand fungicide for control of Pythium Blight. Apply at labeled rates.</p>
SUMMER STRESS COMPLEX/SUMMER DECLINE	<p>Combine Triticonazole SC Fungicide with CHIPCO® Signature™ for diseases associated with Summer Decline. Apply at labeled rates.</p>
<p>Do not apply more than 6 fluid ounces per 1000 sq. ft of Triticonazole SC Fungicide annually.</p>	

10/11

MIXING RATES

at 0.5 ounce per 1000 FT.²

SPRAY VOLUME (Gallons Per 1000 FT.²)

	1 Gallon	2 Gallons	3 Gallons	4 Gallons	5 Gallons	
Spray Tank Capacity	25 Gallons	12.5 fl.oz.	6.3 fl.oz.	4.2 fl.oz.	3.1 fl.oz.	2.5 fl.oz.
	50 Gallons	25 fl.oz.	12.5 fl.oz.	8.3 fl.oz.	6.3 fl.oz.	5 fl.oz.
	100 Gallons	50 fl.oz.	25 fl.oz.	16.7 fl.oz.	12.5 fl.oz.	10 fl.oz.
	200 Gallons	100 fl.oz.	50 fl.oz.	33.3 fl.oz.	25 fl.oz.	20 fl.oz.

at 0.75 fluid ounce per 1000 FT.²

SPRAY VOLUME (Gallons Per 1000 FT.²)

	1 Gallon	2 Gallons	3 Gallons	4 Gallons	5 Gallons	
Spray Tank Capacity	25 Gallons	18.8 fl.oz.	9.4 fl.oz.	6.3 fl.oz.	4.7 fl.oz.	3.75 fl.oz.
	50 Gallons	37.5 fl.oz.	18.8 fl.oz.	12.5 fl.oz.	9.4 fl.oz.	7.5 fl.oz.
	100 Gallons	75 fl.oz.	37.5 fl.oz.	25 fl.oz.	18.8 fl.oz.	15 fl.oz.
	200 Gallons	150 fl.oz.	75 fl.oz.	50 fl.oz.	37.5 fl.oz.	30 fl.oz.

® at 1.0 fluid ounce per 1000 FT.²

SPRAY VOLUME (Gallons Per 1000 FT.²)

	1 Gallon	2 Gallons	3 Gallons	4 Gallons	5 Gallons	
Spray Tank Capacity	25 Gallons	25 fl.oz.	12.5 fl.oz.	8.3 fl.oz.	6.3 fl.oz.	5 fl.oz.
	50 Gallons	50 fl.oz.	25 fl.oz.	16.7 fl.oz.	12.5 fl.oz.	10 fl.oz.
	100 Gallons	100 fl.oz.	50 fl.oz.	33.3 fl.oz.	25 fl.oz.	20 fl.oz.
	200 Gallons	200 fl.oz.	100 fl.oz.	66.7 fl.oz.	50 fl.oz.	40 fl.oz.

NOTES

- Apply with sufficient water to obtain adequate coverage (1 to 5 gallons of spray solution per 1000 square feet.
- Apply with a properly calibrated sprayer and maintain agitation during spray operations.

RESTRICTIONS AND LIMITATIONS

- Only for use on labeled grasses.
- For golf courses only, do not apply to turf cut higher than 1" on golf holes where water bodies are present.
- This product is compatible with most commonly used fungicides, however, check compatibility before tank mixing.
- Mixture with stickers, extenders, or wetting agents is not necessary.
- Do not graze animals on treated turf.
- Do not feed clippings from treated turf to livestock or poultry.

11111

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCES ELECTION, THE REPLACEMENT OF PRODUCT.

Net Content:



Bayer CropScience

**Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)**

Triticonazole SC Fungicide (PENDING) Submitted 07/18/02, Resubmitted 05/16/05, Resubmitted 09/22/06, Resubmitted 01/08/07