

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0080. Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 264-622	2. EPA Product Manager Joanne Miller	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) DROPP® 50WP Cotton Defoliant	PM# Team 23	
5. Name and Address of Applicant (Include ZIP Code) Bayer CropScience P.O. Box 12014, 2 T.W. Alexander Drive Research Triangle Park, NC 27709 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION
JAN 20 2006

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of minor labeling changes per PR Notice 98-10 (see attached letter for details) - REGFEE - Not applicable
CERTIFICATION STATEMENT: This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III


1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container	<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Danielle A. Larochelle	Title Registration Product Manager	Telephone No. (Include Area Code) (919) 549-2718

Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Registration Product Manager	
4. Typed Name Danielle A. Larochelle	5. Date December 22, 2005	

CropScience



December 22, 2005

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504C)
U.S. Environmental Protection Agency
1801 South Bell Street
Arlington, Virginia 22202-4501
(PM: Ms. Joanne Miller, Team 23)

Dear Reviewer,

Re: **DROPP® 50WP Cotton Defoliant - EPA Reg. No. 264-622**
Notification of Minor Labeling Change per PR Notice 98-10

In accordance with PR Notice 98-10, we are notifying the Agency of labeling amendments for DROPP® 50WP Cotton Defoliant (EPA Reg. No. 264-622). Minor editorial changes were made in the "IMPORTANT CLEANOUT INSTRUCTIONS" section of the label (page 5) to clarify the instructions and remove references to a separate technical information bulletin. Please note also that the second paragraph of the "SEQUENTIAL APPLICATION PRECAUTION" section was amended and moved to the "IMPORTANT CLEANOUT INSTRUCTIONS".

The following documents are enclosed in support of this action:

- A completed Application for Registration (EPA Form 8570-1) which includes the Certification Statement required by PR Notice 98-10.
- One (1) copy of the label with changes highlighted.

Please call me at 919 549-2718 if you need any additional information.

Sincerely,

Danielle A. Larochelle
Registration Product Manager

Corr = dal.078-05

3 7 8

NOTIFICATION

JAN 20 2006

Dropp® 50WP COTTON DEFOLIANT

Packed In Water Soluble Bags

For Agricultural Use Only

ACTIVE INGREDIENT: Thidiazuron*..... 50%

INERT INGREDIENTS:..... 50%

*N-phenyl-N'-1,2,3-thiadiazol-5-ylurea

TOTAL 100%

EPA Reg No. 264-622

EPA Est. No. 33689-WG-1

U.S. Patent No. 4,294,605

KEEP OUT OF REACH OF CHILDREN
CAUTION – CAUCION

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • Do not give anything by mouth to an unconscious person.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if Swallowed, Inhaled, or Absorbed Through the Skin. Avoid Breathing Spray Mist. Avoid Contact with Skin, Eyes, and Clothing. Do Not Contaminate Food or Feedstuffs.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target area.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Do not store box under wet conditions. Handle box carefully when stored at temperatures of less than 50°F to avoid breakage of soluble bags.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty box into application equipment. Then dispose of empty box in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DO NOT REUSE EMPTY CONTAINER

GENERAL INFORMATION

This box contains DROPP® 50WP cotton defoliant in water soluble bags. Inner bags dissolve in water and contents disperse. After opening outer box, open foil liner, immediately dump entire unopened inner bags into the mix tank. Do not handle soluble bags or expose them to moisture, since this may cause breakage.

DROPP® 50WP cotton defoliant is used for removal of leaves prior to harvesting. DROPP® disperses easily in water and can be used in any ground and aircraft equipment designed for application of wettable powders. It may take several days before the effect of DROPP® becomes noticeable. Adverse conditions, such as low temperatures, may require higher dosages and/or longer times for more complete defoliation. DROPP® inhibits regrowth after defoliation and thus reduces staining of lint during harvesting and ginning.

USE PRECAUTIONS

Do not apply this product through any type of irrigation system.

Rainfall within 24 hours after application will reduce the effectiveness of DROPP®. Use only freshly prepared sprays. Do not store spray mixture overnight.

Do not feed foliage from treated cotton plants or gin trash to livestock.

Do not plant the following crops **earlier than** the specified periods after the application of DROPP® 50WP:

Small grain, sorghum, corn.....	two (2) weeks
or root crops (except carrots, onions, and sugar beets)	
Legumes (including alfalfa).....	two (2) months
or leafy vegetables (except lettuce and spinach)	
Sugar beets	four (4) months
Carrots, onions, or spinach.....	nine (9) months
Lettuce.....	nine (9) months
	and only following deep plowing (12-15 inches).

Do not use immature crops for food or feed.

Do not allow spray drift to contact crops other than the target crop of mature cotton, or cotton which you desire to defoliate, as this product may injure or defoliate other crops.

Particular care should be taken when applying DROPP® 50WP adjacent to lettuce, citrus, or cantaloupe. Tank mixes with organophosphates may increase nontarget crop phytotoxicity.

PRACTICES TO LOWER THE POTENTIAL FOR DRIFT ONTO NONTARGET CROPS

During applications, particularly under windy conditions, DROPP® 50WP may drift to nontarget crops.

To help reduce the drift potential, use the following practices:

- Do not apply DROPP® 50WP by ground or air when wind speeds exceed ten (10) miles per hour at the time of application. Follow local recommendations if wind speeds of less than ten (10) miles per hour are specified in those recommendations.
- Use of low nozzle pressure (20–30 psi) is recommended.
- Do not apply DROPP® 50WP when a temperature inversion is present or when conditions favor an inversion prior to completing application(s).
- Additionally, do not apply DROPP® 50WP by air within one-half (1/2) mile of lettuce. Do not apply DROPP® 50WP by ground equipment within 100 feet of lettuce.

PRACTICES TO LOWER DRIFT POTENTIAL ONTO NON-TARGET CROPS IN THE RIO GRANDE VALLEY OF TEXAS

To help reduce the drift potential, use the following practices:

- Do not apply DROPP® 50WP by ground or air when wind speeds exceed ten (10) miles per hour at the time of application. Follow local recommendations if wind speeds of less than ten (10) miles per hour are specified in those recommendations.
- Use of low nozzle pressure (20–30 psi) is recommended.
- Do not apply DROPP® 50WP when a temperature inversion is present or when conditions favor an inversion prior to completing application(s).
- Use the largest nozzle orifice possible which permits proper deposition and coverage of product.
- In addition for citrus crops, do not apply DROPP® 50WP by air when citrus in flush is within five (5) miles downwind of the point of application. Do not apply DROPP® 50WP by ground when citrus in flush is within one-half (1/2) mile downwind of the point of application.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

TIME OF APPLICATION

Apply DROPP® 50WP cotton defoliant only to mature cotton plants when the last boll you expect to harvest is mature. A boll can be described as mature when it is too hard to be dented when squeezed between thumb and fingers, is difficult to slice with a sharp knife, and/or when seeds cut in cross sections have fully developed cotyledons, as evidenced by an absence of jelly within the seed.

Apply DROPP® 50WP at least 5 days prior to anticipated harvest.

NOTE: Good defoliation with DROPP® is dependent upon three factors:

1. Actively growing cotton
2. High humidity
3. High moisture content in leaf of cotton plant

Use of DROPP® 50WP alone (without a tank mix partner) when nighttime temperatures are expected to fall below 60°F can result in less than desirable defoliation and/or regrowth inhibition.

Performance of DROPP® 50WP is variable under low temperatures. Ideally nighttime temperatures 2–3 days prior to and following application should be above 60°F or total defoliation and regrowth inhibition can be reduced.

Use of DROPP® 50WP on heat and drought-stressed cotton (low leaf moisture, thick cuticle, etc.) may result in less than satisfactory defoliation and regrowth inhibition.

8 8 8

USE OF ADJUVANTS (CROP OILS)

The use of adjuvants such as petroleum-based crop oils or penetrating oils approved for use on growing crops has been shown to improve performance during low nighttime temperatures (60°–65°F) or when defoliating drought-stressed cotton. Refer to specific adjuvant label for rate recommendations.

The use of adjuvants is recommended when applying DROPP® 50WP in the desert Southwest (i.e., Arizona and California).

PRECAUTION: The addition of adjuvants can cause desiccation and/or leaf freezing during periods of high temperatures. The use of compounds that desiccate leaf tissue is not recommended.

MIXING INSTRUCTIONS

Fill the spray tank with one-half of the total amount of water to be used. After opening each box to be used, open its foil liner. Immediately dump the entire contents of the required number of boxes of soluble bags into the mix tank. Fill the tank with the additional quantity of water required and then start the agitator. Use entire contents of each box.

(WHEN PRINTING FREEFALL PACKAGING): Fill the spray tank with one-half of the total amount of water to be used. Open each bag to be used and immediately dump the entire contents of the required number of bags into the mix tank. Fill the tank with the additional quantity of water required and then start the agitator. Use entire contents of each bag.

TANK MIX ADJUVANTS

When using tank mixes of DROPP® 50WP and organophosphates, the use of a surfactant or compatibility agent is recommended to improve tank clean-out and overall defoliation. Prior to mixing in the spray tank, a small scale compatibility test should be conducted.

Use only products which are exempt from tolerance under 40 CFR 180.1001.

Always follow the mixing instructions on the label of the appropriate adjuvant.

APPLICATION

Use the specified dosage of DROPP® 50WP cotton defoliant in sufficient amount of water to give complete coverage of the foliage with uniform wetting of leaf surfaces. Agitate the spray mix during application.

DOSAGE

One box (5 x 1lb bags) of DROPP® treats 12.5–25 acres (0.2–0.4lbs DROPP® per acre). Apply in 10–25 gallons of spray per acre by ground equipment and 2–10 gallons per acre by air. Use the higher rates during periods of low temperatures (60°–65°F).

Two applications not exceeding a total of 1 box per 8.3 acres (0.6 lbs./acre) may be necessary to defoliate rank cotton.

BOTTOM DEFOLIATION FOR USE IN ARIZONA ONLY: DROPP® 50WP may be used as an aid to improve air circulation and help reduce boll rot caused by moisture.

Apply DROPP® 50WP at 0.1 lbs. formulated product per acre in 10–25 gallons of water per acre.

Application to **bottom** 1/3 of plant **only** recommended.

Timing of application should be based upon the latest date when ground application equipment can enter the field without crop damage.

PRECONDITIONING FOR USE IN ARIZONA AND CALIFORNIA ONLY: DROPP® 50WP may be used as a pre-conditioner to enhance the activity of a defoliant application.

Apply DROPP® 50WP at 0.1–0.2 lbs. formulated product per acre. Apply in 10–25 gallons of water per acre by ground application or 2–10 gallons of water per acre by aerial application.

Timing of application is recommended to be 7–14 days prior to the use of another defoliant. Refer to the second product label prior to use for complete recommendations.

TANK MIX OF DROPP® 50WP PLUS PREP™ FOR BOLL OPENING: The tank mix of DROPP® 50WP plus Prep™ is recommended to improve overall defoliation, and as an aid in accelerating the opening of mature, unopened cotton bolls. Best activity will be obtained where the tank mix is applied to mature cotton plants. (See Time Of Application section of this label.) Do not apply tank mix before sufficient unopened bolls have matured to produce the desired cotton yield. If nighttime temperatures are expected to fall below 60°F, unsatisfactory defoliation and regrowth inhibition may result. Certified cotton seed producers should determine boll and seed maturity prior to treatment.

Apply DROPP® 50WP at a rate of 0.1–0.4 lbs. formulated product plus Prep™ at 1.0–2.0 lbs. active ingredient per acre. The higher rate of DROPP® should be used where excessive regrowth is anticipated, or during cooler temperatures.

Apply as a dilute spray as follows:

<u>Application Method</u>	<u>Gallons of Water/Acre</u>
Aerial	3–10
Ground	10–50

TANK MIX OF DROPP® 50WP PLUS PREP™ FOR DEFOLIATION ENHANCEMENT: Lower rates of Prep™ may be tank mixed with DROPP® 50WP to enhance defoliation without providing boll opening activity. Prep™ may still be applied as a sequential treatment for boll opening following an application of DROPP® 50WP plus Prep™ at a defoliation enhancement rate. Read and follow the Prep™ label, and do not exceed a maximum of 2.0 lb. ai/acre for combined uses of PREP™ per acre per year.

7 7 8

Apply DROPP® 50WP at a rate of 0.1–0.4 lbs. formulated product plus Prep™ at 1/3 pint per acre (0.25 lbs. ai/acre). The higher rates of DROPP® 50WP should be used where excessive regrowth is anticipated, or during cooler temperatures.

TANK MIX OF DROPP® 50WP PLUS DEF® 6 OR DROPP® 50WP PLUS FOLEX® 6EC: The tank mix of DROPP® 50WP and Def® 6 or Folex® 6EC is recommended to improve defoliation and inhibit regrowth under less than ideal conditions. Less than ideal conditions are those when the nighttime temperatures are expected to be 60°F on the date of application and for three days following application.

Best activity will be obtained when the tank mix is applied to mature cotton plants with 60% or more open bolls.

RATE OF APPLICATION

DROPP® 50WP + Def® 6 = Use 0.1–0.4 lbs

DROPP® 50WP + 0.5–2.0 pints Def® 6

DROPP® 50WP + Folex® 6EC = Use 0.1–0.4 lbs

DROPP® 50WP + 0.5–2.0 pints Folex® 6EC

Higher rate of DROPP® 50WP should be used where excessive regrowth is anticipated.

Maximum rates of DROPP® and phosphate defoliant applied in a tank mix during very high air temperatures can cause desiccation .

Use lower rate of Def® 6 or Folex® 6EC under ideal nighttime temperature conditions (65°F); higher rate during less than ideal or adverse conditions.

Apply as a dilute spray in 5–10 gallons of water per acre by air or 10–25 gallons of water per acre with ground equipment.

When applying Def® 6 or Folex® 6EC defoliant plus DROPP® 50WP as a tank mix, the following mixing sequence is recommended:

1. Add required water to spray tank, agitate.
2. Add DROPP® 50WP according to label directions (see **mixing instructions**).
3. Add Def® 6 or Folex® 6EC according to rate desired **after** the DROPP® 50WP has completely dispersed.

A second application of the labeled tank mix may be made where necessary, but not exceeding a total of 0.6 lbs DROPP® per acre.

Refer to Def® 6 and Folex® 6EC labeling for additional use directions and cautions when using tank mixtures of DROPP® 50WP and those products.

DO NOT USE TANK MIX OF DROPP® 50WP PLUS DEF® 6 OR FOLEX® 6EC IN THE RIO GRANDE VALLEY COUNTIES OF STARR, HIDALGO, WILLACY, AND CAMERON.

IMPORTANT CLEANOUT INSTRUCTIONS

Tank mixes of DROPP® 50WP with organophosphates may form a residue in application equipment. **While still fresh and moist**, this residue can be removed by flushing the entire system with a commercial tank cleaner. Consult your State Extension cotton specialist for recommended tank cleaners and cleaning procedures.

Do not allow the spray solution to dry in the application equipment. Immediately following application, clean all equipment (mix tanks, pumps, transfer lines, application tanks, sumps, booms, nozzles and all related equipment) thoroughly with a commercial tank cleaner and water.

The procedure for removing dried residues in the application equipment requires allowing the diluted solution of a commercial tank cleaner to stand in equipment, filled to capacity, for 7 days followed by thorough flushing.

Should small quantities of DROPP® 50WP remain in inadequately cleaned equipment, they may be released during subsequent applications and may cause damage to crops. Bayer CropScience accepts no liability for damage to crops due to inadequately cleaned equipment.

SEQUENTIAL APPLICATION PRECAUTION

Do not apply sequential application of DROPP® 50WP following any defoliant or desiccant **except** DROPP® 50WP used alone or DROPP® 50WP tank mixes. Reduced activity of the second treatment will result.

8 2 8

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

NET CONTENTS:

Prep is a trademark and Def is a registered trademark of Bayer.
Folex is a registered trademark of Amvac Chemical.



Bayer CropScience

Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

Dropp® 50WP (MASTER) Approved 10/1/02, Notification 3/17/04, Notification 05/25/05