



264-652

12-27-2000

1/7

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 2-28-95

| | | | |
|--|---|---|-----------------------|
| | United States Environmental Protection Agency Washington, DC 20460 | <input type="checkbox"/> Registration <input checked="" type="checkbox"/> Amendment <input type="checkbox"/> Other | OPP Identifier Number |
|--|---|---|-----------------------|

Application for Pesticide - Section I

| | | |
|---|---|--|
| 1. Company/Product Number 264-652 | 2. EPA Product Manager Joanne I. Miller | 3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted |
| 4. Company/Product (Name) Relv®Herbicide | PM# 23 | |
| 5. Name and Address of Applicant (Include ZIP Code) <input type="checkbox"/> Check if this is a new address | 6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____ | |

Section - II

NOTIFICATION

| | |
|--|--|
| <input checked="" type="checkbox"/> Amendment - Explain below. | <input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated <u>DEC 27 2000</u> |
| <input type="checkbox"/> Resubmission in response to Agency letter dated _____ | <input type="checkbox"/> "Me Too" Application. |
| <input checked="" type="checkbox"/> Notification - Explain below. | <input type="checkbox"/> Other - Explain below. |

Explanation: Use additional page(s) if necessary. (For section I and Section II.)
 Rely Herbicide registration for use on apples, grapes, and tree nuts was recently approved by California Department of Pesticide Regulation. The potato vine desiccation use was not included in the application. Consequently, a statement was added to the Application Directions for Potato Vine Desiccation to indicate that this use is not registered in California.

Section - III

1. Material This Product Will Be Packaged In:

| | | | |
|---|--|---|---|
| Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No | Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No | Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No | 2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____ |
| * Certification must be submitted If "Yes" Unit Packaging wgt. No. per container | | If "Yes" Package wgt No. per container | |

3. Location of Net Contents Information
 Label Container

4. Size(s) Retail Container

5. Location of Label Directions

6. Manner in Which Label is Affixed to Product
 Lithograph
 Paper glued
 Stenciled Other _____

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

| | | |
|-----------------------------|--------------------------------------|---|
| Name Paul A. Cain | Title Registration Manager | Telephone No. (Include Area Code) _____ |
|-----------------------------|--------------------------------------|---|

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

| | | |
|--------------------------------------|---|---|
| 2. Signature | 3. Title Registration Manager | 6. Date Application Received (Stamped) . |
| 4. Typed Name Paul A. Cain | 5. Date December 14, 2000 | |

Rely[®] HERBICIDE

For Control of Emerged Weeds in Apples, Grapes, and Tree Nuts; and for Potato Vine Desiccation

| | |
|--|---------------|
| ACTIVE INGREDIENT: glufosinate-ammonium* | 11.33%** |
| OTHER INGREDIENTS: | 88.67% |
| * CAS Number 77182-82-2, Protected by U.S. Patent No. 4,400,196 | TOTAL 100.00% |
| **Equivalent to 1.00 pound of active ingredient per U.S. gallon. | |

EPA Reg No. 264-652

EPA Est. No. 264-MI-001
407-IA-2

KEEP OUT OF REACH OF CHILDREN WARNING — AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-888-AVENTIS (1-888-283-6847)

FIRST AID

| | |
|-------------------------------|--|
| IF SWALLOWED | <ul style="list-style-type: none"> Rinse mouth thoroughly with plenty of water. Do not induce vomiting. Get medical attention immediately. |
| IF IN EYES | <ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Get medical attention if irritation develops or persists. |
| IF ON SKIN OR CLOTHING | <ul style="list-style-type: none"> Take off contaminated clothing. Wash skin immediately with plenty of soap and water. Get medical attention. |
| IF INHALED | <ul style="list-style-type: none"> Move person to fresh air. Get medical attention if breathing difficulty develops. |

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or when going for treatment. Call 1-800-334-7577 for emergency medical treatment information.

NOTE TO PHYSICIAN

If this product is ingested, endotracheal intubation and gastric lavage should be performed as soon as possible, followed by charcoal and sodium sulfate administration. Additionally, call 1-800-334-7577 immediately for further information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Causes substantial but temporary eye injury. Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- long-sleeved shirt and long pants

- chemical-resistant gloves such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton[®] ≥ 14 mils
- shoes plus socks
- protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not clean equipment or dispose of equipment washwaters in a manner that will contaminate water resources or arable land. Glufosinate-ammonium and its degradates have those properties normally associated with pesticides that have been detected in groundwater. Use of this product in areas with coarse soils and high water tables may result in groundwater contamination.

GENERAL INFORMATION

RELY[®] Herbicide is a nonselective herbicide for application as a foliar spray for the control of a broad spectrum of emerged annual and perennial grass and broadleaf weeds. RELY[®] will also control certain woody species. Plants that have not yet emerged at the time of application will not be controlled. THOROUGH SPRAY COVERAGE IS IMPORTANT. Visual effects and control from application of RELY[®] occur within 2 to 4 days after application under good growing conditions.

Avoid all contact with foliage or green tissue of desirable vegetation. Do not spray during windy conditions. This product may injure or kill growing plants that receive spray drift or if they receive applications of a spray mixture containing RELY[®] Herbicide by error or accident. If desirable vegetation is contacted, rinse the sprayed portion with water immediately to reduce potential injury.

RELY[®] Herbicide works best when weeds are actively growing. Warm temperatures, high humidity, and bright sunlight improve the performance of RELY[®]. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. Weeds under stress or in dense populations will require application at the highest rate recommended. Refer to the *How to Apply* section of this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not use this product until you have read the entire label. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

IN THE STATE OF NEW YORK ONLY: NOT FOR USE IN NASSAU AND SUFFOLK COUNTIES.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Do not use or store near heat or open flame.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Empty containers should be triple rinsed (or equivalent), then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill, or by incineration; or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry-interval (REI) of 12 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls; chemical-resistant gloves such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton[®] ≥ 14 mils; shoes plus socks; protective eyewear.

APPLICATION DIRECTIONS FOR USE ON TREE AND VINE CROPS

Apply RELY® Herbicide as a directed spray to control undesirable vegetation in apple orchards, grape vineyards, and in tree nut groves for tree nuts listed below. When applied as recommended in this label, RELY® controls annual and perennial weeds. Refer to the *How to Apply* section of this labeling for recommended rates and a list of weeds controlled. Apply as a broadcast, banded, or spot treatment application depending on the situation. Avoid direct spray or drift to desirable vegetation. Regrowth may occur due to the weed stage of growth at application, low use rate, or environmental conditions. Repeat applications of RELY® may be necessary to control plants generating from underground parts or seed.

| Tree Nuts | Vineyards | Tree Fruits |
|-------------------------------|---------------|-------------|
| almonds | all grape | apples |
| pecans | varieties | |
| walnuts | (table, wine, | |
| filberts | and raisins) | |
| hickory nuts | | |
| macadamia nuts (bush nuts) | | |

Do not apply on, or allow spray to drift onto, desirable foliage of trees and vines, as damage will occur. Avoid contact with green or uncallused bark on young trees and vines as injury may occur. Do not apply RELY® Herbicide on trees within one year of transplanting.

Refer to the *When to Apply*, *How to Mix*, and *How to Apply* sections of this label for information on how and when to properly mix and apply this product. Do not apply more than 18 quarts of RELY® Herbicide per acre to tree and vine crops in a 12-month period.

WHEN TO APPLY

RELY® Herbicide is a foliar-active material with little or no perceptive soil-residual activity. Best results are obtained when applications are made to actively growing weeds. RELY® is rainfast for most weeds within 4 hours after application. Rainfall within 4 hours may necessitate retreatment. Do not irrigate within 4 hours of treatment.

Apply RELY® Herbicide at the rate recommended in the *How to Apply* section of this label. Repeat applications of RELY® or tank mixes of RELY® with appropriate residual herbicides will be needed to control weeds emerging from underground parts or seeds.

When using RELY® Herbicide in a weed control program with other herbicides, follow the precautions and "Directions For Use" for those herbicides.

HOW TO MIX

Mix RELY® Herbicide with water to make a finished spray solution as follows:

1. Fill the spray tank with the required amount of water.
2. Add the proper amount of RELY® Herbicide.

HOW TO APPLY

Apply RELY® Herbicide by ground equipment only. Use backpack, pump-up, or hydraulic sprayers. Do not apply this product aerially. Do not apply this product through any type of irrigation system. Good spray coverage of target weeds is necessary for optimum control.

Broadcast Application

Use the recommended rate of RELY® Herbicide as shown in the following sections. Use a minimum of 20 gallons of water per acre with a minimum of 30 psi spray pressure. Properly calibrate your spraying equipment to assure applications at the correct rates and volumes. As density of weeds increases, increase spray gallonage and/or pressure to ensure complete coverage.

Banded Application

Banded applications may be used using the following formula to calculate the amount of herbicide needed:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Rate per acre broadcast} = \text{Amount of herbicide needed for treatment}$$

For best results with ground broadcast or banded application equipment, use flat fan nozzles. Check for even distribution of spray droplets. Do not apply when weather conditions favor drift, such as when winds are gusty. Applications under conditions which cause drift of this product will result in damage to any vegetation contacted.

Spot or Directed-Spray Application

For spot or spray application, use RELY® Herbicide at 1.5 to 4 fluid ounces per gallon of water. Mix 1.5 to 4 fluid ounces per gallon of water depending upon the weeds to be controlled, see list of weeds in the following sections on this label. Spray undesirable vegetation foliage on a spray-to-wet basis prior to runoff. Ensure uniform and complete coverage. Use a coarse spray. Do not spray during windy conditions. Use a backpack, pump-up, or hydraulic sprayer. Thoroughly clean the sprayer following use.


DO NOT make spot spray applications to apple suckers, as tree injury may occur.

Weeds Controlled by RELY® Herbicide

For spot application: Apply 1.5 fluid ounces per gallon of water.

For broadcast application: Apply 3 quarts per acre when the weeds are less than 6 inches tall.

Apply 5 quarts per acre when the weeds are 6 inches or taller.



Broadleaf Weeds

- chickweed
- clover
- common cocklebur
- common mallow
- filaree
- jimsonweed
- kochia
- London rocket
- malva (little mallow)
- maretail
- purslane
- shepherdspurse
- smartweed

Grasses and Sedges

- barnyardgrass
- cupgrass
- fall panicum
- giant foxtail
- goosegrass
- green foxtail
- Johnsongrass
- lovegrass
- shattercane
- stinkgrass
- windgrass
- yellow foxtail

For spot application:

Apply 3 to 4 fluid ounces per gallon of water.

For broadcast application:

Apply 4 quarts per acre when the weeds are less than 8 inches tall. Apply 6 quarts per acre when the weeds are 8 inches or taller.

Broadleaf Weeds

- annual sowthistle
- bindweed
- buffalobur
- burdock
- Canada thistle
- curly dock
- dandelion
- dogbank (hemp)
- fiddleneck
- field gromwell
- fleabane
- goldenrod
- groundsel
- horsetail
- lambquarters
- leafy spurge
- mugwort
- musk thistle
- nettle
- nightshade
- pennycress

- pigweed, red root
- plantain
- prickly lettuce
- ragweed
- redstem filaree
- Russian thistle
- tansy mustard
- velvetleaf
- vervain
- vetch
- Virginia cooperleaf
- white clover
- white heath aster
- wild buckwheat
- wild mustard
- wild onion
- wild radish
- wild rose
- wild turnip
- woodsorrel
- yellow rocket

Grasses and Sedges

- annual bluegrass
- bahiagrass
- barley
- Bermudagrass
- carpetgrass
- crabgrass
- dallisgrass
- downy brome
- fescue
- guineagrass
- Kentucky bluegrass
- nutsedge
- paragrass
- quackgrass
- ryegrass
- sandbur
- smooth brome
- torpedograss
- vaseygrass
- wheat
- wild oat

Woody Species


- Rubus spp.
- poison ivy/oak

Sucker Control with RELY® Herbicide

RELY® Herbicide will give excellent control of suckers without injury to mature trees or vines. For sucker control, apply a split application approximately 4 weeks apart at 4 quarts per acre. Coverage of all sucker foliage is necessary for optimum control. Suckers should not exceed 12 inches in length.

Tank Mixtures

Always predetermine the compatibility of labeled tank mixtures of this herbicide with water carrier by mixing small proportional quantities in advance. Always add individual formulations to the spray tank according to the following sequence (first to last): wettable powder, flowable, emulsifiable concentrate, drift control additive, water-soluble liquid followed by surfactant.



Mixing Procedures:

1. Fill the spray tank one-half full with water.
2. Add the tank mix product to the spray tank.
3. Fill the spray tank with water to the desired level.
4. Add the recommended amount of RELY® Herbicide; always add the RELY® last to help reduce foaming.
5. Nonionic antifoaming agents may be used.
6. Additional surfactant is not needed.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed.

Keep bypass line on or near bottom of tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh. Carefully select proper nozzle to avoid spraying a fine mist. For best results with conventional ground application equipment, the use of flat fan nozzles is suggested.

Tank Mix Recommendations for Tree and Vine Crops

RELY® Herbicide does not provide residual weed control. RELY® will not control underground, unexposed plants, or plant parts. Delay application until plants have emerged. Tank mix applications of RELY® plus the following residual herbicides are recommended for broad spectrum control and residual activity. Refer to label of companion tank mix products for directions, limitations, warnings, and list of weeds controlled.

Simazine 4L (Do not use on tree nuts.)

Simazine 80W (Do not use on tree nuts.)

Simazine 90 (Do not use on tree nuts.)

Karmex DF

Sinbar-80W* (Do not use on grapes.)

Solicam DF

Surflan A.S.

Devrinol 50WP

Goal 1.6E

Princep 4L

*For tank mix applications to tree nuts, Sinbar 80W may be mixed with RELY® Herbicide for use on pecans only.

RESTRICTIONS AND PRECAUTIONS FOR TREE AND VINE CROPS

1. DO NOT apply more than 4.5 gallons of this product per acre to any site in any calendar year.
2. DO NOT graze, harvest, and/or feed treated orchard cover crops to livestock.
3. DO NOT apply this product through any type of irrigation system.
4. DO NOT apply this product aerially.
5. DO NOT apply this product within 14 days of nut, apple, or grape harvest.
6. Compatibility of RELY® Herbicide and clay-based, dry-flowable herbicides should be determined before tank mixing.

APPLICATION DIRECTIONS FOR USE ON FARMSTEADS

When applied as recommended, this product controls undesirable plant vegetation in noncrop areas around farmstead building foundations, shelter belts, along fences, and general nonselective farmstead weed control. Refer to the *How to Apply* section of this labeling for appropriate application rates to control specific weeds.


APPLICATION DIRECTIONS FOR POTATO VINE DESICCATION*

Apply RELY® Herbicide at a rate of 3 pints per acre in 20 to 100 gallons of water per acre with ground equipment, or in 5 to 10 gallons per acre with aerial equipment. Use sufficient water for thorough coverage of potato vines. Where crop canopy is dense, better spray coverage will be achieved with the higher spray volumes.

For best results, apply RELY® Herbicide at the beginning of natural senescence of potato vines. Do not harvest potatoes earlier than 9 days after application. Do not apply to potatoes grown for seed stock.

Do not plant rotation crops in a field treated with RELY® Herbicide for 120 days after the last application of this product, with the exception of wheat, barley, buckwheat, millet, oats, rye, sorghum, and triticale which may be planted 70 days after the last application of this product. Corn and soybeans may be planted at any time.

* Not registered for use in California for Potato Vine Desiccation.



7/7

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Aventis CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: AVENTIS CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Aventis CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Aventis CropScience disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT AVENTIS CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

- Rely is a registered trademark of the Aventis Group.
- Karmex, Sinbar, and Viton are registered trademarks of DuPont Company
- Princep and Solicam are registered trademarks of Novartis.
- Surflan is a registered trademark of DowAgroSciences
- Devrinol is a registered trademark of Zeneca Inc., Ag Products
- Goal is a registered trademark of Rohm and Haas Co.

Aventis CropScience USA LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

RELY Herbicide (MASTER) Approved 8/17/00.

