



000264-00678-092900

Printed: 07:15:49 Friday, 13 Oct, 2000 # 6 / 11339

Systems Integration Group, Inc.

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:

264-678

Date of Issuance:

SEP 29 2000

Term of Issuance:

Conditional, expires
October 1, 2002

Name of Pesticide Product:

Previcur

NOTICE OF PESTICIDE:
 Registration
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Aventis CropScience
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, NC 27709

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(B) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA sec. 4.
2. Submit and/or cite a livestock feeding study on propamocarb hydrochloride and storage stability data from the livestock feeding study by October 1, 2002. The need for a livestock analytical enforcement method and livestock tolerances will be determined after receipt of the ruminant feeding study and determination of the residues of concern in livestock.
3. Submit and/or cite a corrosion characteristics study.
4. Make the following label change: replace the EPA File Symbol 264-ATI with EPA Registration Number 264-678.

Signature of Approving Official:

Mary J. Walker

Date:

SEP 29 2000

EPA Form 8570-6

CONCURRENCES

SYMBOL	R505C						
SURNAME	Ellwanger						
DATE	9/29/00						

(EPA Reg. No. 264-678, continued)

276

5. Submit one copy of the final printed label before you release the product for shipment.

6. Submit production information (pounds or gallons produced) for this product for the fiscal year in which the use on potatoes is conditionally registered in accordance with FIFRA sec. 29. The fiscal year begins October 1 and ends September 30. The production information will be submitted to the Agency no later than November 15, following the end of the preceding fiscal year. This information must be submitted to:

U.S. Environmental Protection Agency
Registration Support Branch
Registration Division (7505C)
Ariel Rios Building
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec.6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records. If there are any questions concerning this matter contact Dr. Tom Ellwanger at (703) 308-9352.

Sincerely



Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505C)

Enclosure

PREVICUR® Fungicide

For Protection of Potatoes From Late Blight

Active ingredients: Propamocarb hydrochloride, (Propyl [3-(dimethylamino)propyl] carbamate monohydrochloride)*	66.5%
Inert ingredients:	33.5%
*Protected by U.S. Patent No. 4,959,388	Total 100.0%

Contains 6.0 lbs. propamocarb hydrochloride active ingredient per gallon.

EPA Reg No. 264-XXX

EPA Est. No. 407-IA-002

KEEP OUT OF REACH OF CHILDREN CAUTION

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-888-AVENTIS (1-888-283-6847)

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Call a physician or Poison Control Center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow.. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. 	<p>ACCEPTED with COMMENTS In EPA Letter Dated SEP 29 2000</p> <p><i>Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 264-678</i></p>
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. 	
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. 	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p> <p>For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577.</p>		

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling. Avoid contact with skin, eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)


Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.



User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the high water mark. **Do not** contaminate water when disposing of equipment washwaters. **Do not** apply when weather conditions favor drift from the treated areas.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in original container and keep tightly closed. Store in a cool, dry place.

Pesticide disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label directions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container disposal: **Do not** reuse empty container. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements of this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker reentry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

GENERAL INFORMATION

PREVICUR[®] is a soluble fungicide concentrate formulated to give protection against late blight.

PREVICUR[®] is recommended for use in programs which are compatible with principles of Integrated Pest Management (IPM), which include the use of disease-resistant crop varieties, cultural practices, pest scouting, and disease forecasting systems which reduce unnecessary applications of pesticides.

PREVICUR[®] is effective for strategic use in programs that attempt to minimize disease resistance to fungicides. Consult with your Federal or State Cooperative Extension Service representatives for guidance on the proper use of PREVICUR[®] in programs which seek to minimize the occurrence of disease resistance to other fungicides.

APPLICATION TIMING AND DOSAGE


Use PREVICUR[®] in a tank mixture with chlorothalonil or mancozeb as a foliar spray in a **preventive** program for control of late blight in potatoes. Begin applications when conditions are favorable for disease, but before infection, and continue on **7 to 14 day intervals** until the threat of disease is over.

When the risk of disease, or conditions favorable for disease increase, use the shorter spray intervals. Under these conditions and when applying a PREVICUR[®] tank mix at intervals of 10 days or longer, alternate with an application of a contact fungicide midway between PREVICUR[®] tank mix applications. For example, under severe disease conditions, apply a PREVICUR[®] tank mix on a 10-day schedule alternating on the fifth day with a contact fungicide.

Do not apply more than 6.0 pints of PREVICUR[®] (4.5 lb. total ai/A) in a single growing season. Do not apply within 14 days of harvest.

PREVICUR[®] + CHLOROTHALONIL

Apply a tank mix of PREVICUR[®] + chlorothalonil at a rate of 1.2 pints per acre PREVICUR[®] (0.9 lb. ai/A propamocarb) + 0.9 lb. ai/A chlorothalonil (1.2 pints per acre Bravo Weather Stik[®] or equivalent product). This tank mixture may be applied using ground or aerial equipment or via chemigation, following directions for use concerning application timing and volume.



PREVICUR® + MANCOZEB

Apply a tank mix of PREVICUR® + mancozeb at a rate of 1.2 pints per acre PREVICUR® (0.9 lb. ai/A propamocarb) + 1.0 lb. ai/A mancozeb (1.25 lb. per acre DiThane M-45 or equivalent product). This tank mixture may be applied using ground or aerial equipment or via chemigation, following directions for use concerning application timing and volume.

Ground applications: Using ground spray equipment, apply PREVICUR® in a tank mixture with chlorothalonil or mancozeb at the rates specified. PREVICUR® can be used effectively in dilute or concentrate sprays. Thorough uniform coverage is essential for disease control.

Aerial applications: Apply PREVICUR® in a tank mixture with chlorothalonil or mancozeb at the rates specified above using fixed wing or rotary aircraft equipment in a *minimum* of 10 gallons of water per acre to assure uniform coverage of the foliage. Thorough uniform coverage is essential for disease control.

Chemigation applications: See application techniques below. Apply PREVICUR® in a tank mixture with chlorothalonil or mancozeb at the rates specified above in an adequate volume of irrigation water to ensure uniform coverage.

MIXING THE SPRAY

Add one-half of the required amount of water to the spray or mixing tank and start agitation. Add the required quantity of PREVICUR® Fungicide to the water and complete filling with water to the required total volume. Maintain agitation throughout spraying. Do not allow spray mixture to remain in the tank overnight, or for long periods during the day without agitation.

Spray Drift Management

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR.

The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses, or to applications using dry formulations:

1. The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.
3. To avoid spray drift, do not apply when winds exceed 10 mph.

Where States have more stringent regulations, they should be observed. The applicator should be familiar with and take into account the information covered in the *Aerial Spray Drift Reduction Advisory*.

APPLICATION AND CALIBRATION TECHNIQUES FOR SPRINKLER IRRIGATION

Apply this product only through center pivot, motorized lateral move, traveling gun, solid set or portable (wheel move, side roll, end tow, or hand move) irrigation system(s). Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not apply this product through irrigation systems connected to a public water system. 'Public water system' means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days per year.

Controls for both irrigation water and pesticide injection systems must be functionally interlocked, so as to automatically terminate pesticide injection when the irrigation water pump motor stops. A person knowledgeable of the irrigation system and responsible for its operation shall shut the system down and make necessary adjustments should the need arise.

The irrigation water pipeline must be fitted with a functional, automatic, quick-closing check valve to prevent the flow of treated irrigation water back toward the water source. The pipeline must also be fitted with a vacuum relief valve and low pressure drain, located between the irrigation water pump and the check valve, to prevent back-siphoning of treated irrigation water into the water source.

Always inject this product into irrigation water after it discharges from the irrigation pump and after it passes through the check valve. Never inject pesticides into the intake line on the suction side of the pump.

Pesticide injection equipment must be fitted with a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump. Interlock this valve to the power system, so as to prevent fluid from being withdrawn from the chemical supply tank when the irrigation system is either automatically or manually turned off.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur. Do not apply when wind speed favors drift beyond the area intended for treatment.

This product may be used through two basic types of sprinkler irrigation systems as outlined in Sections A and B below. Determine which type of system is in place, then refer to the appropriate directions provided for each type.

A. Center Pivot, Motorized Lateral Move and Traveling Gun Irrigation Equipment

For injection of pesticides, these continuously moving systems must use a positive displacement injection pump, of either diaphragm or piston type, constructed of materials that are compatible with pesticides, capable of being fitted with a system interlock, and capable of injection at pressures approximately 2-3 times those encountered within the irrigation water line. Venturi applicator units cannot be used on these systems.

696

Thoroughly mix recommended amount of this product for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run. Mixture in the chemical supply tank must be continuously agitated during the injection run. Shut off injection equipment after one revolution or run, but continue to operate irrigation system until this product has been cleared from last sprinkler head.

B. Solid Set and Portable (Wheel Move, Side Roll, End Tow, or Hand Move) Irrigation Equipment

With stationary systems, an effectively designed in-line Venturi applicator unit is preferred which is constructed of materials that are compatible with pesticides; however, a positive-displacement pump can also be used. Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a thirty to forty-five minute period. Mix desired amount of this product with water for acreage to be covered so that the total mixture of this product plus water in the injection tank is equal to the quantity of water used during calibration. Operate entire system at normal pressures recommended by the manufacturer of injection equipment used, for amount of time established during calibration. Agitation is recommended. This product can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until this product has been cleared from last sprinkler head.

Use Precautions

- Do not rotate to root and leafy vegetables for 30 days following the last application of PREVICUR® Fungicide.
- Do not rotate to winter wheat and all other crops for 120 days following the last application of PREVICUR® Fungicide.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Aventis CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: AVENTIS CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Aventis CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. AVENTIS CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT AVENTIS CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

NET CONTENTS: 2.64 US.Gallons (10 liters)

Previcur is a registered trademark of the Aventis Group.

Aventis CropScience USA LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

PREVICUR Fungicide (PENDING) Submitted 7/27/00, Resubmitted 8/28/00, Resubmitted 9/20/00

