



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
Aerial Rios Building
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

264-695

Date of Issuance:

OCT 8 2004

Term of Issuance:

Conditional

Name of Pesticide Product:

Reason 500 SC

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Bayer CropScience LP
2 T.W. Alexander Drive
Research Triangle Park, NC 27709

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA). Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA Section 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA Section 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA Section 4.
2. Make the following label change before you release the product for shipment: Revise the EPA Registration Number to read, "EPA Reg. No. 264-695."
3. Submit one copy of the revised final printed label before releasing the product for shipment.
4. If the conditions enumerated above are not complied with, the registration will be subject to cancellation in accordance with FIFRA Section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Signature of Approving Official:

Dennis McNeilly
Acting Product Manager (22)
Fungicide Branch

Date:

OCT 8 2004

GROUP 11 FUNGICIDE**REASON[®] 500 SC Fungicide**

For Use On Potatoes and Other Tuberous and Corn Vegetables, Tomatoes, Onions and Bulb Vegetables, Lettuce, and Cucurbit Crops

ACTIVE INGREDIENT:

Fenamidone: (5S)-3,5-dihydro-5-methyl-2-(methylthio)-5-phenyl-3-(phenylamino)-4H-imidazol-4-one 44.4 %

INERT INGREDIENTS: 55.6 %

Equivalent to 500 g/L or 4.13 lbs. active ingredient per gallon.

EPA Reg. No. 264-XXX

EPA Est. No.:

KEEP OUT OF REACH OF CHILDREN

CAUTION

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

FIRST AID

IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
<p>For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577.</p> <p>Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p>	
<p>NOTE TO PHYSICIAN: Possible mucosal damage may contraindicate the use of gastric lavage. May pose an aspiration pneumonia hazard</p>	

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Harmful if inhaled. Avoid breathing spray mist. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants, shoes plus socks, and chemical resistant gloves made of any waterproof material, such as Nitrile, Butyl, Neoprene and/or Barrier Laminate. Remove contaminated clothing and wash before reuse.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, shrimp and oysters. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Draft and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

SURFACE WATER ADVISORIES

This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: long-sleeved shirt and long pants, shoes plus socks, and chemical resistant gloves made of any waterproof material, such as Nitrile, Butyl, Neoprene and/or Barrier Laminate.

STORAGE AND DISPOSAL

PESTICIDE STORAGE

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

REASON® 500 SC Fungicide is a sprayable, broad-spectrum foliar fungicide for control of certain plant diseases of potatoes and other tuberous and corm vegetables, tomatoes, onions and other bulb vegetables, lettuce (leaf and head), squash, melons, cucumbers and other cucurbit crops. See use directions for list of all crops approved for use. Use of REASON® should be integrated into an overall disease, pest management, or IPM program. REASON® may be used with disease forecasting or Extension advisory programs which recommend application timings based on environmental factors favorable to disease development. Consult with your local agricultural authorities for additional IMP strategies established for your area. The higher rates in the rate range or shorter spray intervals may be required under conditions of heavy infection pressure, highly susceptible varieties, or when disease conducive environmental conditions exist. **FAILURE TO FOLLOW THE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN ILLEGAL RESIDUES, POOR DISEASE CONTROL, AND/OR CROP INJURY.**

Applications may be made at the longer spray intervals under low to moderate disease pressure. When environmental conditions are conducive for disease development, when disease has been detected in the area, or under moderate to high disease pressure, the shorter application interval and the higher rates are recommended.

FUNGICIDE RESISTANCE STATEMENT

REASON® 500 SC Fungicide is an imidazolinone fungicide that exhibits no known cross-resistance to fungicide chemistry such as sterol-inhibitors, dicarboximides, benzimidazoles, anilinopyridines, or phenylamides. REASON® is an inhibitor of the Qo (quinone

outside) site within the electron transport system (QOI inhibitor) in several plant pathogenic fungi species. REASON® exhibits cross-resistance in certain plant-pathogenic fungi to fungicides of the QOI Group, which includes certain strobilurin compounds such as azoxystrobin and trifloxystrobin. Fungal pathogens can develop resistance to products with the same mode of action when used repeatedly. Because resistance development cannot be predicted, use of this product should conform to resistance management strategies established for the crop and use area. Consult your local or State agricultural authorities for resistance management strategies that are complementary to those in this label. Resistance management strategies may include rotating and/or tank-mixing with products having different modes of action or limiting the total number of applications per season. Bayer encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

Do not use a fungicide from the QOI group for more than 1/2 of the total sprays in any one season. Do not use a QOI fungicide for more than 3 sequential applications or more than 6 times per season including mixtures on potatoes and other tuberous and corm vegetables, tomatoes, bulb vegetables, or lettuce. Do not use a QOI fungicide in sequential applications on cucurbit crops. Avoid alternation of REASON® 500 SC Fungicide with other fungicides in the QOI group. REASON® should not be alternated or tank mixed with any fungicide to which resistance has already developed.

HOW TO USE REASON® 500 SC FUNGICIDE

Ground Application

Apply in a minimum of 15 gallons of water per acre. Thorough uniform coverage is essential for effective disease control.

Aerial Application

Apply REASON® using fixed wing or rotary aircraft equipment in a minimum of 5 gallons of water per acre. Thorough and uniform coverage is essential for effective disease control.

Mixing Instructions

Prepare no more spray mixture than is needed for immediate operation. Add □ of the required amount of water to the mix tank. Start the agitator running before adding the required amount of REASON® 500 SC Fungicide. Continue agitation while filling the tank to ensure thorough mixing. Maintain agitation during application and apply with properly calibrated application equipment. Do not allow spray mixture to stand overnight or for prolonged periods. A high quality spreader/sticker, approved for use on growing crops, should be used with REASON®. REASON® should be added to the tank before the addition of any adjuvant. Consult the adjuvant label or manufacturer for crop tolerance and safety information when used with REASON®.

Compatibility

REASON® is compatible with most commonly used fungicide, herbicide, insecticide, and foliar nutrient products. However, the physical compatibility of REASON® with all potential tank mix partners has not been fully investigated. If tank mixing with other pesticides is desirable, conduct a jar test with the volumes and rates typically used in agricultural application. Using a small container of water, add the proportionate amounts of the products, wettable powders and water-dispersible granular products first, then liquid flowables and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 15 minutes. Look for signs of separation, globules, sludge, flakes, or other precipitates. Physical compatibility is indicated if the combination remains mixed or can be remixed readily. THE CROP SAFETY OF ALL POTENTIAL TANK MIXES WITH REASON®, INCLUDING ADDITIVES AND OTHER PESTICIDES HAS NOT BEEN TESTED ON ALL CROPS. BEFORE APPLYING ANY TANK MIXTURE NOT SPECIFICALLY RECOMMENDED ON THIS LABEL, SAFETY TO THE TARGET CROP(S) SHOULD BE CONFIRMED.

POTATOES AND OTHER TUBEROUS AND CORM VEGETABLES

(Arracacha, arrowroot, artichoke (chinese), artichoke (jerusalem), canna (edible), cassava (bitter and sweet), chayote (root), chufa, dasheen, ginger, leren, potato, sweet potato, tanier, turmeric, yam (bean and true)

HOW TO USE	DISEASE	USE RATE	COMMENTS
		FL OZ PER ACRE	
Apply using ground equipment, chemigation, or by air.	Early Blight (<i>Alternaria solani</i>)	5.5 – 8.2	For optimum results, begin applications as soon as crop and/or environmental conditions become favorable for disease development. Applications should be made on a 5 -10 day interval depending upon disease conditions. Do not apply more than 33 fl oz of REASON® (1.07 lb. ai/A) per growing season to potatoes and other tuberous and corm vegetables. Do not apply within 14 days of harvest.
Under conditions favorable for disease development, shorten the spray intervals and/or switch to the higher rate for improved control.	Late Blight (<i>Phytophthora infestans</i>)	1b. ai/A 0.178 – 0.267	

TOMATOES

HOW TO USE	DISEASE	USE RATE	COMMENTS
		FL OZ PER ACRE	
<p>Apply using ground, air, or chemigation equipment.</p> <p>Under conditions favorable for disease development, shorten the spray intervals and/or switch to the higher rate for improved control.</p>	<p>Early Blight (<i>Alternaria solani</i>)</p> <p>Late Blight (<i>Phytophthora infestans</i>)</p>	<p>5.5 - 8.2</p> <p>lb. ai/A</p> <p>0.178 - 0.267</p>	<p>For optimum results, begin applications as soon as crop and/or environmental conditions become favorable for disease development. Applications should be made on a 5 -10 day interval depending upon disease conditions.* Do not apply more than 33 fl oz of REASON® (1.07 lb. ai/A) per growing season to tomatoes. Do not apply within 14 days of tomato harvest.</p>

BULB VEGETABLES

(Onion (green, dry bulb and welsh), garlic, great-headed garlic, leek, and shallot)

HOW TO USE	DISEASE	USE RATE	COMMENTS
		FL OZ PER ACRE	
<p>Apply using ground, air, or chemigation equipment.</p> <p>Under conditions favorable for disease development, utilize a 5 day spray interval for improved control.</p>	<p>Onion Downy Mildew (<i>Peronospora destructor</i>)</p> <p>Purple Blotch (<i>Alternaria pom</i>)</p>	<p>5.5</p> <p>lb. ai/A</p> <p>0.178</p>	<p>For optimum results, begin applications as soon as crop and/or environmental conditions become favorable for disease development. Applications should be made on a 5 -10 day interval depending upon disease conditions. Do not apply more than 33 fl oz of REASON® (1.07 lb. ai/A) per growing season to onions. Do not apply within 7 days of onion harvest. For improved spray coverage, REASON® can be tank mixed with spray adjuvants approved for use on growing crops.</p>

LETTUCE (HEAD AND LEAF)

HOW TO USE	DISEASE	USE RATE	COMMENTS
		FL OZ PER ACRE	
<p>Apply using ground, air, or chemigation equipment.</p> <p>Under conditions favorable for disease development, shorten the spray intervals and/or switch to the higher rate for improved control</p>	<p>Downy mildew (<i>Bremia lactucae</i>)</p>	<p>5.5 - 8.2</p> <p>lb. ai/A</p> <p>0.178 - 0.267</p>	<p>For optimum results, begin applications as soon as crop and/or environmental conditions become favorable for disease development. Applications should be made on a 5 -10 day interval depending upon disease conditions. Do not apply more than 33 fl oz of REASON® (1.07 lb. ai/A) per growing season to lettuce. Do not apply within 2 days of lettuce harvest.</p>

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

OCT 8 2004

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.

264-695

CUCURBIT VEGETABLES:

(Chayote, chinese waxgourd, citron melon, cucumber, gherkin, edible gourds, momordica spp., muskmelon, pumpkin, squash (summer and winter) and watermelon

HOW TO USE	DISEASE	USE RATE	COMMENTS
		FL OZ PER ACRE	
Apply using ground, air, or chemigation equipment. Under conditions favorable for disease development, utilize a 5 day spray interval for improved control.	Downy mildew (<i>Pseudoperonospora cubensis</i>)	5.5	For optimum results, begin applications as soon as crop and/or environmental conditions become favorable for disease development. Applications should be made on a 5 -10 day interval depending upon disease conditions. Do not apply more than 33 fl oz of REASON® (1.07 lb. ai/A) per growing season. Do not apply within 14 days of harvest.
	Alternaria leaf spot (<i>Alternaria cucumerina</i>)	lb. ai/A 0.178	

SPRAY DRIFT

SENSITIVE AREAS: This pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitats for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulation.

1. The distance of the outer most nozzles on the boom must not exceed $\frac{1}{4}$ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed. The applicator should be familiar with and take into account the information covered in the *Aerial Drift Reduction Advisory Information*.

INFORMATION ON DROPLET SIZE: (This section is advisory in nature and does not supersede the mandatory label requirements)

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions below).

CONTROLLING DROPLET SIZE: (This section is advisory in nature and does not supersede the mandatory label requirements)

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM LENGTH: (This section is advisory in nature and does not supersede the mandatory label requirements)

For some use patterns, reducing the effective boom length to less than $\frac{1}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.

APPLICATION HEIGHT: (This section is advisory in nature and does not supersede the mandatory label requirements)

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

SWATH ADJUSTMENT: (This section is advisory in nature and does not supersede the mandatory label requirements)

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.)

WIND: (This section is advisory in nature and does not supersede the mandatory label requirements)

Drift potential is lowest between wind speeds of 2 - 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY: (This section is advisory in nature and does not supersede the mandatory label requirements)

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS: (This section is advisory in nature and does not supersede the mandatory label requirements)

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

DIRECTIONS FOR USE THROUGH SPRINKLER IRRIGATION SYSTEMS

Apply this product only through sprinkler irrigation systems including microjet, solid set, wheel lines and center pivot. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

For specific information about calibration, contact State Extension Service Specialists, equipment manufacturers or other irrigation experts.

SPRAY PREPARATION: Remove scale, pesticide residues, and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

APPLICATION INSTRUCTIONS: First prepare a suspension of REASON® 500 SC Fungicide in a mix tank. Fill tank with 1/2 to 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Add the required amount of REASON® and then the remaining volume of water. Then set sprinkler to deliver no more than 0.4 inch of water per acre. Start sprinkler and uniformly inject the suspension of REASON® into the irrigation water line so as to deliver the desired rate per acre. The suspension of REASON® should be injected with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing. If you should have any other questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

NOTE: When treatment with REASON® has been completed, further field irrigation over the treated area should be avoided for 24 hours to prevent washing the chemical off the crop.

GENERAL PRECAUTIONS FOR APPLICATIONS THROUGH SPRINKLER IRRIGATION SYSTEMS

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension. Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute solution per unit time. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shutdown. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e. g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment. If you are unsure of wind conditions, contact your local extension agent.

Do not apply when wind speed favors drift, when system connection or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled and drained. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from non-uniform distribution of treated water.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation shall shut the system down and make necessary adjustments should the need arise.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.

8/8

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.


LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

NET CONTENTS: 2.5 GALLONS

REASON® is a registered trademark of Bayer.



Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

REASON® 500 SC Fungicide (PENDING) Submitted 3/28/01, Resubmitted 10/1/03.