

264-700

12/14/2009

Jacket
1/9



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number:

264-700

Date of Issuance:

DEC 14 2009

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Name of Pesticide Product:

Dropp SC Cotton Defoliant

Name and Address of Registrant (include ZIP Code):

Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, NC 27709

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA provided that you:

- 1) Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.

Signature of Approving Official:

/s/

Joanne Miller
Senior Registration Specialist
Herbicide Branch, Team 23
Registration Division (7505P)

Date:

DEC 14 2009

Page 2
EPA Reg. 264-700

- 2) Delete "General" from "General Information". On page 3, top of page, change "recommendations" to "directions".

- 3) On page 5, add "to the extent consistent with applicable law" in front of "Bayer CropScience accepts".

A stamped copy of the label is enclosed for your records. You must submit one copy of the final printed label before you release the product for shipment. Products shipped after 12 months from the date of this letter or the next printing of the label whichever occurs first, must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA. Your release for shipment of the product constitutes acceptance of these conditions. This label supersedes all other previously accepted labels. If you have any questions please call Erik Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov.

Dropp[®] SC COTTON DEFOLIANT

For Agricultural Use Only

ACTIVE INGREDIENT: Thidiazuron (N-phenyl-N'-1,2,3-thiadiazol-5-ylurea)*	41.0%
INERT INGREDIENTS:	59.0%
TOTAL 100%	

*Contains 4.0 lbs of thidiazuron per gallon.

EPA Reg No. 264-700

EPA Est. No. 33689-WG-1

KEEP OUT OF REACH OF CHILDREN CAUTION — CAUCION

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • Do not give anything by mouth to an unconscious person.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing spray mist. Avoid contact with skin, eyes, and clothing. Do not contaminate food or feedstuffs.

PERSONAL PROTECTIVE EQUIPMENT (PPE)


All mixers, loaders, and applicators must wear:

- Long-sleeved shirt
- Long pants, and
- Shoes and socks

See Engineering Controls for Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ACCEPTED
with **COMMENTS**
In EPA Letter Dated
DEC 14 2009
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
264-700



ENGINEERING CONTROLS

Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)].

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

This chemical has properties and characteristics associated with chemicals detected in ground water. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

GENERAL INFORMATION

Dropp® SC Cotton Defoliant is used for removal of leaves prior to harvesting. Dropp® disperses easily in water and can be used in any ground and aircraft equipment. It may take several days before the effect of Dropp® becomes noticeable. Adverse conditions, such as low temperatures, may require higher dosages and/or longer times for more complete defoliation. Dropp® inhibits regrowth after defoliation and thus reduces staining of lint during harvesting and ginning.

USE PRECAUTIONS

Do not apply this product through any type of irrigation system.

Rainfall within 24 hours after application will reduce the effectiveness of Dropp®. Use only freshly prepared sprays. Do not store spray mixture overnight.

Do not feed foliage from treated cotton plants or gin trash to livestock.

Do not plant the following crops **earlier than** the specified periods after the application of Dropp® SC:

Small grain, sorghum, corn..... or root crops (except carrots, onions, and sugar beets)	two (2) weeks
Legumes (including alfalfa)..... or leafy vegetables (except lettuce and spinach)	two (2) months
Sugar beets.....	four (4) months
Carrots, onions, or spinach.....	nine (9) months
Lettuce.....	nine (9) months and only following deep plowing (12-15 inches)

Do not use immature crops for food or feed.

Do not allow spray drift to contact crops other than the target crop of mature cotton, or cotton which you desire to defoliate, as this product may injure or defoliate other crops.

Particular care should be taken when applying Dropp® SC adjacent to lettuce, citrus, or cantaloupe. Tank mixes with organophosphates may increase nontarget crop phytotoxicity.

PRACTICES TO LOWER THE POTENTIAL FOR DRIFT ONTO NONTARGET CROPS

During applications, particularly under windy conditions, Dropp® SC may drift to nontarget crops.

To help reduce the drift potential, use the following practices:

- Do not apply Dropp® SC by ground or air when wind speeds exceed ten (10) miles per hour at the time of application. Follow local recommendations if wind speeds of less than ten (10) miles per hour are specified in those recommendations.
- Use of low nozzle pressure (20–30 psi) is recommended.
- Do not apply Dropp® SC when a temperature inversion is present or when conditions favor an inversion prior to completing application(s).
- Additionally, do not apply Dropp® SC by air within one-half (1/2) mile of lettuce. Do not apply Dropp® SC by ground equipment within 100 feet of lettuce.

PRACTICES TO LOWER DRIFT POTENTIAL ONTO NON-TARGET CROPS IN THE RIO GRANDE VALLEY OF TEXAS

To help reduce the drift potential, use the following practices:

- Do not apply Dropp® SC by ground or air when wind speeds exceed ten (10) miles per hour at the time of application. Follow local recommendations if wind speeds of less than ten (10) miles per hour are specified in those recommendations.
- Use of low nozzle pressure (20–30 psi) is recommended.
- Do not apply Dropp® SC when a temperature inversion is present or when conditions favor an inversion prior to completing application(s).
- Use the largest nozzle orifice possible which permits proper deposition and coverage of product.
- In addition, for citrus crops, do not apply Dropp® SC by air when citrus in flush is within five (5) miles downwind of the point of application. Do not apply Dropp® SC by ground when citrus in flush is within one-half (1/2) mile downwind of the point of application.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

TIME OF APPLICATION

Apply Dropp® SC Cotton Defoliant only to mature cotton plants when the last boll you expect to harvest is mature. A boll can be described as "mature" when it is too hard to be dented when squeezed between thumb and fingers, is difficult to slice with a sharp knife, and/or when seeds cut in cross sections have fully developed cotyledons, as evidenced by an absence of jelly within the seed.

Apply Dropp® SC at least 5 days prior to anticipated harvest.

NOTE: Good defoliation with Dropp® is dependent upon three factors:


1. Actively growing cotton
2. High humidity
3. High moisture content in leaf of cotton plant

Use of Dropp® SC alone (without a tank mix partner) when nighttime temperatures are expected to fall below 60°F can result in less than desirable defoliation and/or regrowth inhibition.

Performance of Dropp® SC is variable under low temperatures. Ideally, nighttime temperatures 2–3 days prior to and following application should be above 60°F or total defoliation and regrowth inhibition can be reduced.

Use of Dropp® SC on heat and drought-stressed cotton (low leaf moisture, thick cuticle, etc.) may result in less than satisfactory defoliation and regrowth inhibition.

USE OF ADJUVANTS (CROP OILS)



6/9

The use of adjuvants such as petroleum-based crop oils or penetrating oils approved for use on growing crops has been shown to improve performance during low nighttime temperatures (60°–65°F) or when defoliating drought-stressed cotton. Refer to specific adjuvant label for rate recommendations.

The use of adjuvants is recommended when applying Dropp® SC in the desert Southwest (i.e., Arizona and California).

PRECAUTION: The addition of adjuvants can cause desiccation and/or leaf freezing during periods of high temperatures. The use of compounds that desiccate leaf tissue is not recommended.

MIXING INSTRUCTIONS

Fill the spray tank with one-half of the total amount of water to be used. Begin agitation and add the recommended amount of Dropp® SC. Fill the tank with the additional quantity of water required. Maintain sufficient agitation to ensure a uniform spray mixture during application.

TANK MIX ADJUVANTS

When using tank mixes of Dropp® SC and organophosphates, the use of a surfactant or compatibility agent is recommended to improve tank clean-out and overall defoliation. Prior to mixing in the spray tank, a small scale compatibility test should be conducted.

Use only products which are exempt from tolerance under 40 CFR 180.1001.

Always follow the mixing instructions on the label of the appropriate adjuvant.

APPLICATION

Use the specified dosage of Dropp® SC Cotton Defoliant in sufficient amount of water to give complete coverage of the foliage with uniform wetting of leaf surfaces. Apply in 10–25 gallons of spray per acre by ground equipment and 3–10 gallons per acre by air. Agitate the spray mix during application.

DOSAGE

One gallon of Dropp® SC treats 20 – 80 acres (0.05–0.20 lb. active ingredient/acre).

For cutout and mature cotton under normal weather patterns, use DROPP® SC at 0.05 to 0.10 lb. active ingredient per acre. The 0.05 lb. rate is most effective when used in a tank mix with FINISH® 6 PRO or PREP®.

Increased rates of DROPP® SC above 0.10 lb. of active ingredient per acre may be needed to defoliate and control regrowth during periods of rank growth/high fertilizer conditions, extreme weather conditions, such as extended periods of rain and/or low temperatures (60°–65°F), and on full-season cotton varieties.

Two applications not exceeding a total of 9.6 ounces (0.3 lb. active ingredient/acre) may be necessary to defoliate rank cotton.

DROPP® SC COTTON DEFOLIANT APPLICATION RATE TABLE

To Achieve an Application Rate of:	Use This Amount of DROPP® SC *	At the Indicated Rate, One Gallon of DROPP® SC Will Treat:
0.05 lbs. ai/Acre	1.6 ounces/Acre	80 Acres
0.07 lbs. ai/Acre	2.3 fluid oz/A	56 Acres
0.10 lbs ai/Acre	3.2 ounces/Acre	40 Acres
0.15 lbs ai/Acre	4.8 ounces/Acre	26.6 Acres
0.17 lbs. ai/Acre	5.4 fluid oz/A	23.5 Acres
0.20 lbs ai/Acre	6.4 ounces/Acre	20 Acres

* Maximum of 2 applications not exceeding a total of 9.6 ounces of DROPP® SC (0.3 lb. a.i.) per acre per season

BOTTOM DEFOLIATION FOR USE IN ARIZONA ONLY: Dropp® SC may be used as an aid to improve air circulation and help reduce boll rot caused by moisture.

Apply Dropp® SC at 1.6 ounces (0.05 lb. active ingredient/acre) formulated product per acre.

Application to **bottom** 1/3 of plant **only** recommended.

Timing of application should be based upon the latest date when ground application equipment can enter the field without crop damage.

PRECONDITIONING: Dropp® SC may be used as a pre-conditioner to enhance the activity of a defoliant application.

Apply Dropp® SC at 1.6 – 3.2 ounces (0.05 – 0.1 lb. active ingredient/acre) formulated product per acre.

Timing of application is recommended to be 7–14 days prior to the use of another defoliant. Refer to the second product label prior to use for complete recommendations.

TANK MIX OF DROPP® SC PLUS FINISH® 6 PRO OR PREP™ FOR BOLL OPENING: The tank mix of Dropp® SC plus FINISH® 6 PRO or Prep™ is recommended to improve overall defoliation, and as an aid in accelerating the opening of mature, unopened cotton bolls. Best activity will be obtained where the tank mix is applied to mature cotton plants. (See Time of Application section of this label.) Do not apply tank mix before sufficient unopened bolls have matured to produce the desired cotton yield. If nighttime temperatures are expected to fall below 60°F, unsatisfactory defoliation and regrowth inhibition may result. Certified cotton seed producers should determine boll and seed maturity prior to treatment.

Apply Dropp® SC at a rate of 1.6 – 6.4 ounces (0.05 – 0.2 lb. active ingredient/acre) formulated product plus FINISH® 6 PRO at 1.3 – 2 pt/A (1.0 – 1.5 lbs. active ingredient per acre) or Prep™ at 1.3 – 2.6 pt/A (1.0–2.0 lbs. active ingredient per acre). The higher rate of Dropp® SC should be used where excessive regrowth is anticipated, or during cooler temperatures.

TANK MIX OF DROPP® SC PLUS FINISH® 6 PRO FOR DEFOLIATION ENHANCEMENT: Lower rates of FINISH® 6 PRO may be tank mixed with Dropp® SC to enhance defoliation without providing boll opening activity. FINISH® 6 PRO may still be applied as a sequential treatment for boll opening following an application of Dropp® SC plus FINISH® 6 PRO at a defoliation enhancement rate. Read and follow the FINISH® 6 PRO label, and do not exceed a maximum of 2.0 lb. ethephon ai/acre for combined uses per acre per year.

Apply Dropp® SC at a rate of 1.6 – 6.4 ounces (0.05 – 0.2 lb active ingredient/acre) formulated product plus FINISH® 6 PRO at 5 ounces per acre (0.23 lbs. active ingredient/acre). The higher rates of Dropp® SC should be used where excessive regrowth is anticipated, or during cooler temperatures.

TANK MIX OF DROPP® SC PLUS DEF® 6 OR DROPP® SC PLUS FOLEX® 6 EC: The tank mix of Dropp® SC and Def® 6 or Folex® 6 EC is recommended to improve defoliation and inhibit regrowth under less than ideal conditions. Less than ideal conditions are those when the nighttime temperatures are expected to be 60°F on the date of application and for three days following application.

Best activity will be obtained when the tank mix is applied to mature cotton plants with 60% or more open bolls.

RATE OF APPLICATION

Dropp® SC + Def® 6 = 1.6 – 6.4 ounces Dropp® SC + 0.5 – 2.0 pints Def® 6.

Dropp® SC + Folex® 6 EC = 1.6 – 6.4 ounces Dropp® SC + 0.5 – 2.0 pints Folex® 6 EC.

Higher rate of Dropp® SC should be used where excessive regrowth is anticipated.

Maximum rates of Dropp® and phosphate defoliants applied in a tank mix during very high air temperatures can cause desiccation.

Use lower rate of Def® 6 or Folex® 6 EC under ideal nighttime temperature conditions (65°F); higher rate during less than ideal or adverse conditions.

When applying Def® 6 or Folex® 6 EC defoliant plus Dropp® SC as a tank mix, the following mixing sequence is recommended:

1. Add required water to spray tank, agitate.
2. Add Dropp® SC according to label directions (see **mixing instructions**).
3. Add Def® 6 or Folex® 6 EC according to rate desired **after** the Dropp® SC has completely dispersed.

A second application of the labeled tank mix may be made where necessary, but not exceeding a total of 0.3 lb. active ingredient of Dropp® SC per acre.

Refer to Def® 6 and Folex® 6 EC labeling for additional use directions and cautions when using tank mixtures of Dropp® SC and those products.

DO NOT USE TANK MIX OF DROPP® SC PLUS DEF® 6 OR FOLEX® 6 EC IN THE RIO GRANDE VALLEY COUNTIES OF STARR, HIDALGO, WILLACY, AND CAMERON.

IMPORTANT CLEANOUT INSTRUCTIONS

Tank mixes of Dropp® SC with organophosphates may form a residue in application equipment. **While still fresh and moist**, this residue can be removed by flushing the entire system with a commercial tank cleaner. Consult your State Extension cotton specialist for recommended tank cleaners and cleaning procedures.

Do not allow the spray solution to dry in the application equipment. Immediately following application, clean all equipment (mix tanks, pumps, transfer lines, application tanks, sumps, booms, nozzles and all related equipment) thoroughly with a commercial tank cleaner and water.

The procedure for removing dried residues in the application equipment requires allowing the diluted solution of a commercial tank cleaner to stand in equipment, filled to capacity, for 7 days followed by thorough flushing.

Should small quantities of Dropp® SC remain in inadequately cleaned equipment, they may be released during subsequent applications and may cause damage to crops. Bayer CropScience accepts no liability for damage to crops due to inadequately cleaned equipment.

SEQUENTIAL APPLICATION PRECAUTION

Do not apply sequential application of Dropp® SC following any defoliant or desiccant **except Dropp® SC used alone or Dropp® SC tank mixes**. Reduced activity of the second treatment will result.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Store at temperatures below 100° F. If allowed to freeze, remix before using.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL


[Containers equal to or less than 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in sanitary landfill, or by other procedures approved by State and local authorities.

[Intermediate Bulk Containers – Bottom Discharge (e.g. – Schuetz Caged IBC)]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. End users are authorized to remove tamper evident cables as required to remove the product from the container and to pressure rinse the container with water to remove residual product contents. Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. Empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inch on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

DO NOT REUSE EMPTY CONTAINER



9/9

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: ~~TO THE EXTENT CONSISTENT WITH APPLICABLE LAW~~, BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. ~~TO THE EXTENT CONSISTENT WITH APPLICABLE LAW~~, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: ~~TO THE EXTENT CONSISTENT WITH APPLICABLE LAW~~, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

NET CONTENTS: 1 GALLON


Dropp, Finish and Def are registered trademarks of Bayer.
Prep is a trademark of Bayer.
Folex is a registered trademark of Amvac Chemical.

PRODUCED FOR



Bayer CropScience

Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)



Dropp® SC (PENDING) 03/13/09