



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Mel Toliver
Bayer CropScience
2 TW Alexander Drive
P.O. Box 12014
RTP, NC 27709

JUL 31 2008

Subject: Elite® 45 DF Foliar Fungicide
EPA Reg. No. 264-749
Your amendment dated March 31, 2005 – Adding new uses

Dear Mr. Toliver:

The amended label referred to above, resubmitted on July 8, 2008 in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended is acceptable providing the following changes are made:

1. In the ingredient statement, revise "Inert Ingredients" to read "Other Ingredients".
2. On page 3 of the mixing instructions, revise the sentence to read "Mixing: Add labeled amount of Elite 45 DF. . .".
3. On page 3 of the compatibility instructions, revise the sentence to read "Compatibility: . . . Pour the specified proportions of the products into a suitable. . .".
4. On page 7 of the directions for use, delete the instructions for application to cherries, post-harvest.

One copy of the label stamped "Accepted" is enclosed for your records. Please submit one copy of the final printed label before the product is released for shipment. If you have any questions, please contact Tracy Keigwin by phone at (703) 305-6605 or via email at keigwin.tracy@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Mary L. Waller".

Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505P)

Enclosure

2 8 9

GROUP  FUNGICIDE

Elite® 45 DF Foliar Fungicide

ACTIVE INGREDIENT: Tebuconazole, α -[2-(4-chlorophenyl)ethyl]- α -(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol 45%
 INERT INGREDIENTS: 55%
 100%

EPA Reg. No. 264-749

EPA Est. No.

ACCEPTED
 with **COMMENTS**
 In EPA Letter Dated
 7/31/2008

STOP - Read the label before use
KEEP OUT OF REACH OF CHILDREN

Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under EPA Reg. No. 264-749

WARNING AVISO

usted no entienda la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (TO THE USER: If you do not understand the label, find someone to explain it to you in detail.)

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577
 For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

FIRST AID

If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577.

Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

Note To Physician: No specific antidote. Treat symptomatically.

Symptoms of Poisoning: The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Causes substantial but temporary eye injury. Harmful if swallowed, inhaled or absorbed through the skin. Prolonged or frequently repeated skin contact may cause allergic skin reactions in some individuals. Do not get in eyes or on clothing. Avoid contact with skin. Avoid breathing dust or spray mist.

Personal Protective Equipment (PPE): Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA Chemical Resistant Category Selection Chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

- Protective eyewear

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry interval and notification to workers. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI). The REI for each crop is listed in the application directions associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks
- Protective eyewear

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking or material is spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Dispose of pesticide as directed below. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer CropScience Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer CropScience Emergency Response Telephone No. is 1-800-334-7577, or contact Chemtrec at 1-800-424-9300.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty container into application equipment. Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. If not recycled, then dispose of empty container in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Chemigation: Do not apply this product through any type of irrigation system.

Spray Volume: ELITE 45 DF may be applied in 50 gallons or more of spray solution per acre by ground spray equipment or in 15 gallons or more per acre by aircraft equipment except as noted under "REMARKS" for each crop. Increase spray volume as crop growth increases to ensure thorough coverage of the foliage and fruit. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied.

No aerial application is allowed on Long Island, New York.

Mixing: Add recommended amount of ELITE 45 DF into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the ELITE 45 DF should be thoroughly dispersed prior to the addition of other materials.

Compatibility: To determine the compatibility of ELITE 45 DF with other products, the following procedure should be followed:

Pour the recommended proportions of the products into a suitable container of water, mix thoroughly and allow to stand at least five (5) minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible. For further information, contact your local Bayer CropScience representative.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS; RIVERS; PERMANENT STREAMS, MARSHES OR NATURAL PONDS; ESTUARIES AND COMMERCIAL FISH FARM PONDS.

Spray Drift Management: The interaction of many equipment and weather related factors determines the potential for spray drift. The applicator is responsible for considering all of these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator.

Buffer Zone Requirements: For soil or foliar applications, do not apply by ground within 25 feet, or by air within 150 feet of lakes; reservoirs; rivers; permanent streams, marshes or natural ponds; estuaries and commercial fish farm ponds.

Recommendations For Aerial Applications: The spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Importance of Droplet Size: An important factor influencing drift is droplet size. Small droplets (<150 - 200 microns) drift to a greater extent than large droplets. Within typical equipment specifications, applications should be made to deliver the largest droplet spectrum that provides sufficient control and coverage. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure.

Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Wind Speed Restrictions: Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size, canopy and equipment specifications determine drift potential at any given wind speed. Do not apply when winds are greater than 15 mph and avoid gusty and windless conditions. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Restrictions During Temperature Inversions: Do not make aerial or ground applications during temperature inversions. Drift potential is high during temperature inversions. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however if fog is not present, inversions can also be identified by the movement of smoke from a ground source. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical mixing.

Airblast (Air Assist) Specific Recommendations for Tree Crops and Vineyards: Airblast sprayers carry droplets into the canopy of trees/vines via a radially or laterally directed air stream. The following specific drift management practices should be followed:

- * Adjust deflectors and aiming devices so that spray is only directed into the canopy;

- * Block off upward pointed nozzles when there is no overhanging canopy;
- * Use only enough air volume to penetrate the canopy and provide good coverage;
- * Do not allow the spray to go beyond the edge of the cultivated area (i.e., turn off sprayer when turning at end rows);
- * Only spray inward, toward the orchard or vineyard, for applications to the outside rows.

Runoff Management: Do not cultivate within 10 feet of the aquatic areas to allow growth of a vegetative filter strip. When used on erodible soils, best management practices for minimizing runoff should be employed. Consult your local Soil Conservation Service for recommendations in your use area.

ROTATIONAL CROPS

Any crop not specified on this label may be planted into treated areas 120 days after last application.

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF ELITE 45 DF
Grape	Black rot (<i>Guignardia bidwellii</i>) Powdery mildew (<i>Uncinula necator</i>)	4 oz./A
<p>Powdery Mildew Apply in a preventive spray schedule. Make the first application of ELITE 45 DF before bloom and continue applications using spray intervals of up to 21 days in low to moderate disease pressure. Use a 14-day schedule when disease pressure is severe.</p> <p>Black Rot <u>Preventive Schedule:</u> Make the first application at 1 to 3 inches of new shoot growth and continue at 7- to 14-day intervals through 5° Brix stage or until veraison (berry coloring) is complete. Apply at 1 inch new shoot growth and at 7- to 10-day intervals on highly susceptible varieties or under severe disease conditions. <u>Post-Infection Schedule:</u> A post-infection schedule may be followed from 1-inch new shoot growth through 5° Brix stage. Apply within 72 hours after the beginning of an infection period. ELITE 45 DF applications should not be closer than 7 days apart. Continue ELITE 45 DF applications using the preventive schedule if the post-infection schedule is discontinued.</p> <p>Botrytis ELITE 45 DF, applied in a powdery mildew spray schedule, will enhance the activity of registered fungicides used for control of Botrytis bunch rot. Applications must be made on a 14-day schedule for suppression of Botrytis.</p> <p>GENERAL REMARKS: ELITE 45 DF may be applied up to 14 days before harvest. A maximum of 2 pounds of ELITE 45 DF may be applied per acre per crop season. The minimum interval between applications is 7 days.</p> <p>By ground application only in the state of New York. Apply product in sufficient water for thorough coverage of vines and fruit. Increase spray volume as vine growth increases. Restricted-entry interval (REI) = 12 hours.</p> <p>RESISTANCE MANAGEMENT STRATEGY: ELITE 45 DF is a sterol demethylation inhibitor (DMI) fungicide. It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Tank-mixing ELITE 45 DF with other DMI fungicides is not recommended.</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF ELITE 45 DF
Banana Plantain	Black Leaf Streak (<i>Mycosphaerella fijiensis</i>) Yellow Sigatoka (<i>Mycosphaerella musicola</i>)	3.2 oz./A
<p>Apply specified dosage as a foliar spray in a minimum of 20 gallons or more of water per acre to assure thorough coverage of the plants. A spray oil recommended for use on banana or plantain should be utilized at 4.3 pints per acre in tank-mix with ELITE 45 DF. Thoroughly mix ELITE 45 DF in the spray solution prior to the addition of a spray oil. Make the first application before disease becomes established and repeat applications at 14-day intervals if necessary to maintain disease control.</p> <p>Do not apply ELITE 45 DF to non-bagged bananas. A maximum of 16 ounces of ELITE 45 DF may be applied per acre per harvest cycle. ELITE 45 DF may be applied up to the day of harvest.</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF ELITE 45 DF
Tree Nuts: Almond (see specific use directions) Beechnut Brazil nut Butternut Cashew Chestnut Chinquapin Filbert (see specific use directions) Hickory nut Macadamia nut Pistachios (see specific use directions) Walnut	Brown rot blossom blight (<i>Monilinia laxa</i> , <i>M. fructicola</i>) Botryosphaeria panicle and shoot blight (<i>Botryosphaeria dothidea</i>) Eastern filbert blight (<i>Anisogramma anomala</i>)	8 oz/A
<p>Begin applications when conditions are favorable for disease but before infection. Apply on a 7- to 14-day spray schedule.</p> <p>A maximum of 32 ounces of product may be applied per acre per crop season. Do not cut cover crops in treated areas for feed, or allow livestock to graze treated areas. ELITE 45 DF may be applied up to 35 days before harvest.</p> <p>Restricted-entry interval (REI) = 12 hours.</p> <p>RESISTANCE MANAGEMENT STRATEGY: ELITE 45 DF is a demethylation inhibitor (DMI) fungicide. It may be applied in a tank mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Tankmixing ELITE 45 DF with other DMI fungicides is not recommended.</p>		
Almond	Brown rot blossom blight (<i>Monilinia laxa</i> , <i>M. fructicola</i>)	8 oz/A
<p>Blossom blight: Begin application at pink bud. If the bloom period is extended, and/or severe disease conditions exist, make a second application at full bloom. If conditions remain favorable for disease, make another application at petal fall.</p> <p>GENERAL REMARKS: Apply ELITE in a minimum spray volume of 15 gallons per acre by air or 50 gallons per acre by ground. Reduce the application interval for varieties that are highly susceptible to the indicated diseases or when severe disease conditions exist. The use of ground application after petal fall is preferred because of difficulty in penetrating the canopy and obtaining thorough coverage of the foliage and fruit by air.</p> <p>A maximum of 32 ounces of product may be applied per acre per crop season. Do not cut cover crops in treated areas for feed, or allow livestock to graze treated areas. ELITE 45 DF may be applied up to 35 days before harvest.</p> <p>Restricted-entry interval (REI) = 12 hours.</p> <p>RESISTANCE MANAGEMENT STRATEGY: ELITE 45 DF is a demethylation inhibitor (DMI) fungicide. It may be applied in a tank mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Tankmixing ELITE 45 DF with other DMI fungicides is not recommended.</p>		
Pistachio	Botryosphaeria panicle and shoot blight (<i>Botryosphaeria dothidea</i>)	8 oz/A
<p>Botryosphaeria: Begin application at 25 to 50% bloom and repeat again 10 to 14 days later to protect young flower clusters and fruit. Make two additional applications of ELITE 14 days apart beginning 49 days before harvest.*</p> <p>GENERAL REMARKS: Apply ELITE in a minimum spray volume of 15 gallons per acre by air or 50 gallons per acre by ground. Reduce the application interval for varieties that are highly susceptible to the indicated diseases or when severe disease conditions exist. The use of ground application after flowering is preferred because of difficulty in penetrating the canopy and obtaining thorough coverage of the foliage and fruit by air.</p> <p>A maximum of 32 ounces of product may be applied per acre per crop season. Do not cut cover crops in treated areas for feed, or allow livestock to graze treated areas. ELITE 45 DF may be applied up to 35 days before harvest.</p> <p>Control of insect vectors and a thorough pruning program to remove plant tissue infected by <i>Botryosphaeria</i> are critical for optimum control of this disease.</p> <p>Restricted-entry interval (REI) = 12 hours.</p> <p>RESISTANCE MANAGEMENT STRATEGY: ELITE 45 DF is a demethylation inhibitor (DMI) fungicide. It may be applied in a tank mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Tankmixing ELITE 45 DF with other DMI fungicides is not recommended.</p>		

Filbert (Hazelnut)	Eastern filbert blight (<i>Anisogramma anomala</i>)	8 oz/A
<p>Apply ELITE 45 DF in a preventive spray schedule when conditions are favorable for disease (such as budbreak when the first green tissue is visible). Continue applications of ELITE at 7- to 14-day intervals if weather conditions are conducive to disease development. Use shorter spray intervals during budbreak and rapid shoot elongation. The expanding shoot tip must be protected. Reduce the application interval for varieties that are highly susceptible to the indicated disease or when severe disease conditions exist.</p> <p>Apply ELITE in sufficient spray volume for thorough coverage. Tankmixing the lowest labeled rate of a spray surfactant with ELITE may improve spray coverage and penetration of the active ingredient into plant tissue. The use of ground application is preferred because of the difficulty in penetrating the canopy and obtaining thorough coverage of the foliage and stems by air. A maximum of 32 ounces of ELITE 45 DF may be applied per acre per crop season. Do not cut cover crops in treated areas for feed, or allow livestock to graze treated areas. ELITE 45 DF may be applied up to 35 days before harvest</p> <p>ELITE 45 DF is a sterol demethylation inhibitor (DMI) fungicide. It may be applied in tank mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Tankmixing ELITE with other DMI fungicides is not recommended.</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

APPLICATION DIRECTIONS			
CROP	DISEASE	RATE OF ELITE 45 DF	
Pome Fruits: Apple Crabapple Loquat Mayhaw Pear Pear oriental Quince	Scab (<i>Venturia spp.</i>) Cedar-apple rust (<i>Gymno-sporangium juniperi-virginianae</i>) Powdery mildew (<i>Podosphaera leucotricha</i>)	4 to 8 oz./acre	2 oz./100 gal.**
<p>Scab: Preventive Schedule: Apply at 7- to 10-day intervals from green tip through the second cover spray. Post-Infection Schedule: Apply within 72 hours after the beginning of a scab infection period. ELITE 45 DF applications should not be closer than 7 days apart. Continue ELITE applications using the preventive schedule if the post-infection schedule is discontinued.</p> <p>Cedar-apple rust: Apply at 7- to 10-day intervals from pink stage through the second cover spray.</p> <p>Powdery mildew: Apply at 7- to 10-day intervals from tight cluster through the second cover spray.</p> <p>GENERAL REMARKS: *The amount of ELITE 45 DF required per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees. For smaller trees, multiply 2 ounces times the number of 100 gallons of spray solution required to thoroughly wet, to the point of runoff, one acre of the trees being treated. **For concentrate sprays, apply the same amount of product per acre as would be applied in a dilute spray based on tree size and foliage volume.</p> <p>ELITE may be applied up to 75 days before harvest. A total of 3 pounds of ELITE 45 DF may be applied per acre per crop season. Restricted-entry interval (REI) = 5 days.</p> <p>RESISTANCE MANAGEMENT STRATEGY: ELITE 45 DF is a demethylation inhibitor (DMI) fungicide. It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Do not mix or alternate ELITE 45 DF with other DMI fungicides.</p>			

APPLICATION DIRECTIONS			
CROP	DISEASE	RATE OF ELITE 45 DF	
Stone Fruits: Apricot Cherry Nectarine Peach Plum Plumcot Prune (fresh)	Brown rot (blossom blight, fruit rot) (<i>Monilinia laxa</i> , <i>M. fructicola</i>) Cherry Leaf Spot (<i>Blumeriella jaapii</i>) Powdery mildew (<i>Podosphaera</i> spp., <i>Sphaerotheca pannosa</i>)	4 to 8 oz./acre*	2 oz./100 gal.**
Plum Plumcot Prune (fresh) Peach	Rust (<i>Tranzschelia discolor</i>)	5 to 8 oz./acre*	
<p>Blossom blight: Begin application at white bud on cherry, pink bud on peach and nectarine, and green tip on plums and prunes. Apply again at 50% bloom and at petal fall if conditions continue to be favorable for disease development.</p> <p>Fruit rot: Begin application at two to three weeks before harvest and continue at 7-day intervals through the day of harvest. The blossom and fruit stages must be protected for optimum control of brown rot. If ELITE 45 DF is applied during only one of these stages, another registered fungicide should be applied to the other stage to provide optimum protection. Additional cover sprays during the early post-bloom period are also important for preventing quiescent fruit infections on sweet cherry and peach.</p> <p>Leaf spot: Begin application at petal fall or when first leaves unfold and continue applications at 7- to 14-day intervals. Applications should be made at 7-day intervals early in the growing season when terminal growth is rapid and/or under severe disease conditions. A post-harvest application may be made to maintain control and reduce overwintering inoculum.</p> <p>Powdery mildew: Follow leaf spot schedule until terminal growth ceases.</p> <p>Rust: Begin applications after canker emergence and continue applications at 14-day intervals under severe disease conditions.</p> <p>GENERAL REMARKS: ELITE 45 DF may be applied up to and including the day of harvest (0-day PHI). Do not apply more than 3 lb. of product per acre per crop season. * The amount of ELITE 45 DF required per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees. For smaller trees, multiply 2 ounces times the number of 100 gallons of spray solution required to thoroughly wet, to the point of runoff, one acre of the trees being treated. For concentrate sprays, apply the same amount of product per acre as would be applied in a dilute spray based on tree size and foliage volume, but not less than 4 ounces of ELITE 45 DF per acre. Apply 6 to 8 ounces per acre of ELITE 45 DF when severe disease conditions exist. ** When using the 2 oz./100 gal. rate, use at least 4 ounces of ELITE 45 DF per acre.</p> <p>Restricted-entry interval (REI) = 5 days.</p> <p>RESISTANCE MANAGEMENT STRATEGY: ELITE 45 DF is a demethylation inhibitor (DMI) fungicide. It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Do not mix ELITE 45 DF with other DMI fungicides.</p>			
Cherry (Post-Harvest)	Brown rot (<i>Monilinia</i> spp.) Gray mold (<i>Botrytis cinerea</i>) Rhizopus rot (<i>Rhizopus</i> spp.)	8 to 16 oz. per 25,000 lb. of fruit	
<p>Mix the specified rate of ELITE 45 DF in 100 gallons of water or wax/oil emulsion (or aqueous dilution of a wax/oil emulsion). Apply through a spray or flood system on a commercial packing line after stem cutting, sizing, sorting, or washing. For spray treatments, use conventional spray equipment such as T-Jet, flat-fan, or flood jet spray nozzles. Use enough fruit coating material to fully coat fruit. Workers handling treated fruit need to wear waterproof gloves. Do not apply more than once prior to boxing and do not rinse.</p>			
Plums (Post-Harvest) Including Chickasaw, Damson, Japanese, Plumcot, Prune (Fresh)	Brown rot (<i>Monilinia</i> spp.) Grey Mold (<i>Botrytis cinerea</i>) Rhizopus rot (<i>Rhizopus</i> spp.)	8.0 oz. per 200,000 lb. of fruit	
<p>Mix specified rate of ELITE 45 DF in 100 gallons of water, with or without wax. Maintain ELITE 45 DF suspension by direct or by-pass agitation sufficient to prevent settling of product in spray mixture. Apply as spray to fruit as they travel along packing line. After sizing, sorting or washing, ELITE 45 DF application may be incorporated into a waxing system using conventional equipment that provides for even and complete coverage of fruit. Do not apply more than once prior to boxing and do not rinse.</p>			

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

Net Contents:

PRODUCED FOR



Bayer CropScience

**Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)**

Elite 45 DF Foliar Fungicide (PENDING) Submitted 07/14/04, Resubmitted 08/21/06, Resubmitted 07/08/08