

Bayer CropScience LP P.O. Box 12014 2 T.W. Alexander Drive Research Triangle Park, North Carolina 27709 1-866-99BAYER (1-866-992-2937) <http://www.bayercropscienceus.com>

Provado® 1.6 Flowable
Insecticide EPA Reg. No. 264-763 Label Revisions: Added use directions for Bush Berries to include Blueberry (including lowbush and highbush).

Supplemental Label

Provado® 1.6 Flowable Insecticide

For control of certain insects infesting various crops.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read this label and the product package label before using this product. This Supplemental Label must be in possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the registered product label for Provado® 1.6 Flowable Insecticide.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes,

Applicators and Other Handlers Must Wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.
- Shoes plus socks

Follow manufacturer's instructions for cleaning/ maintaining personal protective equipment, PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RECOMMENDED APPLICATIONS

BUSH BERRIES

CROP	PEST	RATE PER APPLICATION
Blueberry (including highbush and lowbush) Currant Elderberry Gooseberry	Aphids Leafhoppers	Foliar Application 3 - 4 fl oz per acre
(including Glassywinged sharpshooters)		
Huckleberry Juneberry Lingonberry Salal	Thrips Japanese beetle (adults)	6 - 8 fl oz per acre

FOLIAR APPLICATION Do not apply prior to bloom or during bloom. Aphid Control and Japanese beetle (adult) control - Apply specified dosage of Provado® in one of the following methods: • As a ground application in a minimum of 20 gal of water per acre. • As an aerial application in a minimum of 5 gal of water per acre. Applications must achieve thorough coverage for optimum control. Initially, Provado® will provide lethal activity against Japanese beetle adults. As Provado® is absorbed into the foliage, lethal activity will diminish but sub-lethal effects, including "knockdown", will persist for 7-10 days. Under conditions of heavy beetle pressure, re-infestation, or adverse environmental conditions, reapplication of Provado® 1.6 may be needed. Allow at least 7 days between Provado® treatments. No more than 5 applications of Provado® may be made per season. Allow 3 days between last application and harvest.

NOTE: Regardless of formulation or application type (foliar or soil), do not apply more than a total of 0.5 lb active ingredient per acre Provado® or Admire® or a combination of Provado® or Admire® per season. Do not apply more than a total of 40 fl oz/acre (0.5 lb ai/acre) per season of Provado® (foliar application).

NOTE: Regardless of formulation or type of application (soil or foliar), do not apply more than a total of

ACCEPTED

MAY 17 2004

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide Registered under EPA Reg. No. 264-763

2/2

0.5 lb active ingredient of Admire® or Provado® per acre per season.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at www.bayercropscienceus.com

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).