

**UNITED STATES ENVIRONMENTAL PROTECTION AGENCY**

WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

OCT 24 2007

Mr. Mike See  
Bayer CropScience  
P.O. Box 12014  
2 T.W. Alexander Drive  
Research Triangle Park, NC 27709

Subject: Axiom DF Herbicide  
EPA Registration No. 264-766  
Supplemental labeling submitted September 12, 2007 for uses on wheat and  
grasses grown for seed

Dear Mr. See:


The supplemental labeling referred to above is acceptable provided that you adhere to the following conditions:

1. Make all of the changes specified in the document "Summary of Comments on Reg # 264-766".
2. Submit no later than December 30, 2009, a special comparative sensitivity study on thyroid hormone levels in neonatal and adult rats. The registrant should consult with the Agency in developing the protocol for the study.
3. Submit a 28-day inhalation toxicity study in rats no later than January 31, 2008.
4. Submit a standard for the FOE thioglycolate sulfoxide metabolite to the National Pesticide Standards Repository/Analytical Chemistry Branch/OPP, 701 Mapes Road, Fort George G. Meade, MD 20755-5350.
5. Submit two copies of final printed labeling within 30 days of receipt of this letter.

A stamped copy of the label is enclosed for your records.

If you have any questions about this letter, you may call Tobi Colvin-Snyder at 703-305-7801.

Sincerely,

A handwritten signature in black ink, appearing to read "Jim Tompkins", is written over the typed name.

Jim Tompkins  
Product Manager 25  
Herbicide Branch  
Registration Division (7505P)

# Summary of Comments on untitled

---

Page: 1

---

Author: tsnyder  
Subject: Highlight  
Date: 10/11/2007 11:30:40 AM

**T** Move this text to immediately following "DIRECTIONS FOR USE".

---

Author: tsnyder  
Subject: Inserted Text  
Date: 10/11/2007 11:29:37 AM

**T** package

---

Author: tsnyder  
Subject: Replacement Text  
Date: 10/11/2007 11:29:56 AM

**T** Axiom DF

---

Author: tsnyder  
Subject: Cross-Out  
Date: 10/11/2007 11:31:44 AM

**T** Replace with text highlighted above. Stating this twice is redundant.

---

4/9

ACCEPTED  
with COMMENTS  
In EPA Letter Dated  
OCT 24 2007  
under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
amended, for the pesticide  
registered under EPA Reg. No.  
264-766



**Bayer CropScience LP**  
P.O. Box 12014  
2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
1-866-99BAYER (1-866-992-2937)  
<http://www.bayercropscienceus.com>

**Axiom® DF Herbicide**  
**EPA Reg. No.: 264-766**  
**EPA Approval: May 10, 2007**  
**For use on winter wheat and grasses grown for seed**

### Supplemental Label

# Axiom® DF Herbicide

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read this label and the product package label before using this product. This Supplemental Label must be in possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the registered product label for Rely® Herbicide.

#### DIRECTIONS FOR USE

~~It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Please refer to the Rely® Herbicide product label for complete directions for use.~~

#### GENERAL USE RESTRICTIONS

Axiom® DF Herbicide is for use on soybeans, field corn, white corn, corn grown for silage, perennial grasses grown for seed, wheat and triticale and field corn grown for seed. Do not apply to popcorn or sweetcorn.

Do not apply this product through any type of irrigation system. Do not apply aerially.

Do not use flood irrigation to apply, activate or incorporate Axiom®.

Do not apply more than 23 ounces Axiom® per acre per use season in corn, 13 ounces per acre per use season in soybeans or perennial grasses grown for seed, and 10 ounces per acre per use season in winter wheat.

For all metribuzin products used, do not apply more than a total of 1 lb ai/A per use season of metribuzin.

Do not apply Axiom® or Axiom® tank mixtures when environmental conditions may favor drift to non-target sites.

Corn seed should be planted a minimum of 1-1/2 inches deep and soybean seed should be planted a minimum of 1 inch deep.

If any crop treated with Axiom® is lost, corn or soybeans may be replanted immediately. Do not make a second application of Axiom®.

This product may not be mixed or loaded within 50 feet of any wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing / loading areas.

Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or washwater, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specific minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing / loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

This product must be used in a manner which will prevent back siphoning in wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

#### RATE SELECTION / SOIL TEXTURE

The specified rates of Axiom® DF Herbicide are defined by texture and organic matter content of the soil being treated. Unless a specific soil texture is mentioned, rate tables throughout this label refer to the following three soil texture groups: coarse, medium and fine. If you are not sure how to classify your soil, contact your Bayer CropScience representative, the Cooperative Extension Service or other knowledgeable person. The following chart includes a complete listing of soil textures included in each of the soil textures groupings:

COARSE	MEDIUM	FINE
Sand	Loam	Silty clay loam
Loamy sand	Silt loam	Silty clay
Sandy loam	Silt	Clay loam
	Sandy clay loam	Clay
	Sandy clay	

**MIXING INSTRUCTIONS**

**Liquid Carriers:**

Axiom® DF Herbicide is a dry flowable herbicide that must be mixed in water or sprayable fluid fertilizer. Compatibility of Axiom® or its labeled tank mix products with these liquid carriers should always be predetermined prior to spraying. Refer to the SPRAYABLE FLUID FERTILIZER COMPATIBILITY TEST (Appendix I) of this label to determine product compatibility in fluid fertilizer carriers.

Before mixing Axiom® and its labeled tank mixtures, examine the spray equipment making sure it is completely clean and free of rust or corrosion. Be sure the equipment is free of any residues from previously used pesticides. Flush lines with clean water or recommended detergents after the last application. Use an approved method for disposing of rinsate.

For optimal spray tank mixing convenience and efficiency, Axiom® is recommended to be added to the spray tank via an eductor system.

The proper mixing sequence for Axiom® and recommended tank mixtures with the appropriate liquid carrier is as follows:

1. Fill the spray tank or nurse tank 1/4 full with the appropriate liquid carrier.
2. Start recirculation and agitation system and continue throughout mixing and application.
3. If the compatibility test indicates the need of a compatibility agent, add the recommended amount of compatibility agent to the spray tank.
4. If ammonium sulfate is to be used, add it now.
5. Next add the specified quantity of Axiom® through the eductor system or to the spray tank [slowly add Axiom® if water or sprayable grade nitrogen fertilizers (28-0-0, 32-0-0) are the carriers; for other sprayable grade fertilizers first check compatibility and then either mix directly or preslurry first in water depending on the results of the compatibility test].
6. If tank mixing with wettable powders or other dry flowable products in water as the carrier, they may be added now. If tank mixing these products in a sprayable grade fertilizer carrier, first preslurry these products with water and then slowly add them to any sprayable grade fertilizer carrier.
7. If tank mixing with emulsifiable concentrates or soluble products, add the products to the spray tank.
8. If tank mixing with a glyphosate- containing product or Touchdown®, add the products to the spray tank.
9. If mixing spray adjuvants in the mixture, add them after all other products have been mixed.
10. Fill the spray tank to the desired level with the appropriate liquid carrier.
11. Continue agitation during transport and application until the spray tank is empty.

Axiom® and all registered mixtures should be kept agitated once mixed and then sprayed immediately. Do not allow mixtures to stand for prolonged periods of time. Water quality, pH, temperature and/or other components of the mixture may affect how long the mixture may stand before application.

**Dry Fertilizer Carriers:**

Axiom® may be impregnated or coated on dry bulk fertilizers for application. Refer to Appendix II of this label for details.

**APPLICATION INFORMATION**

**SPRAYER APPLICATION**

**Ground Broadcast Treatment:** Accurately calibrate the sprayer prior to mixing the herbicide treatments. Apply Axiom® and the labeled tank mixtures in a minimum of 10 gallons of total spray volume per acre using broadcast boom equipment. Use a pump with capacity to maintain 30 to 40 psi at the nozzles, maintain adequate in-tank agitation to keep the spray mixture in suspension and provide a minimum of 20% bypass at all times. Application must be made at a sufficient spray pressure and volume to provide accurate and uniform application of spray particles to a given area without causing spray drift to non-target areas. If mixed with other labeled herbicides, the spray volume may be no less than the minimum volume recommended by the tank mix product or 10 gallons, whichever is greater. The use of screens to protect the pump and nozzles is recommended. Screens placed on the suction side of the pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles. Refer to the nozzle manufacturer for additional recommendations. Agitate thoroughly before and during application with either bypass or mechanical agitation. Rinse the sprayer thoroughly with clean water immediately after each use.

**Band Treatment:** Axiom® and the labeled tank mixtures may be applied as a band treatment. Use the following formula to calculate the amount of herbicide needed for band treatments:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \frac{\text{Broadcast rate}}{\text{per acre}} = \frac{\text{Band rate}}{\text{per acre}}$$

**DRY FERTILIZER APPLICATION**

Apply using a minimum of 200 lb of dry fertilizer per acre. See Appendix II for additional details.

**APPLICATION METHODS AND TIMING**

Axiom® may be applied either alone or in tank mixtures with or sequentially with certain other registered and recommended herbicides provided that, for tank mixes, the labeling for the other herbicide does not prohibit such mixing. Applications may be made preplant surface, preplant incorporated or preemergence. Axiom® may be used in either a single or split application program. Do not apply when environmental conditions favor drift.

**Preplant Surface:** For use in conservation, minimum or no-tillage crop production systems, Axiom® alone or as a recommended tank mixture may be applied as a broadcast spray up to 45 days before planting in corn and 14 days before planting in soybeans. A sequential application is recommended for applications made 30 to 45 days prior to planting, where 2/3 of the highest specified broadcast rate for the crop and soil texture is applied initially and the remaining 1/3 is applied at planting. Treatments made less than 30 days before planting may use either a single or sequential application. If weeds are present at the time of application, apply the Axiom® treatment in tank mixture with a recommended nonselective herbicide such as a glyphosate-containing product or Touchdown. If possible, do not move treated soil out of the row or move untreated soil to the soil surface during planting, as weed control may be reduced.

**Preplant Incorporation:** Apply Axiom® alone or in combination with recommended tank mixes as a broadcast spray and incorporate into the upper 1 to 2 inches of the soil surface up to 14 days before planting. Avoid deep incorporation since reduced weed control and/or crop injury may result. Incorporate with implements which provide uniform, shallow incorporation (example -finishing disk, harrow, rolling cultivator, etc.)

**Pre-emergence:** Axiom® DF Herbicide alone and its recommended tank mixes may be applied to the soil surface as a broadcast spray or band application after planting of the crop but prior to weed or crop emergence. If weeds are present at the time of application, apply Axiom® with a nonselective herbicide such as a glyphosate-containing product or Touchdown. Rainfall and/or overhead sprinkler irrigation is necessary to move Axiom® into the upper soil surface where weed seeds germinate. Dry weather conditions following application may reduce weed control. If adequate moisture is not received within 7 to 10 days after applications and weeds begin to emerge from the soil, a light rotary hoeing or shallow incorporation (no deeper than 1/2 inch deep) will improve performance and minimize crop damage. Excessive rainfall or irrigation after application may reduce weed control.

**Special Applications: Fall Application** (for use only in IA, MN, ND, SD, WI, north of Route 20 in NE, north of Route 136 in IL, and north of Interstate 70 in OH):

Following harvest of crops in the fall, Axiom® DF Herbicide may be applied to crop stubble after October 15, when the sustained soil temperature at the four-inch soil depth is less than 50 °F, but before the ground is frozen. This application is limited to only medium- and fine-textured soils with an organic matter of 2.5% or greater and which will be planted to corn the following spring. The soil may be tilled before or after application with incorporation depth no more than two to three inches following herbicide application. If a spring application of Axiom® follows the fall application, the total Axiom® rate for both applications must not exceed 23 ounces.

**WINTER WHEAT**

Axiom Herbicide is recommended for control of certain annual grass and broadleaf weeds when applied early postemergence in winter wheat. Axiom may be applied either alone, in tankmix combination with and/or sequentially with other registered herbicides.

**SPECIAL PRECAUTIONS AND RESTRICTIONS**

1. Apply by ground application only.
2. Certain winter wheat varieties are sensitive to Axiom. Use of Axiom is not recommended on the following sensitive winter wheat varieties: AgriPro Clemens, AgriPro Foster, AgriPro Mallard, AgriPro Mitchell, Arise 766, Arise W33, AT 90W, AT 91W, Arapaho, Baker Seed 33, Beck 101, Beck 102, Beck 108, Brodbeck Celebrity, Brodbeck Stealth, Brodbeck Viper, Century, Cimarron, Coker 833, Coker 916, Coker 983, Coker 9024, Coker 9105, Coker 9323, Coker 9474, Coker 9663, Coker 9835, Coker 9704, Coker 9766, Coker 9835, Coker 9877, CountryMark 522W, CountryMark 558W, Delta Grow 3300, Delta King 7777, Delta King 9121, EK 102, Erwin Keith 102, Erwin Keith 114, Exsegen Rebekah, FFR 552, FFR 535, FFR 555, FFR 558, Florida 304, Freedom, FS 417, FS 423, FS 425, FS 430, GA Dozier, GA Fleming, GA Roberts, GA Stuckey, Gateway Daybreak, Glory, Gore, Hazen, HBK 9650W, Hickory, Jackson, Jay Pee, Julie III, KAS Hamilton, KAS Independence, KY-49-25, Linden, Madison, Mesa, MO Sisson, Mustang, MWS Seeds 30, NE Arapahoe, NE Wahoo, OH Honey, Pacer, Patriot, Patriot 911, Patriot 942, Patriot 948, PI XW 522, PI 2551, PI 2163, Pioneer 2643, Pioneer 2691, Princeton 733, Progeny 156, PSR W71, PSR 226, PSR 278, Rosen, Rupp Seed RS 931, Savannah, Schlessman GR 1620, Seed Consultants 1358, Sierra, Southern States 522, Southern States 535, Southern States 550, Southern States 560, Steyer Seeds Podach, TAM 107, Terra SR204, TR 101, TR 211, TR 1011, TR 8822, TR TV 8825, TR TV 8450, TR TV 8557, TR TV 8565, Triumph 64, Vona, Vigoro RC901, USG 3209, TSG 3709, University of Arkansas Sabbe, Wakefield, Wings, Winridge, Yamhill.  
Not all available winter wheat varieties and hybrids may have been tested for tolerance. If a variety or hybrid is not listed (especially newly released varieties), treat only a small area until tolerance is established before treating large acreages.

3. Weed control may be reduced if excessive straw from the previous crop is present at AXIOM application.
4. Do not apply more than 10 ounces of AXIOM per acre per use season.
5. Do not apply in tank mixtures with crop oil concentrates (COC), vegetable and/or petroleum oils as crop injury may result.
6. Do not graze winter wheat within 30 days of AXIOM application.
7. Do not apply this product by air.

**WEED SPECIES CONTROLLED BY AXIOM**

AXIOM herbicide applied early postemergence at specified rates will control many important annual grass and broadleaf weeds:

POSTEMERGENCE BURNDOWN WEED CONTROL BY AXIOM	
ANNUAL GRASSES	ANNUAL BROADLEAF WEEDS
Bluegrass, annual ( <i>Poa annua</i> ) Brome, soft ( <i>Bromus mollis</i> ) Bromegrass, California ( <i>Bromus carinatus</i> ) Canarygrass, hood ( <i>Phalaris paradoxa</i> ) Fescue, rattail ( <i>Vulpia myuros</i> )	Chickweed, mouseear ( <i>Cerastium vulgatum</i> ) Falseflax, smallseed ( <i>Camelina microcarpa</i> ) Fiddleneck, coast ( <i>Amsinckia intermedia</i> ) Fiddleneck, tarweed ( <i>Amsinckia lycopsoides</i> ) Henbit ( <i>Lamium amplexicaule</i> ) Mustard, blue ( <i>Chorispora tenella</i> ) Radish, wild ( <i>Raphanus raphanistrum</i> )

**WEED SPECIES SUPPRESSED BY AXIOM**

AXIOM herbicide applied early postemergence at specified rates will suppress many important annual grass and broadleaf weeds:

POSTEMERGENCE BURNDOWN WEED SUPPRESSION BY AXIOM	
ANNUAL GRASSES	ANNUAL BROADLEAF WEEDS
Barley, foxtail ( <i>Hordeum jubatum</i> ) Blackgrass ( <i>Alopecurus myosuroides</i> ) Brome, downy ( <i>Bromus tectorum</i> ) Brome, ripgut ( <i>Bromus rigidus</i> ) Crabgrass, Large ( <i>Digitaria sanguinalis</i> ) Foxtail, spp. ( <i>Setaria spp.</i> ) Oats, wild ( <i>Avena fatua</i> ) Ryegrass, annual ( <i>Lolium multiflorum</i> ) Windgrass ( <i>Apera spica-venti</i> )	Chickweed, common ( <i>Stellaria media</i> ) Lambsquarters, common ( <i>Chenopodium album</i> ) Mustard, tumble ( <i>Sisymbrium altissimum</i> ) Pigweed spp. ( <i>Amaranthus spp.</i> ) Shepherdspurse ( <i>Capsella bursa-pastoris</i> )
Weed suppression indicates that weed control may vary from poor to excellent depending on soil characteristics, stage of growth at application and environmental conditions at and following application. For maximum control, apply at the highest specified rate. Weed suppression appears as a reduction in weed size and growth as compared to a non-treated area in the same field.	

**AXIOM USE RATES**

AXIOM use rates and application timings are listed in the following table:

AXIOM IN WINTER WHEAT		
AXIOM DF Broadcast Rates Per Acre		APPLICATION TIMINGS AND REMARKS
Soil Texture	Rate (Ounces per acre)	
Coarse	4 – 6	Apply specified rate per acre as a uniform broadcast spray using 5 or more gallons of spray carrier per acre as a ground application. Application timings are from spike to the 3 leaf growth stage of winter wheat (Early Postemergence). For best control of weeds, 1) plant wheat after the first flush of weeds has emerged and was destroyed by tillage or in no-till winter wheat, apply a nonselective herbicide to control emerged weeds at or prior to planting 2) make AXIOM applications prior to the 1-leaf stage of the weeds and 3) use the highest specified rate.
Medium / Fine	6 – 10	

**AXIOM and SENCOR IN WINTER WHEAT**

AXIOM contains metribuzin which is the active ingredient in SENCOR. AXIOM may be tank-mixed with SENCOR but do not exceed a total of more than the equivalent amount of metribuzin active ingredient specified on the SENCOR label for the soil type and crop stage at application.

For sequential applications of AXIOM and SENCOR, do not exceed a total of 0.5 pounds active ingredient (or the equivalent of 10 2/3 oz SENCOR 75 DF product) per acre per year of metribuzin. Allow a minimum of 21 days between sequential applications of AXIOM and SENCOR if wheat is actively growing, or 45 days between applications if wheat has entered dormancy, or is stressed by frost, disease, insects, drought, excessive moisture, or any other factor.

Refer to the individual product labels for detailed explanations on use rates, directions and special precautions/restrictions.

**AXIOM USE IN TANK MIXTURES AND/OR SEQUENTIAL APPLICATIONS**

If a broader spectrum of weed control is desired, AXIOM may be tank mixed with and/or applied sequentially with any other herbicides which are registered for use in winter wheat unless specifically prohibited by the label of the tank mix or sequentially-applied product. Three-way or multiple tank mixtures are permitted unless restricted by the product label. Use of crop oil concentrates, vegetable oils and/or petroleum oils are not recommended as crop injury may occur. Refer to the individual product labels for specified use rates, precautions and/or restrictions. Always follow the most restrictive label.

<b>AXIOM® DF HERBICIDE SEQUENTIAL APPLICATION:</b>	
<b>SEQUENTIAL APPLICATION (Labeled Use Rate)</b>	<b>ADDITIONAL WEEDS CONTROLLED</b>
Osprey™ Herbicide (3.2 – 4.75 oz/A)	Annual/Italian ryegrass, wild oats, blackgrass, roughstalk bluegrass, littleseed canarygrass, Persian darnel, wild mustard, volunteer canola
Postemergence application only. Requires an adjuvant as specified in the product label Applications should be made to actively growing weeds Do not make more than one application in one winter wheat growing season Do not apply more than 4.75 oz per acre in one winter wheat growing season	

**PERENNIAL GRASSES GROWN FOR SEED ONLY**

**(Washington, Oregon [except for Jefferson Co.] and Idaho only)**

AXIOM may be applied to established perennial grasses grown for seed for the control/suppression of weeds and volunteer grasses. Recommended perennial grasses for AXIOM use include established stands of tall fescue, fine fescue, perennial ryegrass, orchardgrass, and bluegrass.

**Special Precautions and Restrictions:**

1. Apply by ground application only.
2. Do not apply this product through any type of irrigation equipment.
3. Weed control may be reduced if excessive straw from the previous crop is present at AXIOM application. If excessive straw is present, uniformly spread the straw across the field, and/or remove/burn the straw prior to application.
4. Use only on fields which are at least one year old or have had at least one seed harvest.
5. Use on perennial ryegrass after the second harvest may result in crop injury. Use the rates of 8 to 10 oz/acre to minimize injury.
6. Do not use on soils with less than 1.0% organic matter, or on soils with pH greater than 7.5.
7. Do not use on clay knobs or poorly covered subsoils.
8. Do not tank mix AXIOM with Diuron, Karmex or other photosynthetic inhibitor herbicides. Allow a minimum of four weeks between application of AXIOM and photosynthetic inhibitor herbicides.
9. AXIOM may be tank-mixed with oxyflourfen, however sequential applications of AXIOM and oxyflourfen must be separated by a minimum of four weeks.
10. Allow at least 120 days between application and harvest of seed grass.
11. Do not allow animals to graze treated fields for a minimum of 30 days following application.
12. Do not make more than one application per use season.
13. Do not apply more than 13 oz AXIOM per acre per use season.

**GRASSES CONTROLLED OR PARTIALLY CONTROLLED BY AXIOM**

AXIOM applied at specified use rates will provide control/suppression of the following weeds:

GRASSES CONTROLLED/SUPPRESSED*	
Barley, foxtail* ( <i>Hordeum jubatum</i> )	Canarygrass, hood ( <i>Phalaris paradoxa</i> )
Bluegrass, annual ( <i>Poa annua</i> )	Crabgrass, Large* ( <i>Digitaria sanguinalis</i> )
Brome, downy* ( <i>Bromus tectorum</i> )	Fescue, rattail ( <i>Vulpia myuros</i> )
Brome, ripgut* ( <i>Bromus rigidus</i> )	Foxtail spp.* ( <i>Setaria spp.</i> )
Brome, soft ( <i>Bromus mollis</i> )	Oats, wild* ( <i>Avena fatua</i> )
Bromegrass, California ( <i>Bromus carinatus</i> )	Ryegrass, annual ( <i>Lolium multiflorum</i> )
Annual Volunteers of the established crop	
Bluegrass ( <i>Poa spp.</i> )	Orchardgrass ( <i>Dactylis glomerata</i> )
Fescue, tall ( <i>Festuca arundinacea</i> )	Ryegrass, perennial ( <i>Lolium perenne</i> )
Fescue, fine ( <i>Festuca tenuifolia</i> )	
* Suppressed weeds may be stunted in growth and/or be reduced populations as compared to non-treated areas but control will generally not be commercially acceptable.	

**CORN, SOYBEAN AND WINTER WHEAT CROP ROTATION INSTRUCTIONS**

Waiting period after an AXIOM application for planting of the following crops in rotation. In the event of a crop failure after application of Axiom®, only the following crops can be planted, after the waiting period as listed below. Do not make a second application of Axiom®.

Immediately			
Corn	Soybean	Wheat <sup>1</sup>	Triticale <sup>1</sup>
1 month			
Potato			
4 months			
Carrot	Wheat <sup>2</sup>	Triticale <sup>2</sup>	
8 months			
Cotton			
12 months			
Alfalfa	Clover	Rice	Wild Rice
Barley	Fescue	Rye	Other crops not listed (except root crops)
Bermudagrass	Millet, pearl	Sorghum	
Bluegrass	Millet, prose	Teosinte	
Bromegrass	Oats		
Buckwheat	Popcorn		
18 months			
Onions	Sugar beets	Other root crops not listed	
<sup>1</sup> Where rates of 10 oz/A or less have been applied.			
<sup>2</sup> Where any other labeled rate of AXIOM >10 oz/A has been applied.			

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577  
 For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at [www.bayercropscienceus.com](http://www.bayercropscienceus.com)

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).