

RESTRICTED USE PESTICIDE

Due to Toxicity to Fish and Aquatic Organisms

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

GROUP	3	INSECTICIDE
GROUP	4A	INSECTICIDE

Leverage[®] 2.7

Suspension Emulsion Insecticide

For control of insect pests on conventional and BT cotton and other crops.

ACTIVE INGREDIENT:

Imidacloprid, 1-[(6-chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine.....	17.0%
Cyfluthrin, Cyano(4-fluoro-3-phenoxyphenyl)methyl 3-(2,2-dichloroethenyl)-2,2-dimethylcyclo-propanecarboxylate.....	12.0%

INERT INGREDIENTS:..... 71.0%

Contains 1.6 lb Imidacloprid per gallon plus 1.1 lb Cyfluthrin per gallon 100.0%

(This product contains aromatic petroleum distillates.)

EPA Reg. No. 264-770

EPA Est. No. 3125-MO-001

ACCEPTED

AUG 31 2005

STOP - Read the label before use Keep out of reach of children

WARNING AVISO

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 264-770

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577
For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

FIRST AID

IF IN EYES:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577.

Have a **LEVERAGE** container or label with you when calling a poison control center or doctor, or going for treatment.

Note To Physician: No specific antidote is available. Treat symptomatically. May pose an aspiration pneumonia hazard.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

May be fatal if swallowed. Causes substantial but temporary eye injury. Causes skin irritation. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Do not get in eyes, on skin or on clothing. Wear goggles or face shield. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse.

Do not contaminate feed or food. Keep out of reach of children.

Personal Protective Equipment: Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves, such as barrier laminate or viton
- Chemical-resistant footwear plus socks
- Protective eyewear
- When mixing, loading or cleaning equipment, wear a chemical-resistant apron.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this products concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

In case of poisoning, call a physician immediately. Have patient lie down and keep quiet.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Additional information may be obtained by consulting your Cooperative Extension Service.

This product is highly toxic to aquatic invertebrates.

Imidacloprid demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or viton
- Chemical-resistant footwear plus socks
- Protective eyewear

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking, invert to prevent leakage. If container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer CropScience Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer CropScience Emergency Response telephone number is 1-800-334-7577, or contact Chemtrec at 800-424-9300.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

RESISTANCE MANAGEMENT

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area.

LEVERAGE[®] 2.7 SE Insecticide contains both Group 3 and Group 4A insecticides. Insect biotypes with acquired or inherent tolerance to these types of products may eventually dominate the insect population if Group 3 and / or Group 4A products are used repeatedly as the predominate method of control for targeted species. This may eventually result in partial or total loss of control of those species by LEVERAGE and/ or other Group 3 and 4A products.

One of the active ingredients in LEVERAGE is a member of the neonicotinoid chemical group. Avoid using a block of more than three consecutive applications of LEVERAGE and / or other Group 4A products having the same or similar mode of action. Following a neonicotinoid block of treatments, Bayer CropScience strongly encourages the rotation to a block of applications with effective products of a different mode before using additional applications of neonicotinoid products. Using a block rotation or windowed approach, along with other IPM practices, is considered an effective use strategy for preventing or delaying an insect pest's ability to develop resistance to this class of chemistry.

Foliar applications of LEVERAGE or other Group 4A products from the neonicotinoid chemical class should not be used on crops previously treated with a long-residual, soil-applied product from the neonicotinoid chemical class.

Other Group 4A, neonicotinoid products used as foliar treatments include: Actara, Assail, CALYPSO, Centric, Clutch, Intruder, PROVADO, TRIMAX and Venom.

Other Group 4A, neonicotinoid products used as seed/ soil treatments include: ADMIRE PRO, Belay, Cruiser, GAUCHO, Platinum and Venom.

Other Group 3, synthetic pyrethroids products include: Ambush, Ammo, Asana XL, BAYTHROID, Capture, Danitol, DECIS, Karate, Mustang Max, Pounce, ProAxis, RENOUNCE and Warrior.

Contact your local extension specialist, certified crop advisor and/or product manufacturer for additional insect resistance management recommendations. Also, for more information on Insect Resistance Management (IRM), visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://irac-online.org/>.

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on an imidacloprid or cyfluthrin label.

ROTATIONAL PLANT-BACK INTERVALS*
IMMEDIATE PLANT-BACK
All crops on this label including the following: Barley, Canola, Corn (field, pop and sweet), Cucurbits, Sorghum, Strawberry, Sugarbeets, Tobacco, and Wheat.
FRUITING VEGETABLES
Eggplant, Groundcherry, Okra, Pepinos, Pepper (including Bell, Chili, Cooking, Pimento and Sweet), Tomatillo, Tomato
HEAD and STEM BRASSICA VEGETABLES
Broccoli, Broccoli raab (<i>rapini</i>), Brussels sprouts, Cabbage, Cauliflower, Cavalo broccoli, Chinese (<i>gai lon</i>) broccoli, Chinese (<i>bok choy</i>) cabbage, Chinese (<i>napa</i>) cabbage, Chinese mustard (<i>gai choy</i>) cabbage, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens.
LEAFY VEGETABLES
Amaranth (leafy amaranth, Chinese spinach, tampala), Arugula (Roquette), Chervil, Chrysanthemum (edible leaved and garland), Cilantro, Corn salad, Cress (garden), Cress (upland, yellow rocket, winter cress), Dandelion, Dock (sorrel), Endive (escarole), Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Spinach (including New Zealand and vine (Malabar spinach, Indian spinach)), Watercress
LEAFY PETIOLE VEGETABLES
Cardoon, Celery, Celtuce, Chinese celery (fresh leaves and stalk only), Florence fennel (including sweet anise, sweet fennel, Finocchio), Rhubarb, Swiss chard
LEGUME VEGETABLES (Edible Podded and Succulent Shelled Peas and Beans and Dried Shelled Peas and Beans)
ROOT VEGETABLES
Beet (garden), Burdock (edible), Carrot, Celeriac, Chervil (turnip-rooted), Chicory, Ginseng, Horseradish, Parsley (turnip-rooted), Parsnip, Radish, Oriental radish (diakon), Rutabaga, Salsify (oyster plant), Salsify (black), Salsify (Spanish), Skirret., Turnip.
TUBEROUS and CORM VEGETABLES
Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Canna (edible, Queensland arrowroot), Cassava (bitter & sweet), Chayote (root), Chufa, Dasheen (taro), Ginger, Leren, Sweetpotato, Tanier (cocoyam), Turmeric, Yam bean (jicama, manioc pea), Yam (true)
30-DAY PLANT-BACK
Cereals (including buckwheat, millet, oats, rice, rye, and triticale), Soybeans and Safflower
10-MONTH PLANT-BACK
Onion and bulb vegetables
12-MONTH PLANT-BACK
All Other Crops
* Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed.

FOLIAR APPLICATIONS

Applications may be made using properly calibrated ground sprayers, fixed- or rotary-winged aircraft or through properly designed, sprinkler-type, chemigation equipment. Thorough and uniform coverage of plants, with direct contact of the spray mixture to the target pests, is required for optimum control. The addition of an organo-silicone based spray adjuvant, at a rate not to exceed the adjuvant manufacturer's recommended use rate may improve coverage. Avoid application procedures where thorough coverage of plant is not possible. Applications made with less than thorough coverage may result in slower activity and/or less overall control from a single application than an application made with higher gallonages. For all aphids, apply as pest population begins to build and prior to build up of damaging levels. Two applications at a 7-day interval may be required to achieve control. For aphid control in crop with dense canopy, use ground application equipment which will provide thorough coverage of lower leaves. See general, **spray drift reduction management**, section below for application guidelines on all application methods.

Ground equipment applications should be made in a minimum of 15 gallons/A.

Aerial applications should be made in a minimum of 5 gallons/A, however, 10 gallons/A are recommended. Aerial applications made to dense canopies may not provide sufficient coverage of lower leaves to provide pest control, especially, aphids. Use only the highest labeled rate of LEVERAGE for aerial applications.

Chemigation applications should be made as concentrated as possible. For best results apply at 100% input for center pivots or 0.10 inch (2,716 gallons) up to 0.15 inch (4,073 gallons) of water/A for other systems. See additional directions and precautions given below. Use only the highest labeled rate for chemigation applications.

For all insects, timing of application should be based on careful scouting and local economic thresholds.

SPECIFIC GUIDELINES FOR USE IN CHEMIGATION SYSTEMS

Types of Irrigation Systems: LEVERAGE may be applied only through sprinkler type irrigation systems. These types include; center pivot, lateral move, or solid set irrigation systems. Do not apply LEVERAGE through any other type of irrigation system.

Injection for Chemigation: Inject the specified dosage of LEVERAGE into the irrigation main, water stream: (1) through a constant flow, metering device; (2) into the center of the main line flow via a pitot tube or equivalent; (3) at a point ahead of at least one, right-angle turn in main stream flow such that thorough mixing with the irrigation water is ensured.

Uniform Water Distribution and System Calibration: The irrigation system must provide uniform distribution of LEVERAGE treated water. Crop injury, lack of effectiveness, or illegal pesticide residues in or on the crop can result from non-uniform distribution. The system must be calibrated to uniformly distribute the rates specified for chemigation application to specific crops. If you have questions about calibration, contact your State Extension Service, equipment manufacturers, or other experts.

Chemigation Monitoring: A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Required Injection and Sprinkler System Safety Devices: The system must contain a functional check valve, vacuum relief valve, and low pressure drain, appropriately located on the irrigation pipeline to prevent water source contamination from back-flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor/engine stops or in cases where there is no water pump, when water pressure decreases to the point where pesticide distribution is adversely affected. Injection systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Using Water from Public Water Systems: Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and to top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional automatic quick-closing check valve to prevent the flow of fluid back toward the injection. The pesticide injection pipeline must contain a functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Chemical Supply Tank Dilution and Agitation: For injection of LEVERAGE use a chemical supply tank for pre-mixing LEVERAGE with water before injecting mixture into the irrigation line. Dilution ratio should be at least 4 parts water to 1 part LEVERAGE. Constant mechanical or hydraulic agitation must be maintained in the chemical supply tank during the entire period of application.

Determine the required amounts of LEVERAGE and water to mix in the tank. The amount of LEVERAGE needed equals the number of fluid ounces of LEVERAGE to be applied per acre multiplied by the number of acres to be chemigated. The amount of emulsion needed equals the gallons of emulsion delivered per hour by the injection pump, multiplied by the number of hours chemigation will take place. The amount of water needed equals the amount of emulsion needed minus the amount of LEVERAGE needed.

Cleaning the Chemical Injection System: In order to apply pesticides accurately, the chemical injection system must be kept clean; free from chemical or fertilizer residues and sediments. Refer to your owner's manual or ask your equipment supplier for the cleaning procedure for your injection system.

Flushing the Irrigation System: At the end of the application period, allow time for all lines to flush the pesticide through all nozzles before turning off irrigation water. To ensure the lines are flushed and free of pesticides, a dye indicator may be injected into the lines to mark the end of the application period.

Center-Pivot and Automatic-Move Linear Systems: Inject the specified dosage per acre continuously for one complete revolution or move of the system. The system should be run at maximum speed. It is recommended that nozzles in the immediate area of control panels, chemical supply tanks, pumps and system safety devices be plugged to prevent chemical contamination of these areas. Use of END GUNS is NOT recommended. End guns which provide uneven distribution of treated water can result in crop injury, lack of effectiveness, or illegal pesticide residues in or on the crop

Solid Set and Manually Controlled Linear Systems: Injection should be during the last 30 to 60 minutes of a regular irrigation period or as a separate 30 to 60 minute application not associated with a regular irrigation.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS; RIVERS; PERMANENT STREAMS, MARSHES OR NATURAL PONDS; ESTUARIES AND COMMERCIAL FISH FARM PONDS.

Spray Drift Reduction Management

The interaction of many equipment and weather related factors determines the potential for spray drift. The applicator is responsible for considering all of these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator. Use the following as a guide for reducing drift onto non-target sites.

Buffer Zone Requirements:

Ground, Foliar Applications: Do not apply by ground within 25 feet of lakes; reservoirs; rivers; permanent streams, marshes or natural ponds; estuaries and commercial fish farm ponds.

Aerial Applications: Do not apply by air within 150 feet of lakes; reservoirs; rivers; permanent streams, marshes or natural ponds; estuaries and commercial fish farm ponds. The spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Importance of Droplet Size: An important factor influencing drift is droplet size. Small droplets (<150 to 200 microns) drift to a greater extent than large droplets. Within typical equipment specifications, applications should be made to deliver the largest droplet spectrum that provides sufficient control and coverage. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Wind Speed Restrictions: Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size, canopy and equipment specifications determine drift potential at any given wind speed. Do not apply when winds are greater than 15 mph and avoid gusty and windless conditions. Avoiding applications when wind direction is toward the aquatic area can reduce risk of exposure to sensitive aquatic areas.

Restrictions During Temperature Inversions: Do not make aerial or ground applications during temperature inversions. Drift potential is high during temperature inversions. Temperature inversions restrict vertical air mixing, which causes small, suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog. However, if fog is not present, the movement of smoke from a ground source can also identify inversions. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical mixing.

Runoff Management: Do not cultivate within 10 feet of the aquatic areas to allow growth of a vegetative filter strip. When used on erodible soils, best management practices for minimizing runoff should be employed. Consult your local Soil Conservation Service for recommendations in your use area. Do not apply if soil is saturated with water. Do not apply under conditions that favor drift from runoff. Do not apply in the rain.

COTTON

Pests Controlled	Rate fluid ounces/Acre
Cabbage looper Cotton aphid Cotton leafworm Cotton leaf perforator Cutworm European corn borer Flea beetle Fleahopper Garden webworm Pink bollworm Saltmarsh caterpillar Southern garden leafhopper Stink bug Threecornered alfalfa hopper Thrips	3.0* (42.7 acres per gallon)
Boll weevil Cotton bollworm (lower rate may be used in Bt cotton, higher rate should be used in non-Bt; for ovicidal effects, use high rate) Lygus bug / Plant bug (in areas with suspected pyrethroid resistance, use high rate) Whitefly (other than sweetpotato whitefly)	3.0* - 3.75 (42.7 - 34.1 acres per gallon)
Grasshopper Tobacco budworm (pyrethroid resistance may limit activity)	3.75 (34.1 acres per gallon)
Pests Suppressed	
Sweetpotato whitefly	3.75 (34.1 acres per gallon)
<p>NOTES and RESTRICTIONS Pre-Harvest Interval (PHI): 14 days. Maximum LEVERAGE allowed per 7 day interval: 3.75 fl oz/A Maximum LEVERAGE allowed per crop season: 22.5 fl oz/A Maximum imidacloprid allowed per crop season: 0.5 lb AI/A. Maximum cyfluthrin allowed per crop season: 0.5 lb AI/A. Applications may be made by ground, aerial or chemigation equipment. Do not graze treated fields after any application of LEVERAGE. See general "DIRECTIONS FOR USE" section for additional information. Do not make more than a total of 10 synthetic pyrethroid applications (one product or combination of products) to a cotton crop in one growing season.</p> <p>* Rate specified for ground sprayer application only. For aerial or chemigation application use only 3.75 fl oz/A.</p>	

HOP

Pests Controlled	Rate fluid ounces/Acre
Aphids	5.8 (22.1 acres per gallon)
<p>NOTES and RESTRICTIONS Pre-Harvest Interval (PHI): 28 days. Maximum LEVERAGE allowed per 21 day interval: 5.8 fl oz/A Maximum LEVERAGE allowed per crop season: 17.4 fl oz/A Maximum imidacloprid allowed per crop season: 0.31 lb AI/A Maximum cyfluthrin allowed per crop season: 0.25 lb AI/A Applications may be made by ground or aerial equipment. See general "DIRECTIONS FOR USE" section for additional information.</p>	

PEPPER

Pests Controlled	Rate fluid ounces/Acre
Aphid Celery leaf-tier Colorado potato beetle (use high rate if pyrethroid resistance suspected) Corn earworm European corn borer Garden webworm Potato leafhopper	3.0* to 3.75 (42.7 - 34.1 acres per gallon)
Beet armyworm (1 st - 2 nd instars) Cabbage looper Thrips (except <i>Thrips palmi</i>) Western yellowstriped armyworm (1 st - 2 nd instars) Whitefly (including sweetpotato whitefly)	3.75 (34.1 acres per gallon)
Pests Suppressed	
Leafminer Pepper weevil <i>Thrips palmi</i>	
<p>NOTES and RESTRICTIONS Pre-Harvest Interval (PHI): 7 days Maximum LEVERAGE allowed per 7 day interval: 3.75 fl oz/A Maximum LEVERAGE allowed per crop season: 18.75 fl oz/A Maximum imidacloprid allowed per crop season: 0.5 lb AI/A. Maximum cyfluthrin allowed per crop season: 0.25 lb AI/A. Applications may be made by ground, aerial or chemigation equipment. See general, "DIRECTIONS FOR USE" section for additional information.</p> <p>* Rate specified for ground sprayer application only. For aerial or chemigation application use only 3.75 fl oz/A.</p>	

POTATO

Pests Controlled	Rate fluid ounces/Acre
Cabbage looper Colorado potato beetle (use high rate if pyrethroid resistance suspected) Cutworm European corn borer Flea beetle Potato leafhopper Potato tuberworm Tarnished plant bug	3.0* to 3.75 (42.7 - 34.1 acres per gallon)
Aphid ¹¹ Potato psyllid	3.75 (34.1 acres per gallon)
<p>NOTES and RESTRICTIONS Pre-Harvest Interval (PHI): 7 days Maximum LEVERAGE allowed per 7 day interval: 3.75 fl oz/A Maximum LEVERAGE allowed per crop season: 15.00 fl oz/A Maximum imidacloprid allowed per crop season: 0.31 lbs AI/A. Maximum cyfluthrin allowed per crop season: 0.26 lbs AI/A. Applications may be made by ground, aerial or chemigation equipment. See general, "DIRECTIONS FOR USE" section for additional information.</p> <p>¹¹ For all aphids, apply as pest population begins to build and prior to buildup of damaging levels. Two applications at a 7-day interval may be required to achieve control. For aphid control in crop with dense canopy use ground application equipment which will provide thorough coverage of lower leaves.</p> <p>* Rate specified for ground sprayer application only. For aerial or chemigation application use only 3.75 fl oz/a.</p>	

TOMATO

Pests Controlled	Rate fluid ounces/Acre
Aphid Colorado potato beetle (use high rate if pyrethroid resistance suspected) European corn borer Stinkbug Tomato fruitworm Tomato hornworm	3.0* to 3.75 (42.7 - 34.1 acres per gallon)
Beet armyworm (1 st - 2 nd instars) Cabbage looper Southern armyworm (1 st - 2 nd instars) Thrips (except <i>Thrips palmi</i>) Tomato pinworm Variegated cutworm Western yellowstriped armyworm (1 st - 2 nd instars) Whitefly (including sweetpotato whitefly)	3.75 (34.1 acres per gallon)
Pests Suppressed	
Dipterous leafminer Flea beetle	3.75 (34.1 acres per gallon)
<p>NOTES and RESTRICTIONS Pre-Harvest Interval (PHI): 7 days Maximum LEVERAGE allowed per 7 day interval: 3.75 fl oz/A. Maximum LEVERAGE allowed per crop season: 18.75 fl oz/A Maximum imidacloprid allowed per crop season: 0.5 lb AI/A. Maximum cyfluthrin allowed per crop season: 0.25 lb AI/A. Applications may be made by ground, aerial or chemigation equipment. See general, "DIRECTIONS FOR USE" section for additional information.</p> <p>* Rate specified for ground sprayer application only. For aerial or chemigation application use only 3.75 fl oz/A.</p>	

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

Net Contents:

ADMIRE, ADMIRE PRO, BAYTHROID, GAUCHO, LEVERAGE, PROVADO, RENOUNCE and TRIMAX are registered trademarks of Bayer CropScience.

Ambush, Actara, Centric, Cruiser, Karate and Platinum are trademarks of a Syngenta Group Company.

Assail is a registered trademark of Nippon Soda Company, LTD.

Ammo, Capture, Mustang Max and Pounce are registered trademarks of FMC Corp.

Asana XL and Intruder are registered trademarks of E. I. DuPont de Nemours & Co.

Danitol and Venom are registered trademarks of Valent USA Corp

Clutch and Belay are registered trademarks of Arvesta Corporation.

ProAxis is a registered trademark of PyTech Chemicals GmbH



Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

Leverage 2.7 Suspension Emulsion Insecticide (PENDING) Submitted 06/24/05.