

264-771

7/30/2003

Page 1 of 10



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

JUL 30 2003

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Mr. R. M. See
Bayer CropSciences, LP
P.O. Box 12014
2 T. W. Alexander Drive
Research Triangle Park, NC 27709

Dear Mr. See:

Subject: Domain DF Herbicide (Update First Aid and Removal of Expiration Date)
EPA Registration No. 264-771
Application Dated July 3, 2003

The expiration date of the conditional registration of the above product is removed. The following conditions are in effect for conditional registration of this product.

1. Submission of a 28-Day inhalation toxicity study in rats.
2. Submission of a special comparative sensitivity study on thyroid hormone levels in neonatal and adult rats.
3. Submission of any data and/or information required to resolve deficiencies remaining once The Agency completes review of submitted data.

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended is acceptable, provided you make the following changes before you release the product for shipment.

1. At the beginning of the list of Personal Protective Equipment (PPE) within the Precautionary Statements, add the statements "Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart." In addition, revise the requirement for "waterproof gloves" to a requirement for "chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride."
2. Within the list of PPE for early re-entry in the Agricultural Use Requirements box, revise the requirement for "waterproof gloves" to a requirement for "chemical-resistant gloves made of any waterproof material."

2 7 10

Submit three (3) copies of final printed labeling incorporating the above changes before you release the product for shipment. Amended labeling supercedes all previously accepted ones. A stamped copy of labeling is enclosed for your records.

Sincerely,

James A Tompkins
James A Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505C)

3710

DomainTM DF Herbicide

For control of certain grass and dicot weeds in soybeans.

ACTIVE INGREDIENTS:

- Flufenacet
N-(4-Fluorophenyl)-N-(1-methylethyl)-2-[[5-(trifluoromethyl)-1,3,4-thiadiazol-2-yl]-oxy]acetamide..... 24.0%
- Metribuzin
4-Amino-6-(1,1-dimethylethyl)-3-(methylthio)-1,2,4-triazin-5(4H)-one..... 36.0%

INERT INGREDIENTS: 40.0%
100.0%

EPA Reg. No. 264-771

EPA Est. No.

**STOP - Read the label before use
KEEP OUT OF REACH OF CHILDREN
CAUTION**

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577
For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
<p>For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577. Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p>	
<p>NOTE TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically.</p>	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, or absorbed through skin. Avoid contact with eyes, skin or clothing.


PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Waterproof gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ACCEPTED
with **COMMENTS**
In EPA Letter Dated:
JUL 30 2003
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
264-771



USER SAFETY RECOMMENDATIONS:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwaters.

Do not allow sprays to drift onto adjacent desirable plants.

Metribuzin has properties that may result in ground water contamination. Application in areas where soils are permeable or coarse and groundwater is near the surface could result in groundwater contamination.

Ground Water: Flufenacet is known to leach through soil into ground water under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water: Flufenacet and metribuzin residues can contaminate surface water through spray drift. Under some conditions, they also have high potential for runoff into surface water (primarily via dissolution in runoff water or erosion), for several to many months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlying tile drainage systems that drain to surface waters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours following application.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, or feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Dispose of pesticide as directed below. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer CropScience Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer CropScience Emergency Response Telephone No. is 1-800-334-7577.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

Domain™ DF Herbicide is a selective soil-applied herbicide for early season control/suppression of many annual grass and dicot weeds in soybeans and should be utilized in conjunction with additional tank mix partners and/or a planned postemergence weed control program. Domain™ is formulated as a 60 % dry flowable. Domain™ may be applied preplant surface, preplant incorporated (mix into the top 1 to 2 inch layer of soil) and preemergence. Domain™ will provide its most effective weed control when applied and subsequently moved into the soil by rainfall, sprinkler irrigation or mechanical tillage prior to weed emergence. Domain™ controls weeds by interfering with seed germination and seedling development and by inhibiting photosynthesis.

Domain™ DF Herbicide may be applied using water as a liquid carrier or be impregnated and applied on dry bulk fertilizer.

Domain™ may be applied either alone or in tank mix combination with additional herbicides. When tank mixing, always observe all precautionary statements and limitations on labeling of all products.

Dry weather following preemergence application of Domain™ or recommended tank mixtures may reduce effectiveness. Cultivate if weeds develop where possible.

GENERAL USE RESTRICTIONS

Domain™ DF Herbicide is intended for use **only** on soybeans.

Do not apply more than 16 ounces of Domain™ per acre per use season.

Soybean seed should be planted a minimum of 1.5 inches deep.

Due to sensitivity of certain soybean varieties/cultivars, Domain™ DF Herbicide is not recommended for use on the following soybean varieties or cultivars: A6520, Altona, AP 2122, AP55, AP71, Burlison, Coker 102, Coker 156, Dassel, GH 1265, GL 3202, Govan, Hartz 5545, Hartz 5545 RR, M83-1010, Maple Amber, Maple Ridge, Narow, Nattawa, NB 2560, NB 2776, NB 3350, Poloma 350, Protage, Regal, S81-84, Semmes, SOI 269, SOI 398, TE 606, Tracy, Vansoy and Vinton 81.

Injury to soybean varieties may occur when Domain™ is used under certain conditions such as when 1) soils have a calcareous surface area or a ph of 7.5 or greater, 2) applied in conjunction with a soil-applied organic phosphate pesticide, 3) applied unevenly or poorly incorporated into the soil, 4) applied to any soil with less than 1/2% organic matter, and 5) heavy rains occur soon after application, especially in poorly drained areas where water may stand for several days.

Do not apply this product through any type of irrigation system. Do not apply aerially.

Do not use flood irrigation to apply, activate or incorporate Domain™ DF Herbicide.

Do not apply Domain™ or Domain™ tank mixtures when environmental conditions may favor drift to non-target sites.

If the soybean crop treated with Domain™ DF Herbicide is lost, only soybeans may be replanted immediately. Do not apply a second application of Domain™.

This product may not be mixed or loaded within 50 feet of any well (including abandoned wells and drainage well), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or property diked mixing/loading areas.

Operations that involve mixing, loading, rinsing or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container rinse or washwater, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over from the pad, which means the pad must be self contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specific minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

This product must be used in a manner which will prevent back siphoning in wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

6 8 10

MIXING INSTRUCTIONS

Liquid Carriers:

Domain™ DF Herbicide is a dry flowable herbicide that must be mixed in water.

Before mixing Domain™ and its labeled tank mixtures, examine the spray equipment making sure it is completely clean and free of rust or corrosion. Be sure the equipment is free of any residues from previously used pesticides. Flush lines with clean water or recommended detergents after the last application. Use an approved method for disposing of rinsate.

For optimal spray tank mixing convenience and efficiency, Domain™ is recommended to be added to the spray tank via an eductor system.

The proper mixing sequence of Domain™ and recommended tank mixtures with the appropriate liquid carrier is as follows:

1. Fill the spray tank or nurse tank 1/4 full with the appropriate liquid carrier.
2. Start recirculation and agitation system and continue throughout mixing and application.
3. If ammonium sulfate is to be used, add it now.
4. Next slowly add the recommended quantity of Domain™ through the eductor system or to the spray tank.
5. If tank mixing with wettable powders or other dry flowable products, they may be added now.
6. If tank mixing with emulsifiable concentrates or soluble products, add the products to the spray tank.
7. If tank mixing with Gramoxone Extra®, a glyphosate-containing product or Touchdown®, add the products to the spray tank.
8. If mixing any additional spray adjuvants in the mixture, add them after all other products have been mixed.
9. Fill the spray tank to the desired level with the appropriate liquid carrier.
10. Continue agitation during transport and application until the spray tank is empty.

Domain™ DF Herbicide and all registered mixtures should be kept agitated once mixed and then sprayed out immediately. Do not allow mixtures to stand for prolonged periods of time. Water quality, pH, temperature and/or other components of the mixture may affect how long the mixture may stand before application.

Dry Fertilizer Carriers:

Domain™ DF Herbicide may be impregnated or coated on dry bulk fertilizers for application. Refer to **Appendix I** of this label for details.

APPLICATION INFORMATION

SPRAYER APPLICATION:

Ground Broadcast Treatment:

Accurately calibrate the sprayer prior to mixing the herbicide treatments. Apply Domain™ and the labeled tank mixtures in a minimum of 10 gallons of total spray volume per acre using broadcast boom equipment. Use a pump with capacity to maintain 30 to 40 psi at the nozzles, maintain adequate in-tank agitation to keep the spray mixture in suspension and provide a minimum of 20% bypass at all times. Applications must be made at a sufficient spray pressure and volume to provide accurate and uniform application of spray particles to a given area without causing spray drift to non-target areas. If mixed with other labeled herbicides, the spray volume may be no less than the minimum volume recommended by the tank mix products or 10 gallons, whichever is greater. The use of screens to protect the pump and nozzles is recommended. Screens placed on the suction side of the pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles. Refer to the nozzle manufacturer for additional recommendations. Agitate thoroughly before and during application with either bypass or mechanical agitation. Rinse the sprayer thoroughly with clean water immediately after each use.

Band Treatment:

Domain™ DF Herbicide and the labeled tank mixtures may be applied as a band treatment. Use the following formula to calculate the amount of herbicide needed for band treatments:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \frac{\text{Broadcast rate}}{\text{per acre}} = \frac{\text{Band rate}}{\text{per acre}}$$

DRY FERTILIZER APPLICATION:

Apply using a minimum of 200 lb of dry fertilizer per acre. See **Appendix I** for additional details.

APPLICATION METHODS AND TIMING

Domain™ DF Herbicide may be applied either alone or in tank mixtures with or sequentially with certain other registered and recommended herbicides. Applications may be made preplant surface, preplant incorporated or preemergence.

Preplant Surface:

For use in conservation, minimum or no-tillage crop production systems, Domain™ DF Herbicide alone or as a recommended tank mixture may be applied as a broadcast spray up to 14 days before planting. If weeds are present at the time of application, Domain™ + crop oil concentrate (1% v/v) will provide burndown of certain broadleaf weeds which are up to 3" tall (See Preplant Burndown Directions). To control a broader spectrum of emerged weeds or if emerged weeds are more than 3" tall, Domain™ may be applied in tank mixture with a recommended nonselective herbicide such as Gramoxone Extra, a glyphosate-containing product or Touchdown. 2,4-D LVE may also be added to any of the above treatments to enhance broadleaf control. Apply only those 2,4-D LVE formulations

which are registered and recommended for preplant and burndown use in soybeans. Refer to the 2,4-D LVE label for specific use rate, application timing and additional restriction information. Following herbicide application, if possible, do not move treated soil out of the row or move untreated soil to the soil surface during planting, as weed control may be reduced.

Preplant Incorporation:

Apply Domain™ alone or in combination with recommended tank mixes as a broadcast spray and incorporate into the upper 1 to 2 inches of the soil surface up to 14 days before planting. Avoid deep incorporation as reduced weed control may result. Incorporate with implements which provide uniform, shallow incorporation (example - finishing disk, harrow, rolling cultivator, etc.)

Preemergence:

Domain™ alone and its recommended tank mixes may be applied to the soil surface as a broadcast spray or band application after planting of soybeans but prior to weed or crop emergence. If weeds are present at the time of application, either apply Domain™ + crop oil concentrate (1% v/v, see burndown weed control chart for weeds controlled) or Domain™ with a nonselective herbicide such as Gramoxone Extra, a glyphosate-containing product or Touchdown. Rainfall and/or overhead sprinkler irrigation is necessary to move Domain™ into the upper soil surface where weed seeds germinate. Dry weather conditions following application may reduce weed control. If adequate moisture is not received within 7 to 10 days after application and weeds begin to emerge from the soil, a light rotary hoeing or shallow incorporation (no deeper than 1/2 inch deep) will improve performance. Excessive rainfall or irrigation after application may reduce weed control.

SOYBEAN

Domain™ DF Herbicide is a selective dry flowable herbicide for the control/suppression of annual grasses and selected broadleaf weeds in soybeans. Domain™ may be applied once per use season either alone or in tank mix combination with or sequentially with certain registered herbicides. The following types of applications are allowed: preplant surface, preplant incorporated and preemergence. The preplant surface and preplant incorporated treatments may be applied up to 14 days before planting. The most effective weed control will occur when the applied product is moved into the soil by rainfall, sprinkler irrigation or mechanical tillage prior to weed emergence from the soil.

Do not graze or feed forage, hay or straw to livestock.

WEED SPECIES CONTROLLED BY DOMAIN™ DF HERBICIDE

Domain™ DF Herbicide applied alone at the recommended dosages and application timings will provide early season (3 to 6 weeks) control of many annual grasses and broadleaf weeds. Length of early season residual weed control is dependent upon the rate of Domain™ applied. To broaden the spectrum of weeds controlled or increase residual control, Domain™ is recommended for use in tank mixture with or sequentially with additional registered herbicides.

EARLY SEASON WEED CONTROL	
ANNUAL GRASS WEEDS	
Barnyard grass (<i>Echinochloa crus-galli</i>)	Foxtail, yellow (<i>Setaria glauca</i>)
Crabgrass, large (<i>Digitaria sanguinalis</i>)	Goosegrass (<i>Eleusine indica</i>)
Crabgrass, smooth (<i>Digitaria ischaemum</i>)	Lovegrass, India (<i>Eragrostis sp</i>)
Foxtail, giant (<i>Setaria faberi</i>)	Panicum, fall (<i>Panicum dichotomiflorum</i>)
Foxtail, green (<i>Setaria viridis</i>)	Signalgrass, broadleaf (<i>Brachiaria platyphylla</i>)
ANNUAL BROADLEAF WEEDS	
Beggarweed, Florida (<i>Desmodium tortuosum</i>)	Ragweed, common (<i>Ambrosia artemisiifolia</i>)
Carpetweed (<i>Mullugo verticillata</i>)	Sesbania, hemp (<i>Sesbania exaltata</i>)
Galinsoga spp. (<i>Galinsoga spp.</i>)	Shepherd's-purse (<i>Capsella bourse-a-pasteur</i>)
Jimson weed (<i>Datura stramonium</i>)	Sicklepod (<i>Cassia tora</i>)**
Kochia (<i>Kochia scoparia</i>)*	Sida, prickly (<i>Sida spinosa</i>)
Lambsquarters, common (<i>Chenopodium album</i>)*	Smartweed, Pennsylvania (<i>Polygonum pennsylvanicum</i>)
Ladysthumb (<i>Polygonum persicaria</i>)	Spurge, spotted (<i>Euphorbia maculata</i>)
Mallow, Venice (<i>Hibiscus trionum</i>)	Sunflower, common (<i>Helianthus annuus</i>)**
Marestail (<i>Erigeron canadensis</i>)	Velvetleaf (<i>Abutilon theophrasti</i>)
Pigweed spp (<i>Amaranthus spp.</i>)*	Waterhemp, common (<i>Amaranthus rudis</i>)*
Purslane, common (<i>Portulaca oleracea</i>)	Waterhemp, tall (<i>Amaranthus tuberculatus</i>)*
Pusley, Florida (<i>Richardia scabra</i>)	
* Domain™ will not provide acceptable control of triazine-resistant biotypes.	
** Weed suppression. Suppressed weeds may be stunted in growth or be of reduced populations as compared to non-treated areas but control will generally not be commercially acceptable.	

8 of 10

PREPLANT BURNDOWN OF BROADLEAF WEEDS

If weeds are emerged at the time of preplant surface application, Domain™ DF Herbicide will provide burndown control of many broadleaf weeds which are up to 3" in height. To improve burndown weed control, Domain™ may be tank mixed with crop oil or other surfactants, sprayable grade fluid fertilizers or with a recommended nonselective herbicide such as a glyphosate-containing product, Touchdown or Gramoxone Extra. For additional broadleaf control, an appropriate formulation of 2,4-D may be added to the mixtures.

PREPLANT BURNDOWN WEED CONTROL BY DOMAIN™ DF HERBICIDE ALONE

ANNUAL BROADLEAF WEEDS

Beggarweed, Florida (<i>Desmodium tortuosum</i>)	Pennycress, field (<i>Thlaspi arvense</i>)
Carpetweed (<i>Mollugo verticillata</i>)	Pigweed spp. (<i>Amaranthus spp</i>)*
Chickweed, common (<i>Stellaria media</i>)	Pineappleweed (<i>Matricaria matricariodes</i>)
Cocklebur, common (<i>Xanthium strumarium</i>)	Purselane, common (<i>Portulaca oleracea</i>)
Galinsoga (<i>Galinsoga parviflorum</i>)	Shepherd's-purse (<i>Capsella bursa-pastoris</i>)
Henbit (<i>Lamium amplexicaule</i>)	Sicklepod (<i>Cassia obtusifolia</i>)
Lambsquarters, common (<i>Chenopodium album</i>)*	Thistle, Russian (<i>Salsola kali</i>)
Lettuce, prickly (<i>Lactuca serriola</i>)	Velvetleaf (<i>Abutilon theophrasti</i>)
Marestail (<i>Conyza canadensis</i>)	Waterhemp, common (<i>Amaranthus rudis</i>)*
Mustard, blue (<i>Chorispura tenella</i>)	Waterhemp, tall (<i>Amaranthus tuberculatus</i>)*
Mustard, wild (<i>Sinapis arvensis</i>)	

* Domain™ DF Herbicide will not provide acceptable burndown control of triazine-resistant biotypes.

DOMAIN™ DF HERBICIDE USE RATES

Domain™ DF Herbicide (dry flowable) may be applied preplant surface, preplant incorporated and/or preemergence at broadcast use rates of **9 to 16 ounces per acre** on most soils which contain 0.5% organic matter or greater. At these use rates, Domain™ will provide **only** early season (from 3 to 6 weeks) residual weed control. The greater the Domain™ rate applied, the longer the early season weed control (up to 6 weeks). Use of the higher rates of Domain™ are also recommended when: weed pressures are heavy, applications are to be made preplant incorporated, early applications are planned and/or on soils with 3% or higher organic matter. Domain™ is **not recommended** for use on sand with less than 1% organic matter or on peat or muck soils with 20% or greater organic matter.

To provide increased residual weed control and/or control of a broader spectrum of weeds, Domain™ is recommended for use in tank mixtures with or sequentially with additional registered herbicides.

DOMAIN™ DF HERBICIDE USE IN TANK MIXTURES AND/OR SEQUENTIAL APPLICATIONS

Domain™ DF Herbicide may be used in tank mixture with or sequentially with any other herbicides which are registered for use in soybeans unless specifically prohibited by the label of the tank mix or sequentially-applied product. These mixtures or sequential applications may be used in conventional, conservation, minimum and no-till crop management systems. Three-way or multiple tank mixtures are permitted unless restricted by the product label. Refer to the individual product labels for recommended use rates, precautions and/or restrictions.

CROP ROTATION RECOMMENDATIONS

In the event of a crop failure, soybeans can be replanted immediately. Do not make a second application of Domain™. After application of Domain™, only the following crops can be planted, after the waiting period as listed below.

IMMEDIATELY	Soybeans		
1 MONTH	Corn	Potatoes	
4 MONTHS	Carrots		
12 MONTHS	Alfalfa	Fescue	Rye
	Barley	Millet, pearl	Sorghum
	Bermudagrass	Millet, prose	Teosinte
	Bluegrass	Oats	Triticall
	Bromegrass	Popcorn	Wheat
	Buckwheat	Rice	Wild Rice
	Clover		

9 7 10

APPENDIX I

IMPREGNATION AND APPLICATION ON DRY FERTILIZER

Dry bulk fertilizer may be impregnated or coated with Domain™ DF Herbicide for applications to soybeans. Follow all Domain™ label recommendations, special instructions and precautions. All state regulations relating to dry bulk fertilizer blending, impregnating and labeling must also be followed.

Rates: Select the recommended Domain™ use rate per acre from the crop section of this label and determine the quantity of dry bulk fertilizer to be applied per acre (use a minimum of 200 pounds and a maximum of 450 pounds per acre). Use the equation below to determine the amount of Domain™ needed per ton of fertilizer applied.

$$\frac{\text{Ounces Domain™ per acre} \times 2000}{\text{lb fertilizer per acre}} = \frac{\text{Ounces Domain™}}{\text{per ton fertilizer}}$$

Impregnation: Domain™ DF Herbicide may be impregnated on many commonly used dry fertilizers but should not be impregnated on ammonium nitrate, fertilizers containing ammonium nitrate, potassium nitrate, sodium nitrate or powder limestone.

To impregnate Domain™ on bulk fertilizer, use a closed rotary-drum mixer or other commonly used dry bulk fertilizer blender equipped with suitable spray equipment. Mix Domain™ with sufficient water to form a sprayable slurry mixture. Spray nozzles must be directed to provide uniform fertilizer coverage while avoiding spray contact with mixing equipment. Nonuniform impregnation can cause crop injury or unsatisfactory performance.

Spray the herbicide mixture onto the fertilizer after blending has started. If necessary, include a suitable drying agent to ensure a spreadable herbicide impregnated fertilizer. If the fertilizer materials are excessively dusty, use diesel fuel or other suitable additive to reduce dust prior to impregnation. Apply the treated fertilizer immediately after impregnation to avoid lump formation and spreading difficulties.

Application: Accurate calibration of fertilizer application equipment and uniform fertilizer distribution are essential for satisfactory weed control. Air flow or auger metered application equipment is preferred (one pass application). If other equipment is used, the recommended application method is to apply one-half the recommended rate and overlap 50 percent or to double apply by splitting the middles to obtain the best distribution pattern.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

NET WEIGHT:

SENCOR, Turbo, Option and Liberty are registered trademarks and Domain is a trademark of Bayer.

Pentagon, Prowl, Pursuit, Raptor, and Scepter are trademarks of American Cyanamid Company.

Basagran, Blazer, Galaxy, Poast, Poast Plus, and Storm are trademarks of BASF Corporation.

FirstRate and Sonalan HFP are trademarks of Dow AgroSciences.

Assure II, Canopy, Canopy XL, Classic, Concert, Lorox, Pinnacle, Reliance STS, and Synchrony STS are trademarks of DuPont.

Authority Broadleaf and Command are trademarks of FMC Corporation.

Roundup and Roundup Ultra are trademarks of Monsanto.

Cobra, Resource and Select are trademarks of Valent.

Flexstar, Fusilade, Fusion, Gramoxone Extra, Tornado, and Touchdown are trademarks of Zeneca.



Bayer CropScience

Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)