

264-806

8/13/2009

1 of 9



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

AUG 13 2009

Mr. Larry R. Hodges
Bayer CropScience LP
2 T.W. Alexander Drive,
Research Triangle Park, NC 27709

Subject: Label Notification(s) for Pesticide Registration Notice 2007-4

Dear Registrant:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 dated July 10, 2009 for:

EPA Registration 264-806

Calypto 4 Flowable Insecticide

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-4 and finds that the label change(s) requested falls within the scope of PRN-2007-4. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact me directly at 703-305-6249 or Banza Djapao of my staff at 703-305-7269.

Sincerely,

A handwritten signature in black ink that reads "Rachel C. Halloran".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

2009

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 264-806	2. EPA Product Manager NOTIF	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Calypso 4 Flowable Insecticide	PM# 501	
5. Name and Address of Applicant (Include ZIP Code) Bayer CropScience LP 2 T. W. Alexander Drive Research Triangle Park, NC 27709 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: NOTIFICATION EPA Reg. No. _____ Product Name AUG 13 2009	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

"Notification of label change per PR Notice 2007-4. This notification is consistent with the guidance in PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA."

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	If "Yes" Unit Packaging wgt. No. per container		<input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Package wgt	No. per container		
3. Location of Net Contents Information <input type="checkbox"/> Label <input checked="" type="checkbox"/> Container		4. Size(s) Retail Container 32 oz, 0.5 gal, 1 gal, 10 gal		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			
Name Larry R. Hodges	Title Registration Manager - Insecticides	Telephone No. (Include Area Code) 919-549-2870	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.			6. Date Application Received (Stamped)
2. Signature 	3. Title Registration Manager - Insecticides		
4. Typed Name Larry R. Hodges	5. Date July 10, 2009		

AUG 13 2009

GROUP	4A	INSECTICIDE
--------------	-----------	--------------------

Calypso[®] 4 Flowable Insecticide

FOR CONTROL OF CERTAIN INSECTS INFESTING VARIOUS CROPS

SHAKE WELL BEFORE USING

ACTIVE INGREDIENT: Thiacloprid

[3-[(6-chloro-3-pyridinyl)methyl]-2-thiazolidinylidene]cyanamide 40.4%

INERT INGREDIENTS: 59.6%

Contains 4 pounds of thiacloprid per gallon.

EPA Reg. No. 264-806

EPA Est. No. 264-MO-002

KEEP OUT OF REACH OF CHILDREN WARNING – AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail).

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> Immediately call a poison control center or doctor for treatment advice. Do not induce vomiting unless told to do so by a poison control center or doctor. Have person sip a glass of water if able to swallow. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> Move Person to fresh air. If Person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
<p>For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577.</p> <p>Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p>	
<p>NOTE TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically.</p>	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS (& DOMESTIC ANIMALS)

WARNING

May be fatal if swallowed. Harmful if inhaled or absorbed through the skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid breathing spray mist. Avoid contact with the eyes, skin or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material, such as Nitrile, Butyl, Neoprene and /or Barrier Laminate and shoes plus socks.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturers instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is highly toxic to marine/estuarine invertebrates. Do not apply directly to water, areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Surface Water Label Advisory

This product may contaminate water through run-off following rainfall events.

Groundwater Label Advisory

The degradate YRC2894 sulfonic acid has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Not for use on residential sites. Do not store in or around residential sites.

Do not apply this product through any kind of irrigation equipment.

Do not allow spray to drift from the application site and contact people, structures people occupy at any time and the associated property, parks and recreation areas, nontarget crops, aquatic and wetland areas, woodlands, pastures, rangelands, or animals.

This product is not registered in NASSAU AND SUFFOLK COUNTIES of New York State. State law prohibits distribution, sale or use of the product in these counties.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is coveralls, chemical-resistant gloves made of any waterproof material, such as Nitrile, Butyl, Neoprene and /or Barrier Laminate and shoes plus socks.

5089

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Do not store in or around residential sites.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Non-refillable container. Do not reuse or refill this container.

(5 Gallons or less)

Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Offer for recycling, if available or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

(Greater than 5 Gallons)

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Offer for recycling, if available or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

MIXING INSTRUCTIONS

To prepare the spray, add a portion of the required amount of water to the spray tank and with agitation add Calypso® 4 Flowable. Complete filling tank with balance of water needed. Maintain sufficient agitation during both mixing and application.

Calypso® 4 Flowable may also be used with other pesticides and/or fertilizer solutions as recommended under specific crop use directions (see NOTE below). When tank mixtures of Calypso® 4 Flowable and other pesticides are involved, prepare the tank mixture as recommended above. When pesticide mixtures are needed, add wettable powders first, Calypso® 4 Flowable, or other flowables second, and emulsifiable concentrates last. Ensure good agitation as each component is added. Do not add the second component in the mixture until the tank contains at least $\frac{1}{2}$ of desired amount of water. If a fertilizer solution is added, a fertilizer pesticide compatibility agent may be needed. Maintain constant agitation during both mixing and application to ensure uniformity of spray mixture.

NOTE: Test compatibility of the intended tank mixture before adding Calypso® 4 Flowable insecticide to the spray or mix tank. Add proportionate amounts of each ingredient in the appropriate order, to a pint or quart jar, cap, shake for 5 minutes, and let set for 5 minutes. Poor mixing or formation of precipitates that do not readily redisperse indicates an incompatible mixture that should not be used. For further information, contact your local Bayer CropScience representative.

RESISTANCE MANAGEMENT

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area. Consult your agricultural advisor for resistance management strategies and recommended pest management practices for your area.

Calypso® 4 Flowable Insecticide contains a Group 4A insecticide called thiacloprid. Insect biotypes with acquired or inherent tolerance to Group 4A products may eventually dominate the insect population if Group 4A products are used repeatedly as the predominant method of control for targeted species. This may eventually result in partial or total loss of control of those species by Calypso® and to other Group 4A products.

The active ingredient in Calypso® is a member of the chloronicotinyl/neonicotinoid (CNI) chemical group and is active at the nicotinic acetylcholine receptor site. Avoid using a block of more than three consecutive applications of Calypso®, and/or other Group 4A products with the same or similar mode of action. Following a CNI block of treatments, Bayer CropScience strongly encourages the rotation to a block of applications with effective products of a different mode of action before using additional applications of CNI products. Using a block rotation or windowed approach, along with other IPM practices, is considered an effective use strategy for preventing or delaying an insect pest's ability to develop resistance to this class of chemistry.

Other Group 4A, CNI products include Actara™, Assail™ and Provado®.

Contact your local extension specialist, certified crop advisor and/or product manufacturer for additional insect resistance management recommendations. For more information on managing insecticide resistance, visit Insecticide Resistance Action Committee (IRAC) on the web at <http://plantprotection.org/IRAC/>.

ENDANGERED SPECIES

Under the Endangered Species Act, it is a Federal Offense to use any pesticide in a manner that results in the death of a member of an endangered species.

To protect the Karner Blue Butterfly in Wisconsin, Michigan and New York consult your county extension agent, or pesticide state lead agency for information concerning this endangered species in the area where you intend to use this product. If this species is present, do not apply within one mile of sandy habitats that support wild lupine plants.

ROTATIONAL CROP RESTRICTIONS

Roots, tubers and bulb vegetables whose tops are not used for feed may be rotated with cotton after a plant back interval of 30 days. Leafy vegetables may be rotated after a 6 month plant back interval. Do not plant any other crop in the treated area for at least one year after application.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS; RIVERS; PERMANENT STREAMS, MARSHES OR NATURAL PONDS; ESTUARIES AND COMMERCIAL FISH FARM PONDS.

Spray Drift Management: The interaction of many equipment and weather related factors determines the potential for spray drift. The applicator is responsible for considering all of these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator.

Buffer Zone Requirements: For **aerial applications**, do not apply within **100 feet** of lakes; reservoirs; rivers; permanent streams, marshes or natural ponds; estuaries and commercial fish farm ponds.

Recommendations For Aerial Applications: The spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Importance of Droplet Size: An important factor influencing drift is droplet size. Small droplets (<150 to 200 microns) drift to a greater extent than large droplets. Within typical equipment specifications, applications should be made to deliver the largest droplet spectrum that provides sufficient control and coverage. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

For aerial applications use only ASABE S572 medium-coarse droplet size spectra (250-450 volume medium diameter measured in microns).

Airblast (Air Assist) Specific Recommendations for Tree Crops and Vineyards: Airblast sprayers carry droplets into the canopy of trees/vines via a radially, or laterally directed air stream. The following specific drift management practices should be followed:

- Adjust deflectors and aiming devices so that spray is only directed into the canopy;
- Block off upward pointed nozzles when there is no overhanging canopy;
- Use only enough air volume to penetrate the canopy and provide good coverage;
- Do not allow the spray to go beyond the edge of the cultivated area (i.e., turn off sprayer when turning at end rows);
- Only spray inward, toward the orchard or vineyard, for applications to the outside rows.

Wind Speed Restrictions: Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size, canopy and equipment specifications determine drift potential at any given wind speed.

Do not apply when winds are greater than 15 mph and avoid gusty and windless conditions. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Restrictions During Temperature Inversions: Do not make aerial or ground applications during temperature inversions. Drift potential is high during temperature inversions. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however if fog is not present, inversions can also be identified by the movement of smoke from a ground source. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical mixing.

RECOMMENDED FOLIAR APPLICATIONS-POME FRUIT

CROP	PEST	RATE PER APPLICATION		SPECIFIC DIRECTIONS
Apple Crabapple Loquat Mayhaw Pear (oriental) Quince	Aphids (except wooly apple aphid) Leafminers Leafhoppers Mirid Bugs	0.5 – 1 fl oz per 100 gal	2 – 4 fl oz per acre ¹	Apply specified dosage as a dilute or concentrate foliar spray as needed. Target most sensitive stages of pest for best results. For optimal leafminer control, target when most eggs have been laid and larvae are in the sap-feeding stage, or at the time of egg hatch. Depending on local conditions, for first generation control, make first application at pink and/or around petal fall or as recommended by your local pest control adviser. A second application for that generation may be required 7-14 days after the first under severe pressure, when extended flight is encountered, when more than one species is to be controlled or when generations are overlapping. A single application may result in suppression only. Calypso® will not control tissue feeding larvae.
	Apple Maggot Codling Moth European Apple Sawfly Oriental Fruit Moth Plum Curculio Japanese beetle Suppression Only: Scale insects	1 – 2 fl oz per 100 gal	4 - 8 fl oz per acre ¹	Apply low rate for low to moderate populations of white leafhoppers and high rate for high populations or for other leafhopper species. Two applications are needed for best control of mirid bugs ; make the first application at pink followed by a second application at petal fall. Consult your local Extension Agent and/or University specialists for codling moth and oriental fruit moth recommendations. Utilize pheromone traps to establish biofix and begin applications around peak egg deposition and prior to egg hatch. Prolonged moth flights may require repeat applications. For plum curculio the adult stage is being targeted. Start applications around petal fall (or earlier) and follow with one to two cover sprays to cover the oviposition period. Applications on 7 to 14 day intervals starting at time of adult emergence with succeeding applications to cover the oviposition period are required for best results against apple maggot . For best results against scale insects , time applications to the crawler stage.

RESTRICTIONS AND PRECAUTIONS: Apples; Crabapple; Loquat; Mayhaw; Pear (oriental) and Quince

- Apply low rate for low to moderate populations and high rate for high populations or for improved or prolonged control.
- Allow 7 or more days between applications.
- Allow at least 30 days between last application and harvest.
- Aerial application of Calypso® may result in slower activity and reduced control relative to results from ground application. For Aerial applications use medium-coarse droplet size spectra (250-450 volume medium diameter measured in microns). For **aerial applications**, do not apply within **100 feet** of lakes; reservoirs; rivers; permanent streams, marshes or natural ponds; estuaries and commercial fish farm ponds.
- Do not apply more than 8 fluid ounces (0.25 lb AI) per acre in a single application.
- Do not apply more than a total of 16 fluid ounces (0.5 lb AI) per acre per year.
- Follow good management practices and do not expose bees to direct application.

¹The amount of Calypso® required per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees. For example, to calculate the rate for aphid control on smaller trees, multiply 0.5-1 fluid ounces times the number of 100 gallons of spray solution required to thoroughly wet, just prior to the point of runoff, one acre of the trees being treated. For concentrate sprays, apply the same amount of product per acre as would be applied in a dilute spray based on tree size and foliage volume.

CROP	PEST	RATE PER APPLICATION		SPECIFIC DIRECTIONS
		1 - 2 fl oz per 100 gal	4 - 8 fl oz per acre ¹	
PEAR	Pear Psylla Aphids Codling Moth Mealybugs Leafminers Pear Midge Japanese beetle Suppression Only: Scale Insects			<p>Apply specified dosage as a dilute or concentrate foliar spray as needed.</p> <p>Target most sensitive stages of pest for best results.</p> <p>For pear psylla control the higher end of the rate range is recommended.</p> <p>For optimal control of mealybug use maximum gallonage for tree size applied with ground application equipment. Insure good spray coverage of the trunk and scaffolding limbs or other resting sites of the mealybug.</p> <p>For optimal leafminer control, target when most eggs have been laid and larvae are in the sap-feeding stage, or at the time of egg hatch. Depending on local conditions, for first generation control, make first application at pink and/or around petal fall or as recommended by your local pest control adviser. A second application for that generation may be required 7-14 days after the first under severe pressure, when extended flight is encountered, when more than one species is to be controlled or when generations are overlapping. A single application may result in suppression only. Calypso® 4 Flowable will not control tissue feeding larvae.</p> <p>Consult your local Extension Agent and/or University specialists for codling moth recommendations. Utilize pheromone traps to establish biofix and begin applications around peak egg deposition and prior to egg hatch. Prolonged moth flights may require repeat applications.</p> <p>For best results against scale insects, time applications to the crawler stage.</p>

RESTRICTIONS AND PRECAUTIONS: Pears

- Allow 7 or more days between applications.
- Allow at least 30 days between last application and harvest.
- Aerial application of Calypso® may result in slower activity and reduced control relative to results from ground application. For Aerial applications use medium-coarse droplet size spectra (250-450 volume medium diameter measured in microns). For **aerial applications**, do not apply within **100 feet** of lakes; reservoirs; rivers; permanent streams, marshes or natural ponds; estuaries and commercial fish farm ponds.
- Do not apply more than 8 fluid ounces (0.25 lb AI) per acre in a single application.
- Do not apply more than a total of 16 fluid ounces (0.5 lb AI) per acre per year.
- Follow good management practices and do not expose bees to direct application.

¹ The amount of Calypso® required per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees. For example, to calculate the rate for aphid control on smaller trees, multiply 0.5-1 fluid ounces times the number of 100 gallons of spray solution required to thoroughly wet, just prior to the point of runoff, one acre of the trees being treated. For concentrate sprays, apply the same amount of product per acre as would be applied in a dilute spray based on tree size and foliage volume.

90f9

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. To the extent allowed by law, all such risks shall be assumed by the user or buyer.


DISCLAIMER OF WARRANTIES: BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Bayer CropScience disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: TO THE EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

NET CONTENTS: 1 GALLON

Calypso is a trademark and Provado is a registered trademark of Bayer.
Assail is a trademark of Nippon Soda Co., Ltd.
Actara is a trademark of the Syngenta Group Co.

Produced for



Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

Calypso 4 Flowable Insecticide (MASTER) Approved 02/16/06, Notification 04/25/06, Notification 09/15/06, Notification 06/30/09

NOTIFICATION

AUG 13 2009

Page 7 of 7