



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

Date of Issuance:

264-816

4/12/2004

Term of Issuance:

Conditional

Name of Pesticide Product:

Betamix B Herbicide

NOTICE OF PESTICIDE:
 Registration
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Bayer CropScience
2 T.W. Alexander Dr.
Research Triangle Park, NC 27709

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.


Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided you agree in writing that:

1. Submit the following outstanding product chemistry data requirement: one year storage stability study (830.6317), within one year from the date of this letter.
2. Submit the analytical method for the determination of the active ingredient in this product to the EPA Analytical Laboratory at Ft. Mead, MD, within one year from the date of this letter.
3. Add the appropriate establishment number to the label.
4. Change order of FIRST AID statements so that the most severe routes of exposure are listed first (i.e. IF IN EYES is first).
5. Add the following statements to the USER SAFETY RECOMMENDATIONS section of the label:
 "Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing."
 "Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing."

Signature of Approving Official:


Date:


James Tompkins, Product Manager (25)
Herbicide Branch, Registration Division (7505C)

4/12/2004

6. Add spray drift management text, see attachment.

You will submit three (3) copies of your final printed labeling before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). A stamped copy of labeling is enclosed for your records.


James Tompkins
Product Manager (25)
Herbicide Branch
Registration Division (7505C)

attachment

Betamix® β Herbicide

APR 12 2004
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

FOR AGRICULTURAL USE ONLY

Postemergence Herbicide for Control of Weeds in Sugar Beets

264-816

ACTIVE INGREDIENT: Phenmedipham*	15.0%
Desmedipham**	15.0%
INERT INGREDIENTS***	70.0%
TOTAL: 100.0%	

Contains 2.7 lbs. active ingredient per gallon.
* 3-methoxycarbonylamino-phenyl-3-methylcarbanilate
** Ethyl m-hydroxycarbanilate carbanilate (ester)
*** Contains Petroleum distillates

EPA Reg. No: 264-816

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN WARNING – AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
IF ON SKIN:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

For **MEDICAL** Emergencies Call 24 Hours A Day 1-800-334-7577.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

NOTE TO PHYSICIAN: May pose an aspiration pneumonia hazard. Contains petroleum distillate.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

Contains Petroleum Distillate. Causes substantial but temporary eye injury. Avoid contact with skin or clothing. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves such as Natural Rubber ≥ 14 mils (includes natural rubber blends and laminates).

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical resistance category selection chart.

Applicators and Other handlers must wear:

- Long-sleeved shirt and long pants

4/10

- Chemical-resistant gloves such as Natural Rubber \geq 14 mils (includes natural rubber blends and laminates)
- Shoes plus socks
- Protective eyewear (such as goggles, face shield, or safety glasses)

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

Remove and wash contaminated clothing before reuse.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE; DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep closed. Store in a cool, dry place. Do not use or store near heat or open flame.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

When packaged in Plastic containers:

Container Disposal: Triple rinse (or equivalent), then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or by other procedures approved by State and local authorities.

DO NOT REUSE THIS CONTAINER.

DESTROY WHEN EMPTY.

[These statements are deleted when SVR packaging is used.]

When packaged in SVR containers:

ECHO SYSTEM® SVR Return Procedure: Return the ECHO SYSTEM SVR container clean (outside only) and empty to the place of business from which the Betamix® β Herbicide was purchased.

This ECHO SYSTEM SVR container is the sole property of Bayer CropScience.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read the entire Directions for Use before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves such as Natural Rubber \geq 14 mils (includes natural rubber blends and laminates)
- Shoes plus socks
- Protective eyewear (such as goggles, face shield, or safety glasses)

GENERAL INFORMATION

When used as directed, Betamix® β Herbicide is selective against weeds in sugar beets. Follow label directions carefully to avoid severe injury to sugar beets. For best results, spray weeds in the cotyledon stage which are actively growing and are not under water or heat stress. Betamix® β will control the following weeds:

Annual sowthistle	<i>Sonchus oleraceus</i>
Black nightshade	<i>Solanum nigrum</i>
Coast fiddleneck	<i>Amsinckia intermedia</i>
Common chickweed	<i>Stellaria media</i>
Common lambsquarters	<i>Chenopodium album</i>
Common ragweed	<i>Ambrosia artemisiifolia</i>
Green foxtail	<i>Setaria viridis</i>
Groundcherry	<i>Physalis lanceifolia</i>
Hairy nightshade	<i>Solanum sarrachoides</i>
Kochia	<i>Kochia scoparia</i>
London rocket	<i>Sisymbrium irio</i>
Nettleleaf goosefoot	<i>Chenopodium murale</i>
Yellow foxtail (pigeon grass)	<i>Setaria glauca</i>
Prostrate pigweed*	<i>Amaranthus gracizans</i>
Portulaca	<i>Portulaca oleraceus</i>
Redroot pigweed*	<i>Amaranthus retroflexus</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>
Wild buckwheat	<i>Polygonum convolvulus</i>
Wild mustard	<i>Brassica kaber</i>

*Redroot pigweed and prostrate pigweed control will be improved with a tank mix of Betamix® β and Betanex® Herbicide in Eastern North Dakota and Minnesota (see Chart 4).

GENERAL PRECAUTIONS AND RESTRICTIONS

DO NOT APPLY BETAMIX® β HERBICIDE TO SUGAR BEETS LATER THAN 75 DAYS PRIOR TO HARVEST.

DO NOT EXCEED A TOTAL OF 6 PINTS BETAMIX® β HERBICIDE PER ACRE PER SEASON.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

DO NOT ROTATE WITH CEREAL GRAIN CROPS FOR 120 DAYS FOLLOWING POSTEMERGENCE APPLICATION OF BETAMIX® β HERBICIDE.

BETAMIX® β HERBICIDE MAY CAUSE BEET INJURY IF THE CROP IS UNDER STRESS FROM ONE OR MORE OF THE FOLLOWING CONDITIONS:

- Rapid climatic changes from cool, overcast days, to hot (80°F or over) bright days. When the air temperature is, or is likely to be, above 80°F on the day of spraying, application should be made in the late afternoon when the temperature is decreasing.
- Frost within 3 days prior to application or 7 days following treatment could cause beet injury.
- Windy conditions or drought

6/10

- Use of a preplant or preemergence herbicide or other chemicals
- Insect or disease injury
- Close cultivation

If stress conditions are present, delay application in order to give plants a chance to recover.

IMPORTANT: Betamix® β Herbicide may cause temporary growth retardation and/or chlorosis or tipburn on sugar beets. Sugar beets usually resume normal growth within 10 days.

DO NOT OVERTREAT:

The use of higher than recommended rates may cause beet injury and/or carry over problems when tank mixed with Nortron® SC Herbicide.

Do not spray while dew is present.

Rainfall or sprinkler irrigation within 6 hours of spraying may reduce weed kill.

Do not allow spray drift to contact adjacent crops which may be injured by spray drift.

MIXING THE SPRAY:

Make sure the sprayer is CLEAN.

Betamix® β emulsifiable concentrate formulation contains sufficient wetting agents for optimum coverage. Do not add additional wetting agents or other spray adjuvants except as specified for "Micro-Rate Applications". Add sufficient water to fill the lines. Then add the desired amount of Betamix® β Herbicide and the remaining quantity of water with the bypass agitator running. Bypass agitation is sufficient. Mechanical agitation is not necessary. Only use freshly prepared spray emulsions.

Always spray immediately after preparing the spray solution. Prepare only enough spray solution to last less than four hours.

APPLICATION INSTRUCTIONS

MULTIPLE (LOW RATE) APPLICATIONS:

Multiple (low rate) applications of Betamix® β Herbicide may be applied by air or ground to sugar beets to control early germinating weeds. The first application must be applied when the earliest emerging weeds have reached cotyledon size. See Chart 1 for broadcast rates. For broadcast applications with ground equipment, apply in 10 to 20 gallons of water per acre. Use 5 to 15 gallons of water per acre with aerial application. See Chart 2 for equivalent band rates. Any weeds which are not completely controlled by the first treatment will usually be checked and controlled by repeat applications. The repeat application should be made 5 to 7 days after the preceding application or when another flush of weeds germinates. If the second application is delayed, conventional treatment as described below will be necessary.

To avoid excessive phytotoxicity to fall-planted sugar beets south of the Tehachapi Mountains in California when temperatures are above 85°F, apply Betamix® β Herbicide at the rate of 0.5 pint per acre (broadcast equivalent). Evening applications are recommended.

Betamix® β Herbicide applied postemergence in a tank mix with Nortron® SC Herbicide (see Chart 3) broadens and enhances the control of troublesome weeds, in addition, provides control of Ladysthumb (*Polygonum persicaria*), Pennsylvania smartweed (*Polygonum pennsylvanicum*), Annual bluegrass (*Poa annua*) and Canarygrass (*Phalaris canariensis*).

For further information, contact your County Agricultural Agent, Farm Advisor, or Bayer CropScience.

CHART 1

DOSAGE CHART FOR MULTIPLE (LOW RATE) BROADCAST APPLICATIONS

Weed Stage*	Pints/Acre Broadcast			
	BETAMIX® β ALONE	BETAMIX® β	+	NORTRON® SC
Cotyledon	0.75-1.5	0.75	+	0.25
2 leaf	1.0-1.5	1.0	+	0.33
4 leaf	1.5-2.25	1.5	+	0.5

*Applications should begin at the cotyledon stage of the weeds.

*Higher dosage rates could be required, depending on the advancement of the weed stage.

*Do not exceed 0.75 pt./acre when sugarbeets are at the cotyledon stage.

*Early two true-leaf sugar beets tend to be the most susceptible to phytotoxicity.

CHART 2

BETAMIX® β HERBICIDE DOSAGE CHART FOR BAND APPLICATION

Broadcast Equivalent	Band Width	Band Rate - Row Spacing (fluid ounces)			
		22"	24"	28"	30"
0.75 pints/acre	5"	2.75	2.5	2.1	2.0
	7"	3.8	3.5	3.0	2.8
1.0 pints/acre	5"	3.6	3.3	2.8	2.6
	7"	5.1	4.6	4.0	3.75
1.5 pints/acre	5"	5.4	5.0	4.3	4.0
	7"	7.6	7.0	6.0	5.6
2.25 pints/acre	5"	8.2	7.5	6.4	6.0
	7"	11.4	10.5	9.0	8.4
3.0 pints/acre	5"	10.9	10.0	8.5	8.0
	7"	15.25	14.0	12.0	11.2
3.75 pints/acre	5"	13.6	12.5	10.7	10.0
	7"	19.1	17.5	15.0	14.0

CHART 3

NORTRON® SC DOSAGE CHART FOR MULTIPLE (LOW RATE) BAND APPLICATIONS

Broadcast Equivalent	Band Width	Band Rate - Row Spacing (fluid ounces)			
		22"	24"	28"	30"
0.25 pints/acre	5"	0.9	0.8	0.7	0.7
	7"	1.3	1.2	1.0	0.9
0.33 pints/acre	5"	1.2	1.1	0.9	0.9
	7"	1.7	1.5	1.3	1.2
0.5 pints/acre	5"	1.8	1.7	1.4	1.3
	7"	2.5	2.3	2.0	1.9

CONVENTIONAL APPLICATIONS

By Ground: Apply Betamix® β Herbicide at the rate of 2.25 to 3.75 pints per acre in 20 to 50 gallons of water broadcast basis. For band application, see Dosage Chart 2.

By Air: Apply Betamix® β Herbicide at the rate of 2.25 to 3.75 pints per acre using 5 to 15 gallons of spray per acre.

Apply the 2.25 to 3.75-pint rates only to sugar beets past the two true-leaf stage. Use the 3.75-pint rate only on well-established sugar beets which are not under stress. The stage of growth of the weeds is very important for satisfactory control. For best results, spray when the weeds are at the two true-leaf stage or smaller, are actively growing and are not under water or heat stress.

In order to avoid phytotoxic spray drift to nontarget crops during application of Betamix® β Herbicide, the following buffer zones should be observed:

Cotton, Potatoes, Sunflowers,
Sorghum, Wheat 50 feet

Blackeye beans, Cabbage, Flax..... 100 feet

Lettuce, Rape, Tomatoes 300 feet

DO NOT APPLY WHEN WIND SPEED IS OVER 10 MILES PER HOUR. AVOID APPLICATIONS WHEN CONDITIONS FAVOR DRIFT.

REPEAT APPLICATION OF BETAMIX® β HERBICIDE: For control of later germinating weeds, make a second application of Betamix® β Herbicide. Use 2.25 to 3 pints of Betamix® β . Allow at least 7 days between first and second applications. Apply when sugar beets

have at least 4 leaves. For best results, use the higher rate and spray when weeds are at the two true-leaf stage. Apply lower rates when the sugar beets are under stress as explained in the General Precautions And Restrictions section.

TANK MIX COMBINATIONS

When tank mixing, read and follow the label for each tank mix product used for precautionary statements, directions for use, weeds controlled, geographic and other restrictions. Use in accordance with the most restrictive of label limitations and precautions. No label dosage should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

Betamix® β Herbicide can be tank mixed with the following broadleaf herbicides for improved broadleaf weed control if application timing is correct for the tank mix products.

Herbicide	Use Rate (pt./A)
Stinger®*	0.25–0.50
Betanex®	See Chart 4
Betanex® β	See Chart 5

*The Betamix® β + Stinger® tank mix should be applied when sugar beets are in the two true-leaf stage or larger.

Chart 4

Dosage Chart for Tank Mixes of Betamix® β Herbicide and Betanex® Herbicide

Equivalent Betamix® β Desired Rate (Pints/acre Broadcast)	Betamix® β + Betanex® (Pints/acre Broadcast)		
0.57	0.28	+	0.57
0.62	0.32	+	0.63
0.67	0.33	+	0.67
0.75	0.37	+	0.75
0.87	0.44	+	0.88
1.12	0.57	+	1.13
1.62	0.81	+	1.63
2.50	1.25	+	2.50

Chart 5

Dosage Chart for Tank Mixes of Betamix® β Herbicide and Betanex® β Herbicide

Equivalent Betamix® β Desired Rate (Pints/acre Broadcast)	Betamix® β + Betanex® β (Pints/acre Broadcast)		
0.57	0.28	+	0.28
0.62	0.32	+	0.32
0.67	0.33	+	0.33
0.75	0.37	+	0.37
0.87	0.44	+	0.44
1.12	0.57	+	0.57
1.62	0.81	+	0.81
2.50	1.25	+	1.25

MICRO-RATE APPLICATIONS (EXCEPT CALIFORNIA)

Multiple Micro-rate applications of Betamix® β Herbicide in tank mixtures with reduced rates of UpBeet®, Stinger®, and modified seed oils may be applied by air or ground equipment to sugar beets to control early germinating weeds.

When adding spray adjuvants to Betamix® β Herbicide the rate must not exceed 0.08 lb a.i./A (see Dosage Chart 6 below) when sugar beets are in the cotyledon to 4-true-leaf stage. When the smallest sugar beet plants in the field are in the 4-true-leaf stage, the rate can be increased to 0.12 lb a.i./A (see Dosage Chart 6 below). The use of wetting agents or spray adjuvants with conventional rates (0.73 to 1.22 lb a.i./acre) or multiple low rate (0.24 to 0.73 lb a.i./acre) applications of Betamix® β is prohibited.

Favorable climatic conditions (good conditions for plant growth and development) are essential for adequate weed control.

DOSAGE CHART 6
DOSAGE CHART FOR MULTIPLE MICRO-RATE BROADCAST APPLICATIONS

Sugar Beet Stage	Betamix® β Herbicide Fluid Ounces/Acre Broadcast
Cotyledon to 4-leaf	4.0 (equivalent to 0.08 lb. ai/A)
4-Leaf*	4.0 – 6.0 (equivalent to 0.08 – 0.12 lb. ai/A)

* Rate can be increased when the smallest sugar beet plants in the field are in the 4-true leaf stage or larger.

Application of Betamix® β Herbicide in broadcast applications is strongly recommended. If band applications are used, do not use less than 11-inch bands.

For broadcast applications of Betamix® β with selected tank mix partners, apply in 10 to 20 gallons of water per acre for ground application, or 5 to 15 gallons of water per acre for aerial application. Use the minimum rate recommended on the tank mix partner label, or a reduced rate of the tank mix partner(s), at the discretion of the grower or applicator, as permitted under FIFRA. [Minimum label rate for UpBeet® is 0.5 oz/acre; for Stinger®, 4.0 fl oz/acre.]

Use modified seed oils at a finished spray concentration of 1.5% v/v or a minimum of 1 pt/acre. A minimum of three sequential applications should be used. Accurate timing is essential; make initial application immediately after weeds emerge, and make repeat applications on 5- to 7-day intervals. If weed control is not adequate due to climatic conditions, spray coverage or other factors, return to multiple (low rate) applications.

Betamix® β Herbicide can be mixed with UpBeet®, Stinger®, and modified seed oils for use on sugar beets in accordance with the most restrictive label limitations and precautions. No label dosage rates should be exceeded. Betamix® β Herbicide cannot be mixed with any product containing a label prohibition against such mixing.

Fungicides or insecticides can be tank mixed with Betamix® β plus UpBeet® plus Stinger® plus methylated seed oils, however, do not combine both fungicides and insecticides with micro-rate mixtures.

MIXING INSTRUCTIONS FOR MICRO-RATE MULTIPLE APPLICATIONS OF BETAMIX® β HERBICIDE

1. Start with a clean spray tank.
2. Fill spray tank with one-third of the total amount of clean water needed for application and start gentle agitation.
3. Slurry UpBeet® in water before adding to spray tank, then add slurried UpBeet® to spray tank.
4. Fill spray tank to two-thirds of the total amount of clean water needed for the application.
5. Add Betamix® β Herbicide followed by Stinger®, then modified seed oil.
6. Add remaining amount of water while continuing gentle agitation. Spray immediately. Spray mixture should not remain in spray tank overnight.

USE PRECAUTIONS FOR MICRO-RATE APPLICATIONS

Not all weeds will be adequately controlled, even with favorable climatic conditions. Micro-rate applications of Betamix® β mixed with UpBeet® and Stinger® will not control ALS-resistant kochia. Multiple low rates of Betamix® β and/or hand labor may be required if multiple micro-rate applications do not adequately control weeds.

Modified seed oils must not be added if the Betamix® β rates exceed the rates listed in Dosage Chart 6 above, as the addition of modified seed oils could increase the possibility of crop injury at dosage rates greater than those listed in Dosage Chart 6.

Multiple micro-rate applications may injure sugar beets if climatic conditions rapidly change from cool, wet, overcast days to bright sunny days. Plugging of spray nozzles may be encountered due to the potential formation of a precipitate in the spray solution that is often associated with micro-rate applications. To minimize potential formation of precipitate, start with a clean spray system, use warm spray water for mixing, completely empty spray solution from each tank load, flush tank and lines between loads with fresh water, never leave diluted spray solution in tank overnight, and/or add ammonia (2% household) at 1% v/v or a basic blend additive (as referenced in the most recent North Dakota State University Weed Control Guide) at 1% v/v. DO NOT apply micro-rate treatments when conditions are favorable for drift to nontarget species.

10/10

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Bayer CropScience disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

NET CONTENTS:

Betamix, Betanex, Nortron and Echo System are registered trademarks of Bayer.
Stinger is a registered trademark of DowElanco.
Upbeet is a registered trademark of E.I. duPont de Nemours and Co.



Bayer CropScience

Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

Betamix β Herbicide (PENDING) Submitted 10/8/03, Resubmitted 2/20/04.