2 5 20



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

> OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

FEB 5 2010

Mr. Larry R. Hodges Product Registration Bayer Environmental Science P.O. Box 12014 2T.W Alexander Drive Research Triangle Park, NC 27709

SUBJECT: Application for Pesticide Notification (PRN 98-10) Request General Label Change/California State Requirements EPA Reg. No. 264-935 Application Dated December 9, 2009

Dear Registrant:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 12/09/09 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

Linda Arrington Notifications & Minor Formulations Team Leader Registration Division (7505P) Office of Pesticide Programs

Please read instruction	ns on r	everse before comp.	ja form.		Form Ap	prove	MB No. 20	70-0060	
\$€PA		Environmenta	inited States I Protection ngton, DC 2046			Image: A state of the state	Registrat Amendm Other		OPP Identifier Number
	Application for Pesticide - Section I								
1. Company/Product Number 264-935				2. EPA Product Manager NOTIF			3. Pro	posed Classification	
4. Company/Product (Allegiance-FL Sec		atment Fungicide		PM#		<u></u>			
5. Name and Address of Applicant (Include ZIP Code) Bayer CropScience LP 2 T. W. Alexander Drive Research Triangle Park, NC 27709			de)	6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No.					
				Section	duct Name				
 Amendment - Explain below. Resubmission in response to Agency letter dated ✓ Notification - Explain below. 			dated	[1	ter dat Applica	ation.	IOTIF	ICATION - 5 2010
Explanation: Use additional page(s) if necessary. (For section I and Section II.) Notification per PR Notice 98-10: As required by the state of California, the full product name "Allegiance-FL Seed Treatment Fungicide" has been inserted in place of the abbreviated product name "Allegiance-FL" throughout the label. "This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA."									
				Section	- 111				
				Water Soluble Packaging Yes Vo		2. Type of Containe Metal V Plastic Glass Paper		r	
* Certification mu be submitted	ıst	lf "Yes" Unit Packaging wgt.	No. per container	if "Yes" Package wgt	No. per containe	r		Other (S	pecify) poly drum
3. Location of Net Contents Information 4			4. Size(s) Retai 1, 2.5	I Container , 15, 30, 40, 5	50 gal.	5. Lo	cation of Labe	l Directio	ns
6. Manner in Which La	<u></u>		Lithogra Paper gl Stencile	ph ued d	Othe	r			
				Section -	· IV				
1. Contact Point (Con	nplete i	tems directly below f	or identification	of individual t	be contacted,	if nec	essary, to proc	cess this	application.)
Name Larry R. Hodges			1	1			elephone 919-549-	a No. (Include Area Code) 2870	
	hat any	nents have made on knowlinglly false or aw.		ll attachments					6. Date Application Received (Stamped)
2. Signature LyR. Hodges				3. Title Registration Manager - Insecticides					
4. Typed Name Larry R. Hodges			5.	5. Date December 9, 2009					
		Previous editions are					A File Copy		Yellow - Applicant Conv

NOTIFICATION

FEB - 5 2010

ALLEGIANCE[®]-FL Seed Treatment

Fungicide

A SEED TREATMENT CHEMICAL FOR CONTROL OF SEED ROT AND DAMPING-OFF DISEASES OF CERTAIN CROPS

ACTIVE INGREDIENTS:

Metalaxyl: N-(2,6-dimethylphenyl)-N-(methoxyacetyl)alanine methyl ester	28.35%
INERT INGREDIENTS:	71.65%
TOTAL	100.00%

Contains 2.6 pounds. Metalaxyl per gallon.

EPA Reg. No. 264-935

EPA Est.

KEEP OUT OF REACH OF CHILDREN WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

For <u>MEDICAL</u> And <u>TRANSPORTATION</u> Emergencies <u>ONLY</u> Call 24 Hours A Day 1-800-334-7577 For <u>PRODUCT USE</u> Information Call 1-866-99BAYER (1-866-992-2937)


FIRST AID

IF ON SKIN OR CLOTHING:	Take off contaminated clothing.		
	Rinse skin immediately with plenty of water for 15-20 minutes.		
	Call a poison control center or doctor for treatment advice.		
IF SWALLOWED:	Call a poison control center or doctor immediately for treatment advice.		
	Have person sip a glass of water if able to swallow.		
	Do not induce vomiting unless told to do so by a poison control center or doctor.		
	Do not give anything by mouth to an unconscious person.		
IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15-20 minutes.		
	Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.		
	Call a poison control center or doctor for treatment advice.		

PRECAUTIONARY STATEMENTS

WARNING HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes skin irritation. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Do not get on skin or on clothing. Avoid contact with eyes. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.



ιτευσε

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear: Coveralls over short-sleeved shirt and short pants, socks and chemical-resistant footwear and chemical-resistant gloves (such as nitrile, butyl, neoprene, and/or barrier laminate).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory: This chemical is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where solids are permeable, particularly where the water table is shallow, may result in groundwater contamination

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal or by cleaning of equipment. PROHIBITIONS

Open dumping is prohibited. Do not reuse empty containers.

PESTICIDE STORAGE

Store in a cool dry location. Do not store in direct hot sunlight. Do not store above 90°F for prolonged periods of time. After prolonged storage in intense cold, allow product to return to approximately 75°F before using.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration. or, if allowed by state and local authorities by burning. If burned, stay out of smoke.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. 🗟 also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Warker Protection Standard. ουιςςί ι L

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

Exception: if the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Works, Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with allything that has been treated. 6000 c

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and (ital involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, socks and chemical resistant for wear, chemical resistant gloves, such as barrier laminate or nitrile rubber or neoprene rubber or Viton and protective eyewear.

د ت ر Page 2 of 7

In California - Only for use in commercial seed treating establishments. Not allowed for "on-farm" use.

NOTE: Seed treated commercially or for sale must be dyed an unnatural color using an EPA approved dye before being sold. Federal law requires that bags containing treated seeds shall be labeled with the following information. This seed has been treated with ALLEGIANCE-FL Seed Treatment Fungicide which contains the active ingredient metalaxyl. Do not use for feed, food or oil purposes. Store away from feeds and foodstuffs. If using ALLEGIANCE-FL Seed Treatment Fungicide treated rice do not use Ridomil for control of *Pythium* in water-seeded rice.

ALLEGIANCE-FL Seed Treatment Fungicide is a systemic fungicide seed dressing specifically for control of systemic downy mildews, *Pythium* and *Phytophthora spp*. For control of other soilborne diseases, combination of Captan and VITAVAX® registered formulations are compatible. Do not use with other seed treatment products unless previous experience assures compatibility.

ALLEGIANCE-FL Seed Treatment Fungicide may be applied as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment.

NOTE: ALLEGIANCE-FL Seed Treatment Fungicide is a systemic fungicide having a specific mode of action. ALLEGIANCE-FL Seed Treatment Fungicide could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, Bayer CropScience cannot assume liability for crop damage resulting from resistant strains of fungi. Consult your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible ALLEGIANCE-FL Seed Treatment Fungicide resistant strains of fungi which may occur. FAILURE TO FOLLOW DIRECTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

<u> </u>	
Canola* Mustard seed* Rapeseed*	For <i>Pythium</i> damping-off control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at 0.25-0.5 fl oz per 100 pounds of seed. *Use not permitted in California unless otherwise directed by supplemental labeling.
Cotton	For <i>Pythium</i> seed rot and damping-off: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at the rate of 0.75-1.5 fl oz per 100 pounds of seed.
Soybeans	For <i>Pythium</i> damping-off and early season <i>Phytophthora</i> control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at the rate of 0.75-1.5 fl oz per 100 pounds of seed.
Peanuts*	For <i>Pythium</i> seed rot and damping-off control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at 0.75 fl oz per 100 pounds of seed. *Use not permitted in California unless otherwise directed by supplemental labeling.
Sorghum	For control of <i>Pythium</i> damping-off apply 0.375-0.75 fl oz of ALLEGIANCE-FL Seed Treatment Fungicide per 100 pounds of seed applied as a seed treatment. For control of systemic downy mildew (races susceptible to chemical control), apply 0.75-1.5 fl oz ALLEGIANCE-FL Seed Treatment Fungicide per 100 pounds of seed on cultivars which possess sorghum downy mildew resistance. To those cultivars which do not possess resistance to a particular race of sorghum downy mildew in heavily infected areas with recent severe history, apply 1.5-3.0 fl oz of ALLEGIANCE-FL Seed Treatment Fungicide per 100 pounds of seed as a seed treatment. Only high quality sorghum seed (90% germination and above) should be treated with ALLEGIANCE-FL Seed Treatment Fungicide.
Sunflowers	For control of systemic downy mildew: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at the rate of 1.5-3.0 fl oz per 100 pounds of seed.
Forage Legumes	Alfalfa, clover, lespedeza, beans (for forage), soybeans (for forage), soybean hay, peanuts (for forage), peanut hay, peas (for forage), pea vine hay, cowpeas (for forage), cowpea hay, trefoil, vetch and velvet beans (for forage).
	For <i>Pythium</i> damping-off and early season <i>Phytophthora</i> control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at the rate of 0.75-1.5 fl oz per 100 pounds of seed.
Grain Crops	Wheat, barley, rye, oats, millet, milo, corn (field corn, sweet corn and popcorn), buckwheat and triticale.
	For <i>Pythium</i> damping-off control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at the rate of 0.75 fl oz per 100 pounds of seed. For systemic downy mildew control of sweet corn: Apply &LEGANCE-FL Seed Treatment Fungicide as a seed treatment at the rate of 1.5 fl oz per 100 pounds of seed.
Rice*	For Pythium damping-off control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at the rate of 0.75-1.5 fl oz per 100 pounds of seed. If using ALLEGIANCE-FL Seed Treatment Fungicide treated rice do not use Ridomil for control of Pythium in water-seeded rice. *Use not permitted in California unless otherwise directed by supplemental labeling.
Seed and Pod Vegetables	Black-eyed peas, chickpeas (Garbanzos), cowpeas, dill, edible soybeans, field beans, tield peas, garden peas, green beans, kidney beans, lima beans, lupines, navy beans, okra, peas, pole beans, shap beans, string beans, wax beans, and lentils. For <i>Pythium</i> damping-off control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at the rate of 0.75 fl oz per 100 pounds of seed. For early season <i>Phytophthora</i> control: Apply ALLEGIANCE-FL as a seed treatment at the rate of 0.75 fl oz per 100 pounds of seed. For control of <i>Pythium</i> damping-off and systemic downy mildew of peas: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at the rate of 1.5 fl oz per 100 pounds of seed.

Page 3 of 7

Turf Grasses	For <i>Pythium</i> damping-off control on golf courses, home lawns reclamation and erosion control projects: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treated at the rate of 1.5 fl oz per 100 pounds of seed.					
Forage Grasses	Grasses grown for hay, grazing or silage, corn (fodder or silage), sorghum (hay or silage), and small grain grown for hay, grazing or silage. For <i>Pythium</i> damping-off control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at the rate of 0.75 fl oz per 100 pounds of seed.					
Brassica (Cole) Leafy Vegetables*	Broccoli (including Chinese and raab), Brussels sprouts, cabbage (including Chinese bok choy, Chinese napa and mustard), cauliflower, cavalo broccoli, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, and rape greens					
	For <i>Pythium</i> damping-off control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at the rate of 0.75 fl oz per 100 pounds of seed.					
	*Use not permitted in California unless otherwise directed by supplemental labeling.					
Cucurbit Vegetables*	Chayote (fruit), Chinese waxgourd, citron melon, cucumber, gherkin, edible gourd (includes hyotan, cucuzza Chinese okra, and hechima), Momordica spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), muskmelon (includes true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon), pumpkin, summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), winter squash (includes butternut squash, calabaza, hubbard squash), Cucumis mixta Cucumis pepo (includes acorn squash, spaghetti squash), watermelon (includes hybrids and/or varieties or Citrullus lanatus.)					
	For <i>Pythium</i> damping-off control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at 0.75 to oz per 100 pounds of seed.					
	*Use not permitted in California unless otherwise directed by supplemental labeling.					
Leafy Vegetables*	Amaranth (leafy, Chinese spinach, tampala), cardoon, celery (including Chinese), celtuce, chervil, chrysanthemun (edible-leaved and garland), corn salad, cress (garden and upland), dandelion, dock, endivefennel (finochio) lettuce (head and leaf), orach, parsley, purslane (garden and winter), radicchio, rhubarb, spinach* (including Nev Zealand and vine), and Swiss chard.					
	*Use not permitted in California unless otherwise directed by supplemental labeling. NOTE: Do not apply ALLEGIANCE-FL Seed Treatment Fungicide to spinach which is destined to be grown in the greenhouse Containers or bags containing ALLEGIANCE-FL Seed Treatment Fungicide –treated spinach seed must be labeled with this restriction.					
	For <i>Pythium</i> damping-off control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at 0.75 oz per 100 pounds of seed.					
Fruiting Vegetables*	Eggplant, ground cherry, pepino, pepper (including bell pepper, chili pepper, cooking pepper, pimento, swee pepper), tomatillo, and tomato					
	For <i>Pythium</i> damping-off control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at 0.75 fl oz per 100 pounds of seed.					
	*Use not permitted in California unless otherwise directed by supplemental labeling.					
Onions*	Onion (dry bulb and green)					
	For <i>Pythium</i> damping-off control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at 0.75 fluid ounces per 100 pounds of seed.					
	*Use not permitted in California unless otherwise directed by supplemental labeling.					
Root and Tuber Vegetables* (except potatoes)	Arracacha, arrowroot, artichoke (Chinese and Jerusalem), beet (garden and sugar), burdock (edible), canna (edible), carrot, cassava (bitter and sweet), celery root, chayote, chervil, chicory, chufa, dasheen, ginger, ginseng horseradish, leren, parsley (turnip-rooted), parsnip, radish (includes oriental dalkon), rutabaga, salsify (includes black and Spanish), skirret, sweet potato, tanier, turmeric, turnip, yam bean (jicama, manioc pea), yam					
	For Pythium damping-off control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at 0.79 fl oz per 100 pounds of seed.					
	NOTE: Do not apply ALLEGIANCE-FL Seed Treatment Fungicide to potatoes.					
	*Use not permitted in California unless otherwise directed by supplemental labeling.					

(

۰ د

 ι
 ι
 ι
 ι

 ι
 ι
 ι
 ι

 ι
 ι
 ι
 ι

 ι
 ι
 ι
 ι

 ι
 ι
 ι
 ι

 ι
 ι
 ι
 ι

 ι
 ι
 ι
 ι

 ι
 ι
 ι
 ι

 ι
 ι
 ι
 ι

 ι
 ι
 ι
 ι

 ι
 ι
 ι
 ι

 ι
 ι
 ι
 ι

 ι
 ι
 ι
 ι

 ν
 ν
 ι
 ι

 ν
 ν
 ι
 ι

 ν
 ν
 ι
 ι

 ν
 ν
 ι
 ι

 ν
 ν
 ι
 ι

 ν
 ν
 ι
 ι

 ν
 ν
 ι
 ι

 ν
 ν
 ι
 ι

 ν
 ν
 ι
 ι

 ν
 ν
 ι
 ι

 ν
 ν
 ι
 ι

 ν
 ν
 ι
 ι

 ν
 ν
 ι
 ι

 ν
 ν
 ι

 ν
 ν</

6069

(

SEED TREATMENT-EXPORT USE

For treatment of the following seed for export only to countries requiring treatment with metalaxyl before importation.

Brassica (Cole) Leafy Vegetables	For <i>Pythium</i> damping-off control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at the rate of 0.75-1.50 fl oz per 100 pounds of seed.
Canola	For <i>Pythium</i> damping-off control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at the rate of 0.50-1.70 fl oz per 100 pounds of seed. (32 – 110 ml per 100 kg of seed)
Corn	For <i>Pythium</i> damping –off control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at 0.75-1.50 fl oz per 100 pounds of seed. For systemic downy mildew control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at 3.0 fl oz per 100 pounds of seed.
Cucurbit Vegetables	For <i>Pythium</i> damping-off control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at 0.75 fl oz per 100 pounds of seed.
Fruiting Vegetables	For <i>Pythium</i> damping-off control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at 0.75- 1.50 fl oz per 100 pounds of seed.
Leafy Vegetables	For <i>Pythium</i> damping-off control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at 0.75- 1.50 fl oz per 100 pounds of seed.
Onions	For <i>Pythium</i> damping-off control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at 0.75- 1.50 fl oz per 100 pounds of seed.
Millet	For <i>Pythium</i> damping-off control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at 0.75- 1.50 fl oz per 100 pounds of seed. For systemic downy mildew control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at 2.5 fl oz per 100 pounds of seed.
Peas	For <i>Pythium</i> damping-off and early season Phytophthora control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at 3.5 fl oz per 100 pounds of seed.
Root and Tuber Vegetables	For <i>Pythium</i> damping-off control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at 0.75-1.50 fl oz per 100 pounds of seed.
	Do not apply ALLEGIANCE-FL Seed Treatment Fungicide to potatoes.
Sorghum	For <i>Pythium</i> damping-off control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at 0.75- 1.50 fl oz per 100 pounds of seed. For systemic downy mildew control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at 5.25 fl oz per 100 pounds of seed.
Sunflowers	For systemic downy mildew control: Apply ALLEGIANCE-FL Seed Treatment Fungicide as a seed treatment at 9.0 fl oz per 100 pounds of seed.

Seed treated in accordance with the export use pattern must be labeled as follows:

•

.

FOR EXPORT ONLY

NOT FOR DOMESTIC SALES OR USE.

ιιιίι ε¢ (сc cc c c ... ι ι ι ι ιιιι ιιιι ໍ່. Page 5 of 7


DIRECTIONS FOR USE AT REDUCED RATES IN COMBINATION WITH OTHER FUNGICIDES

· · ·

.

ALLEGIANCE-FL Seed Treatment Fungicide is a systemic fungicide seed dressing which may be applied in combination with other EPA-registered fungicides to aid in control of *Pythium*. ALLEGIANCE-FL Seed Treatment Fungicide may be applied as a water-based slurry through standard slurry or mist-type commercial seed treatment equipment.

Corn (field), Wheat, Oats, Barley, Rye	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply ALLEGIANCE-FL Seed Treatment Fungicide as a commercial seed treatment at the rate of 0.10-0.375 fl oz per 100 pounds of seed. Apply only in combination with EPA registered rates of Bayer CropScience broad-spectrum seed treatment fungicides.
Soybeans, Field beans, Kidney beans, Navy beans, Pinto beans, Lentils, Peas	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply ALLEGIANCE-FL Seed Treatment Fungicide as a commercial seed treatment at the rate of 0.10-0.375 fl oz per 100 pounds of seed. Apply in combination with EPA registered rates of broad-spectrum seed treatment fungicides.
Sunflowers	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply ALLEGIANCE-FL Seed Treatment Fungicide as a commercial seed treatment at the rate of 0.10-0.375 fl oz per 100 pounds of seed. Apply in combination with EPA registered rates of broad-spectrum seed treatment fungicides.
Peanuts*	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply ALLEGIANCE-FL Seed Treatment Fungicide as a commercial seed treatment at the rate of 0.10-0.375 fl oz per 100 pounds of seed. Apply in combination with EPA registered rates of broad-spectrum seed treatment fungicides. *Use not permitted in California unless otherwise directed by supplemental labeling.
Rice*	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply ALLEGIANCE-FL Seed Treatment Fungicide as a commercial seed treatment at the rate of 0.10-0.375 fl oz per 100 pounds of seed. Apply in combination with EPA registered rates of broad-spectrum seed treatment fungicides. *Use not permitted in California unless otherwise directed by supplemental labeling.



IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, disclaimer of Warranties and Limitations of Liability.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Bayer CropScience, no claims are made to guarantee germination of carry-over seed.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

NOTICE TO BUYER

ι,

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

NET CONTENTS:

VITAVAX® is a registered trademark of Chemtura Corporation RIDOMIL® is a trademark of Syngenta Crop Protection Inc. ALLEGIANCE® is a registered trademark of Bayer

PRODUCED FOR

Bayer CropScience

Bayer CropScience LP P.O. Box 12014, 2 T.W. Alexander Drive Research Triangle Park, North Carolina 27709 1-866-99BAYER (1-866-992-2937)

ALLEGIANCE-FL Seed Treatment Fungicide (MASTER) Approved 09/17/07, Notif 11/23/09

