

# UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

AUG 18 2009

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Jamin Huang, Ph.D.
Bayer CropScience
2 TW Alexander Drive
P. O. Box 12014
Research Triangle Park, NC 27709

Subject: Updated Container and Disposal Statement

Dear Dr. Huang:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 dated July 24, 2009 for:

EPA Registration 264-959 EPA Registration 264-963

Gaucho 75 ST Insecticide Gaucho 75 ST FS Insecticide

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-4 and finds that the label change(s) requested falls within the scope of PRN-2007-4. The label has been date-stamped "Notification" and will be placed in our records.

If you have any questions, call me at 703 305-5409 or electronically at daniel.dani@epa.gov

Sincerely,

Dahi Daniel

Registration Division (7505P) Insecticide/Rodenticide Branch

2049

Please read instructions on	reverse beforenple	ting form.	VOTIE	Form App	ved. OMB No.	2070-006	O. Approval expires 2-28-95
C FDA		inted States	*UTIFICAI	IUN	Registra		OPP Identifier Number
<b>\$EPA</b>	Environmental Washi	l <b>Protecti</b> on ngton, DC 20-	on Agency -	=	Amendr	nent	
					✓ Other		
	<del></del>	Application	on for Pestici			-	
1. Company/Product Number	264-959		2. EPA	Product Mana Venus Ea	-	3. Pr	oposed Classification
4. Company/Product (Name Gaucho	) 75 ST Insecticide		PM#	Team 0	1		
5. Name and Address of Ap	plicant (Include ZIP Co	de)	6. Exp	edited Reve	eiw. In accorda	nce with	FIFRA Section 3(c)(3)
Bayer CropScience			(b)(i), n				mposition and labeling
P.O. Box 12014, 2 T		ve	to:	Rea. No.			
Research Triangle P							
Check if thi	s is a new address			ct Name _		·	
			Section - I	1			
Amendment - Explai	n below.		[ ]	•	labels in repsonse	e to	
Resubmission in res	ponse to Agency letter	dated		Agency lette "Me Too" A			
✓ Notification - Explain	ı below.			Other - Expla	in below.		
Explanation: Use addition							
REGFEE: Notification, Fee	amount due: \$ 0 nended Gaucho 75 ST Ir	nsecticide to up	odate container dispo	sal statement	per PRN 2007-4. I	No new da	ta needed to be reviewed by
CONTACT: Jamin Huang,	jamin.huang@bayercro	pscience.com					
			Section - I				
1. Material This Product Wi	li Re Packaged In:	<del> </del>	Jection - II				
Child-Resistant Packaging	Unit Packaging	<del></del>	Water Soluble P	ackaging	2. Type of	Container	
Yes	Yes		Yes			Metal	
✓ No	√ No		√ No			Plastic Glass	
* Certification must be submitted	If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per container		Paper Other (S	specify)
3. Location of Net Contents	Information	A Cirolol Po	tail Container	1	5. Location of Lab	al Directio	
			1 lb, 10 lbs, 50 lbs	1	F	ei Direcuc	ons .
6. Manner in Which Label is	Container  Affixed to Product	r 7	raph glued iled	Other			
	<del></del>	Stenc	Section - I	<u>,,                                   </u>			
1. Contact Point  Complete	items directly below f	or identification			f nacessary to pre	ocess this	application 1
Name			Title				e No. (Include Area Code)
Jamin Huan	g, Ph.D.		Product Registrat	tion Manager		(919) 549	
		Certifica					6. Date Application
I certify that the state I acknowledge that a both under applicable	ments I have made on ny knowlinglly false or i Iaw.	this form and misleading sta	all attachments the atement may be pur	ereto are true, sishable by fin	accurate and con e or imprisonment	hplete.	Received (Stamped)
2. Signature	^		3. Title				( )
( Jan	- Kus-f			egistration Man	ager, Insecticides		( (
4. Typed Name		X 1	5. Date				
Jamin Huang, Ph. D.			July 24, 2009			(	

# Bayer CropScience

BAYER BAYER

July 24, 2009

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, Virginia 22202-4501

ATTENTION: Linda Arrington (Team leader, Notifications & Minor Formulation)

Re: Notification to Update Container Disposal Statement per PR Notice 2007-4 for Gaucho 75 ST Insecticide (EPA Reg. No. 264-959)

Dear Ms. Arrington,

Byaer CropScience is herein submitting a Notification to update the container disposal statement per PR Notice 2007-4 on the 08/13/2007 EPA latest approved label for Gaucho 75 ST Insecticide. Three copies of the amended label dated 07/22/2009 are attached.

Bayer CropScience 2 T.W. Alexander Drive P. O. Box 12014 Research Triangle Park, NC 27709 Tel: 919 549-2000

"Notification of label change per PR Notice 2007-4. This notification is consistent with the guidance in PR Notice 2007-4 and the requirements of EPA regulations at 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the confidential statement of formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the requirements of 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA."

Please let me know at <u>jamin.huang@bayercropscience.com</u> or at 919-549-2634 if you have any questions regarding this submission.

Sincerely,

Jamin Huang, Ph.D.

Product Registration Manager

# NOTIFICATION

4A INSECTICIDE

# GAUCHO® 75 ST Insecticide

**ACTIVE INGREDIENT:** 

EPA Reg. No. 264-959

**EPA Est.** 

# **KEEP OUT OF REACH OF CHILDREN CAUTION**

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577 For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

# **FIRST AID**

IF SWALLOWED:	Call a poison control center or doctor immediately for treatment advice.
	Have person sip a glass of water if able to swallow.
	Do not induce vomiting unless told to do so by a poison control center or doctor.
	Do not give anything by mouth to an unconscious person.
IF ON SKIN OR	Take off contaminated clothing.
CLOTHING:	Rinse skin immediately with plenty of water for 15-20 minutes.
	Call a poison control center or doctor for treatment advice.
IF INHALED:	Move person to fresh air.
	If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
	Call a poison control center or doctor for further treatment advice.
IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15-20 minutes.
	Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
	Call a poison control center or doctor for treatment advice.
In case of eme	ergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577.

# PRECAUTIONARY STATEMENTS

Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

# HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, absorbed through skin or inhaled. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

# **ENVIRONMENTAL HAZARDS**

This pesticide is highly toxic to birds and aquatic invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Cover or incorporate spilled treated seeds. Plant end rows last to cover any exposed seeds.

### **Resistance Management**

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area.

Gaucho 75 ST contains a Group 4A insecticide. Insect biotypes with acquired or inherent resistance to Group 4A insecticides may eventually dominate the insect population if Group 4A insecticides are used repeatedly as the predominant method of control for targeted species.



The active ingredient in Gaucho 75 ST is a member of the neonicotinoid chemical class. Insect pests resistant to other chemical classes have not shown cross-resistance to Gaucho 75 ST. In order to maintain susceptibility to this class of chemistry in insect species with high resistance development potential, it is recommended that for each crop season: 1) only a single seed application of Gaucho 75 ST or other soil applied neonicotinoid product be made to a crop; 2) foliar applications of products from this same class not be made following soil application of other neonicotinoid products.

Other Group 4A, neonicotinoid products used as foliar treatments include: Actara, Assail, CALYPSO, Centric, Clutch, Couraze, Gallant, Impulse, Intruder, LEVERAGE, Nuprid, Pasada, PROVADO, TRIMAX PRO and Venom.

Other Group 4A, neonicotinoid products used as soil/seed treatments include: Advise, Alias, Belay, Couraze, Cruiser, ADMIRE PRO, Macho, Macho Max, Nuprid, Platinum, Venom and Widow.

Contact your Cooperative Extension specialist, certified crop advisor and/or product manufacturer for additional insect resistance management recommendations. Also, for more information on Insect Resistance Management (IRM), visit the Insecticide Resistance Action Committee (IRAC) on the web at <a href="http://www.irac-online.org">http://www.irac-online.org</a>.

# **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use in commercial seed treaters only. Not for use on agricultural establishments in hopper-box, slurry-box, or other seed treatment applications at, or immediately before, planting. This product is to be used in treaters designed for pelleting or coating seed. Mix thoroughly before use or use entire container at one time. All mixes should be pre-tested to determine physical compatibility between formulations. Observe all cautions and limitations on labeling of all products used in mixtures.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal, or by cleaning of equipment.

#### **PESTICIDE STORAGE**

Store in a cool place. Do not store in direct sunlight. Protect from freezing temperatures.

#### **PESTICIDE DISPOSAL**

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

## **CONTAINER DISPOSAL**

Rigid, Non-refillable containers small enough to shake (i.e., with capacities equal to or less than 50 lbs)

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, offer for recycling, if available or dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

#### Fiber canister

Non-refillable container. Do not reuse or refill this container. Completely empty box into application equipment. Then, offer for recycling, if available or dispose of empty box in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Notification of the crop rotational restriction must be conveyed to the grower by appropriate seed tag labeling or bag printing of all seed units.

Treated seed must not be used for or mixed with food or animal feed or processed for oil. Seed commercially treated with GAUCHO 75 ST must be labeled in accordance with all applicable requirements of the Federal Seed Act.

Labels for commercially treated seed should include the following addition to the Environmental Hazards statements:

• Exposed treated seed may be hazardous to birds. Dispose of all excess treated seed and seed packaging by burial away from bodies of water. Cover or incorporate spilled treated seeds.

Do not graze or feed livestock on treated areas (wheat, barley, triticale, sorghum) for 45 days after planting.

**NOTE**: The purchaser of this product is responsible for ensuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals. Refer to 21CFR, Part 2.25. Any dye or colorant added to treated seed must be cleared for use under 40CFR, Part 180.1001. Federal regulations have established official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates. Do not overtreat GAUCHO 480 or GAUCHO 600-treated seed with GAUCHO 75 ST.

#### **Stored Seed Protection:**

Applied at labeled rates equal to or above 0.67 oz. per hundredweight, GAUCHO 75 ST will provide protection to seed against injury from the following insects: Indian Meal Moth (*Plodia interpunctella*), Rice Weevil (*Stilophilus oryzea*), Red Fligur beetle (*Tribolium castaneum*), and Lesser Grain Borer (*Rhizopertha dominica*). It is recommended that seed with existing populations of stored grain pest be fumigated prior to treating and bagging of seed.

# For Early Season Protection Against Certain Sucking Insects:

GAUCHO 75 ST will aid in the protection of seeds and seedlings against injury by certain early season insects.

#### Carrot:

To provide early season protection of seedlings against injury by seed corn maggot and wireworm, apply as a commercial seed treatment at 5.33 oz. per hundredweight of seed.

# Canola, Rapeseed:

To provide early season protection of seedlings against injury by aphids, flea beetle and wireworm apply 8.53 to 21.3 ounces per hundredweight of seed prior to planting. In areas where flea beetles and foliar insects are in high numbers, the higher application rate is recommended. Please consult your local agricultural office for pest patterns, history, and forecasts to assist in determining the appropriate rate for your region. For suppression of Lygus bugs, including *Lygus* spp., in their second generation and Cabbage Seedpod Weevil larvae, including *Ceutorhynchyus assimilis*, apply 12.8 to 21.3 ounces per hundredweight of seed.

#### Field Corn:

For the protection of corn plants from the corn insect pests listed below. Apply as a seed treatment at the recommended rates listed.

	GAUCH	O® 75 ST
Pest	MG. AJ / KERNEL	OZ. / 80,000 UNIT OF SEED
Corn root worm (including Northern, Western, Southern and Mexican) 1	1.34	5.0
Flea beetle	1.54	3.0
Chinch bug .		
Southern green stinkbug		
White grub		
Seed corn maggot		
Thrips		
Wireworm		
Corn leaf aphid		
Imported fire ant		
Southern corn leaf beetle		
Bill bug <sup>2</sup>		
Grape colaspis <sup>2</sup>		
Black cutworm <sup>3</sup>		

	GAUCHO <sup>®</sup> 75 ST			
Pest	MG. AI / KERNEL	OZ. / 80,000 UNIT OF SEED		
Flea beetle	0.6	2.24		
Chinch bug	<b></b>	2.21		
Southern green stinkbug				
Seed corn maggot				
Thrips				
Wireworm				
Corn leaf aphid	i i			
Imported fire ant				
Grape colaspis <sup>2</sup>				
White grub				
	GAUCH	O <sup>®</sup> 75 ST		
Pest	MG. AI / KERNEL	OZ. / 80,000 UNIT OF SEED		
Seed corn maggot (seed protection only)	0.16	0.6		
Wireworm (seed protection only)	0.10	0.0		
Flea beetle (through 1 leaf stage)				
Imported fire ant				
White grub⁴				

In areas of heavy to severe corn rootworm populations, protection will not be adequate. Use only in areas of light to moderate corn rootworm populations. Consult your State Agricultural Extension Service on levels of corn rootworm populations.

3

<sup>&</sup>lt;sup>2</sup> Reduces early season feeding damage.

<sup>&</sup>lt;sup>3</sup> Will reduce feeding damage caused by leaf feeding black cutworms that are ½ inch or less in length.

<sup>&</sup>lt;sup>4</sup> Reduces feeding damage during emergence and seedling stages.

#### Wheat, Barley, Triticale:

To provide early season protection of seedlings against injury by aphids (including Bird cherry-oat, English grain, Greenbug, and Russian wheat aphid), Hessian fly and Wireworms, apply 0.67 to 2.0 ounces per hundredweight of seed prior to planting. Ensure thorough coverage. Low use-rate for "wireworm-only" protection: GAUCHO® 75 ST applied at the 0.11 – 0.22 oz. per hundredweight of seed offers suppression of wireworm activity on seed and young seedlings. To provide early season protection from grasshopper damage, GAUCHO 75 ST should be applied to seed at 1.0 to 2.0 oz. per hundredweight. To reduce early season damage caused by grasshopper, GAUCHO treated seed may be planted as a 50 to 60 foot border around the edges of the field. Consult your local university extension entomologist for details regarding grasshopper control in your area. For maximum effectiveness, seed should be treated uniformly. Ensure thorough coverage. Use the higher rate to provide increased length of protection and from heavy insect pressure and from wireworms, and to reduce potential spread of Barley yellow dwarf virus due to aphid vectors. Do not graze or feed livestock on treated areas for 45 days after planting.

#### Sorghum:

To provide early season protection of seedlings against injury by aphids (including Corn leaf, English grain, Greenbug, and Yellow sugar cane aphid), Chinch bugs, Fire ants and Wireworms, apply 5.3 ounces per hundredweight of seed prior to planting. Ensure thorough coverage. Do not graze or feed livestock on treated areas for 45 days after planting.

#### Cotton (Delinted Seed Only):

To provide early season protection of seedlings against injury by early season thrips and aphids, apply 10.72 ounces per hundredweight of seed prior to planting. Ensure thorough coverage.

#### Sugar Beet:

To provide early season protection of seedlings against injury by Whitefly, Aphids, Leafhoppers (including to reduce potential for spread of Curly top and Yellow mosaic virus due to aphid and leafhopper vectors), Root aphid, Thrips and Wireworms, apply as a commercial seed treatment in a water slurry containing 2.0 to 4.25 ounces of GAUCHO 75 ST in or on a unit of pelleted sugarbeet seed with a weight ratio of 2:1 pelleting mixture to raw seed (seed count 100,000 – approximately 1 kilogram by weight) at the rate of 2.0 ounces per unit. If rates exceed 2.0 ounces per unit, seed must be pelleted. Film coat application and pelleting companies must be approved by Bayer CropScience.

#### **Sweet Corn:**

For the protection of sweet corn plants from the insect pests listed below. Apply as a seed treatment at the recommended rates listed.

PESTS	GAUCHO® 75 ST OZ./CWT OF SEED	
Flea beetle	5.33	
Early season corn leaf aphid		
Seed corn maggot		
Wireworm		
Imported fire ant	2.67 – 5.33	
Early season corn leaf aphid		
Seed corn maggot		
Wireworm		
Imported fire ant	1.33 – 2.67	
Seed corn maggot (seed protection)		
Wireworm (seed protection)		
Wireworm (seed protection)	0.67 - 1.33	

Adzuki Bean, Asparagus Bean, Broad Bean (Succulent or Dry), Catjang Bean, Chinese Longbean, Field Bean, Guar Bean, Jackbean, Kidney Bean, Lablab Bean, Lima Bean (Succulent or Dry), Moth Bean (Succulent or Dry), Mung Bean, Navy Bean, Pinto Bean, Rice Bean, Runner Bean, Snap Bean, Sword Bean, Tepary Bean, Ure Bean, Wax Bean, Yardlong Bean, Blackeyed Pea (Succulent or Dry), Chickpea, Cowpea (Succulent or Dry), Crowder Pea, Dwarf Pea, Edible-Pod, English Pea, Field Pea, Garden Pea, Green Pea, Pigeon Pea (Succulent or Dry), Snow Pea, Southern Pea (Succulent or dry), Sugar Snap Pea, Grain Lupin, Sweet Lupin, White Sweet Lupin, Lentil:

To provide early season protection of seedlings against injury by wireworm, bean leaf beetle, imported fire ant, and aphid, apply as a commercial seed treatment at 1.33 – 2.67 oz. per hundredweight of seed prior to planting.

For treating of seed destined for planting in states other than California. Treated seed is not to be planted in California.

### Peanuts:

To provide early season protection of seedlings against injury by thrips, apply as a commercial seed treatment at 0.67 – 1.33 oz. per hundredweight of seed prior to planting.

Increases in Tomato spotted wilt virus (TSWV) incidence have been observed with applications of imidacloprid as a seed treatment on multiple varieties of peanuts. This may also be the case with other tospoviruses, or other viruses transmitted by various thrips species or perhaps other pests. Prior to applying GAUCHO 75ST to peanuts, Bayer CropScinece recommends consultation with State, Cooperative Extension Service, or Bayer CropScience representative, for recommendations. Growers are advised to weigh insect control benefits against potential increase in viral disease levels. In areas where TSWV or other tosppvirus are endemic, growers are encouraged to use virus resistant varieties and consult the University of Georgia, Tomato spotted wilt virus index, before treating seed with GAUCHO 75ST.

80f9

## **ROTATIONAL CROPS:**

Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application. The following plant-back intervals should be observed for listed crops:

ROTATIONAL PLANT-BACK INTERVALS*					
IMMEDIATE PLANT-BACK				30-DAY PLANT-BACK	
Artichoke' Barley Brassica (cole) leafy veg. Canola Cilantro Corn, Field Corn, Sweet Cotton	Cranberry Cucurbits Eggplant Ground cherry Leaf petiole veg. Leafy veg. Legume veg. (succulent or dried, including soybean)	Mustard Seed Okra Peanut Pepinos Pepper Popcorn Potato Rapeseed	Strawberry Sugar beet Tomatillo Tomato Triticale Veg. Root and Tuber Watercress Wheat	Cereals, including: buckwheat millet rice Safflower	
		Sorghum			

<sup>\*</sup> Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed.

<sup>\*</sup> For all other crops not listed on an imidacloprid label, or for crops for which no tolerance for the active ingredient has been established, a 12-month plant-back interval should be observed.

# **IMPORTANT: READ BEFORE USE**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Bayer CropScience, no claims are made to guarantee germination of carry-over seed.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

#### NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

**NET CONTENTS:** 5 pounds

GAUCHO is a registered trademark of Bayer

Produced for



Bayer CropScience LP P.O. Box 12014, 2 T.W. Alexander Drive Research Triangle Park, North Carolina 27709 1-866-99BAYER (1-866-992-2937)

GAUCHO 75 ST Insecticide (MASTER) EPA APPROVED 08/13/07, Notification 07/22/09