64-964

9/21/2005 jacket

IL® 2.6F Seed Treatment Fungicide

FOR USE ONLY BY COMMERCIAL SEED TREATERS FOR CONTROL OF SPECIFIED DISEASES ON WHEAT, BARLEY, OATS AND TRITICALE **ACTIVE INGREDIENTS:**

Tebuconazole, Alpha-[2-(4-Chlorophenyl)ethyl]-alpha-(1,1-dimethyl-ethyl)-1H-1,2,4-triazole-1-ethanol......

TOTAL

28.3%

100.0%

Contains 2.6 lbs. tebuconazole per gallon

EPA Reg. No. 264-964

EPA Est.

STOP - READ THE LABEL BEFORE USE

KEEP OUT OF REACH OF CHILDREN CAUTION

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577 For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

FIRST AID

SWALLOWED:	Call a poison control center or doctor immediately for treatment advice.		
1	Have person sip a glass of water if able to swallow.		
	Do not induce vomiting unless told to do so by poison control center or doctor.		
	Do not give anything by mouth to an unconscious person.		
IF ON SKIN OR CLOTHING:	Take off contaminated clothing.		
	Rinse skin immediately with plenty of water for 15-20 minutes.		
	Call a poison control center or doctor for treatment advice.		
IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15-20 minutes.		
	Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.		
	Call a poison control center or doctor for treatment advice.		
IF INHALED:	Move person to fresh air.		
	If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.		
	Call a poison control center or doctor for further treatment advice.		
li li	n case of emergency call toil free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577.		

PRECAUTIONARY STATEMENTS

Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes and clothing. Avoid breathing vapor or spray mist. Wear long sleeved shirt, long pants or coveralls, and chemical-resistant gloves made of waterproof material. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to estuarine invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on agricultural establishments in hopper-box, planter-box, slurry-box or other seed treatment applications at or immediately before planting.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Contact Bayer CropScience or your chemical supplier for container return instructions. If container is not returned in accordance with instructions, triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or incineration, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

SEED LABELING REQUIREMENTS

To meet U.S. Federal Seed Act requirements, all seed treated with RAXIL 2.6F should be labeled:

"Treated with alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol (RAXIL). Do not use treated seed for feed, food or oil purposes. Barley, wheat and oat green forage may be grazed or harvested for hay 31 days after seeding."

RAXIL controls or suppresses certain designated seed and soilborne diseases of wheat, barley and oats.

SEED TREATMENT: It is recommended that RAXIL 2.6F be applied as a water-based slurry through standard slurry or mist-type rommercial seed treatment equipment. Prepare a slurry by mixing the specified dosage in 9 to 20 fl. oz. of water per 100 lbs. of seed. .dd dye** to the resulting slurry. Mix the slurry thoroughly with the seeds to provide uniform coverage.

**21 CFR Chapter I, Section 2.25 requires grain seed treated with poisonous substances in excess of recognized tolerances be suitably colored with an EPA approved dye, such as one of the dyes listed in 40 CFR Section 180.1001 (c) and (d), to prevent their subsequent inadvertent use as a food for man or feed for animals.

APPLICATION RATE			
CROP & DISEASE	RATE OF RAXIL 2.6F	APPLICATION DIRECTIONS	
Wheat and Triticale: Stinking Smut, Flag Smut, Loose Smut, Early Season Septoria Disease Complex, Early Season Rhizoctonia Root Rot, Early Season Common Root Rot, Early Season Fusarium Foot Rot, Early Season, Suppression of Powdery Mildew, Early Season Suppression of Wheat Leaf Rust	0.1 fluid ounce per 100 pounds of seed	Refer to the SEED TREATMENT section on this label for the recommended method of treating seed.	
Jovered Smut, Loose Smut, Barley Stripe (suppression), Early Season Rhizoctonia Root Rot, Early Season Common Root Rot, Early Season Fusarium Foot Rot, Early Season Suppression of Powdery Mildew, Early Season Suppression of Barley Leaf Rust			
Oats: Loose Smut, Pythium Damping-off, Early Season Rhizoctonia Root Rot, Early Season Common Root Rot, Early Season Fusarium Foot Rot, Early Season Suppression of Powdery Mildew, Early Season Suppression of Crown Rust	•• •		

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, disclaimer of Warranties and Limitations of Liability.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Bayer CropScience, no claims are made to guarantee germination of carry-over seed.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. To the extent allowable under State law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: BAYER CROPSCIENCE MAKES NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION. THE REPLACEMENT OF PRODUCT.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

NET CONTENTS:

RAXIL is a registered trademark of Bayer



Bayer CropScience LP P.O. Box 12014, 2 T.W. Alexander Drive Research Triangle Park, North Carolina 27709 1-866-99BAYER (1-866-992-2937)

`AXIL 2.6F Submitted _06_/_30_/_05

ACCEPTED

.

SEP 2 1 2005

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 264-9644