RTU®-TRIFLOXYSTROBIN-METALAXYL Flowable Fungicide

EPA Reg. No. 264-1000

EPA Est.

KEEP OUT OF REACH OF CHILDREN CAUTION

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

FIRST AID

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

in case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes slight eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear: Long-sleeved shirt and long pants, socks plus shoes, chemical-resistant gloves made of any waterproof material and protective eyewear.

Engineering Controls: When handlers use closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Cover or incorporate spilled treated seeds in accordance with local regulations.

Groundwater Advisory: Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

ACCEPTED

APR | 0 20**0**6

Under the Federal Insecticide, Fungicide, and Redenticide Act, as amended, for the pesticide relationed under TP1 Tay, No. 264-1600 Page 1 of 3

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling,

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal, or by cleaning of equipment. Do not reuse empty container. Open dumping is prohibited.

STORAGE

Store in a cool dry location.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Contact Bayer CropScience or your chemical supplier for container return instructions. If container is not returned in accordance with instructions, triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restrictedentry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves made of any waterproof material, shoes plus socks and protective eyewear.

LABELING OF TREATED SEED

Seed commercially treated with RTU-Trifloxystrobin-Metalaxyl Flowable must be labeled in accordance with all applicable requirements of the Federal Seed Act. All bags containing treated seed must be labeled or tagged as follows:

- This seed has been treated with trifloxystrobin and metalaxyl.
- DO NOT USE FOR FOOD, FEED OR OIL PURPOSES.
- · Store away from feeds and other foodstuffs.

GENERAL INFORMATION

RTU-Trifloxystrobin-Metalaxyl Flowable provides seed and seedling protection against seedborne fungi causing seed decay and the soilborne pathogens, Rhizoctonia solani and Pythium spp. causing seed and seedling damping-off. RTU-Trifloxystrobin-Metalaxyl Flowable Fungicide is a ready-to-use seed treatment especially formulated for on-farm equipment for direct application, or may be applied using standard slumy or mist-type seed treatment equipment. Consult a seed treatment specialist for proper guidelines in using RTU-Trifloxystrobin-Metalaxyl Flowable Fungicide to insure uniform coverage of seed, thus ensuring seed safety and best disease protection.

USE PRECAUTIONS

When using formulations that do not contain dye, to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored to distinguish it and prevent subsequent inadvertent use as a food for man or feed for animals.

USE RESTRICTION

Do not use treated seed for food, feed, or oil processing.

Rice: Apply at a product rate of 5.7 fl. oz. of product per 100 lbs. of seed.

Soybean, Adzuki Bean, Asparagus Bean, Broad Bean (Succulent or Dry), Catjang Bean, Chinese Longbean, Field Bean, Guar Bean, Jackbean, Kidney Bean, Lablab Bean, Lima Bean (succulent or Dry), Moth Bean, (Succulent or Dry), Mung Bean, Navy Bean, Pinto Bean, Rice Bean, Runner Bean, Snap Bean, Sword Bean, Tepary Bean, Urd Bean, Wax Bean, Yardiong Bean, Blackeyed Pea (Succulent or Dry), Chickpea, Cowpea (Succulent or Dry), Crowder Pea, Dwarf Pea, Edible-Pod Pea, English Pea, Field Pea, garden Pea, Green Pea, Pigeon Pea (Succulent or Dry), Snow Pea, Southern Pea (Succulent or Dry), Sugar Snap Pea, Grain Lupin, Sweet Lupin, White Lupin, white Sweet Lupin, Lentil:

Apply at a product rate of 5.7 fl. oz, of product per 100 lbs, of seed.

ROTATIONAL CROP RESTRICTION

Areas planted with treated seed may be replanted immediately with any crop for which a tolerance exists for trifloxystrobin. Do not plant any crop without trifloxystrobin tolerances until 30 days after planting trifloxystrobin treated seed.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, disclaimer of Warranties and Limitations of Liability.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Bayer CropScience, no claims are made to guarantee germination of carry-over seed.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. To the extent allowable by State law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: BAYER CROPSCIENCE MAKES NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.


LIMITATIONS OF LIABILITY: TO THE EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

NET CONTENTS:

RTU is a registered trademark of Bayer



Bayer CropScience LP P.O. Box 12014, 2 T.W. Alexander Drive Research Triangle Park, North Carolina 27709 1-866-99BAYER (1-866-992-2937)

RTU-TRIFLOXYSTROBIN-METALAXYL FLOWABLE (PENDING) 04/04/06