

<p>ACCEPTED OCT 31 2006</p> <p>Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide Registered under EPA Reg. No. <i>264-1044</i></p>

IMIDACLOPRID-METALAXYL

Seed Treatment

For the protection of Seeds and Seedlings of Field Corn, Sweet Corn and Popcorn against Damping-Off, Seed Decay caused by *Pythium*; also for protection against Seed Corn Beetle, Seed Corn Maggots, Wireworm, White Grub, Flea Beetle (up to first true leaf), and Imported Fire Ants. For the protection of Seeds and Seedlings of Sorghum against Damping-Off, Seed Decay caused by *Pythium*; also for suppression against Wireworm, Seed Corn Maggot, true White Grub, and protection of seed against Imported Fire Ants.

For the protection of Seeds and Seedlings of Soybeans against Damping-Off, Seed Decay caused by *Pythium*; also suppression of Seed Corn Maggots and early season control of over-wintering Bean Leaf Beetles.

ACTIVE INGREDIENTS:	
Imidacloprid	25.00%
Metalaxyl	1.00%
INERT INGREDIENTS	74.00%
TOTAL	100.00%
EPA Reg. No. 264-1044	EPA Est.

KEEP OUT OF REACH OF CHILDREN

CAUTION

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material, and shoes plus socks.

Follow manufacturer's instructions for cleaning / maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds and aquatic invertebrates and toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Exposed treated seeds may be hazardous to birds. In the event that treated seeds are spilled and cannot be salvaged, completely cover them with soil. Do not contaminate water when disposing of equipment washwaters or rinsate.

PRODUCT INFORMATION

Imidacloprid-Metalaxyl may be used on seed previously treated with a full dosage of protective fungicide to give added protection against *Pythium* damping-off or seed decay.

Note: Adjust the seed drill or planter to give the required rate of seed delivery with treated seed. Regularly inspect planter plates and clean as required. At the end of each day, clean the electric-eye monitors as directed by the manufacturer.

Important: Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seed beyond planting time.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks, protective eyewear

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty containers.

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store in a cool dry location. Protect from excessive heat.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty canister into application equipment. Then dispose of empty canister in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Field Corn, Sweet Corn, Popcorn: Use at a rate of 1.5 ounces per 42 pounds of seed.

Sorghum: Use at a rate of 5.0 ounces per 100 pounds of seed.

Soybeans: Use at a rate of 4.0 oz. per 100 pounds of seed.

Mixing Instructions (1.5 oz canister)

Use as a dry mixture in the planter box as a seed treatment prior to planting. Apply Imidacloprid- Metalaxyl to seed at planting time with the canister applicator tube system.

Field corn, Sweet Corn, Popcorn: Apply at a rate of 1.5 oz. canister per 42 pounds of seed.

Sorghum: Apply at a rate of 1.5 oz. canister per 30 pounds of seed.

Soybeans: Apply at a rate of 1.5 oz. canister per 37 pounds of seed.

For best results, fill planter box with seed, add Imidacloprid- Metalaxyl through applicator tube and mix so that all seeds are covered. **DO NOT MIX WITH BARE HANDS.** Use only at the recommended rate. Lower amounts may not give desired control.

Mixing Instructions (4-pound pail)

Use as a dry mixture in the planter box as a seed treatment prior to planting.

Field Corn, Sweet Corn, Popcorn: Use two scoops full (1.5 oz.) per 42 pounds of seed.

Sorghum: Use two scoops full (1.5oz.) per 30 pounds of seed.

Soybeans: Use two scoops full (1.5 oz.) per 37 pounds of seed.

For best results, fill planter box half full of seed, add half the required amount of Imidacloprid- Metalaxyl and mix thoroughly with a stick or paddle. Add remainder of the seed and Imidacloprid- Metalaxyl and mix again so that all seeds are covered. **DO NOT MIX WITH BARE HANDS.** Use only at the recommended rate. Lower amounts may not give desired control.

NOTE: Do not graze or feed livestock on treated areas for 45 days.

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on the Imidacloprid-Metalaxyl label, or for any crop for which a tolerance exists for imidacloprid, as soon as practical following the last application. The following plant-back intervals should be observed for listed crops:

ROTATIONAL PLANT-BACK INTERVALS*				
IMMEDIATE PLANT-BACK				30-DAY PLANT-BACK
Artichoke	Cranberry	Mustard seed	Sugar beet	Cereals (Including buckwheat, millet, oats, popcorn, rice and rye) Safflower
Barley	Cucurbits	Okra	Tomatillo	
Rrassica (cole) leafy veg.	Eggplant	Pepinos	Tomato	
Canola	Ground cherry	Pepper	Triticale	
Cilantro	Leaf petiole veg.	Popcorn	Veg. Root and Tuber	
Corn, Field	Leafy veg.	Potato	Watercress	
Corn, Sweet	Legume veg. (succulent or dried, including soybean)	Rapeseed	Wheat	
Cotton		Sorghum		
		Soybean		
		Strawberry		
* Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed.				
* All other crops not listed on an Imidacloprid label, or for crops for which no tolerance for the active ingredient have been established, a 12-month plant-back interval should be observed.				

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Bayer CropScience, no claims are made to guarantee germination of carry-over seed.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: BAYER CROPSCIENCE MAKES NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.


LIMITATIONS OF LIABILITY: TO THE EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

NET CONTENTS:

PRODUCED FOR



**Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)**

IMIDACLOPRID-METALAXYL Seed Treatment (PENDING) Submitted 07/27/06, Resubmitted 08/23/06