

264-1052

2/15/2011

10/15



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

FEB 15 2011

Melvin K. Tolliver  
Bayer CropScience  
2 T.W. Alexander Drive  
P.O. Box 12014  
RTP, NC 27709

Subject: Adament 50 WG Fungicide  
EPA Reg. No. 264-1052  
Decision Number 444344

Dear Mr. Tolliver:

The Agency is in receipt of your December 2, 2010 Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call Summer Gardner-Jenkins at 703-308-9353 or you may call me at 703-308-9354.

Sincerely,

Mary L. Waller  
Product Manager (21)  
Fungicide Branch  
Registration Division (7504P)

2015

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 2-28-95



United States  
Environmental Protection Agency  
Washington, DC 20460

Registration  
 Amendment  
 Other

OPP Identifier Number

**Application for Pesticide - Section I**

1. Company/Product Number 264-1052	2. EPA Product Manager Mary Waller	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Adament 50 WG Fungicide	PM# Team 21	
5. Name and Address of Applicant (Include ZIP Code) Bayer CropScience LP P.O. Box 12014, 2 T.W. Alexander Drive Research Triangle Park, NC 27709 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

**Section - II**

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)

Notification of other revisions per PR Notice 98-10.

See attached for explanation.

**Section - III**

1. Material This Product Will Be Packaged In:						2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No				<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt.	No. per container		
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container			4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>		
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____			

**Section - IV**

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Melvin K. Tolliver	Title Registration Product Manager, Fungicides	Telephone No. (Include Area Code) (919) 549-2631
<b>Certification</b> I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment, both under applicable law.		6. Date Application Received  (Stamped) 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
2. Signature <i>Melvin K. Tolliver</i>	3. Title Registration Product Manager, Fungicides	
4. Typed Name Melvin K. Tolliver	5. Date December 2, 2010	

Adament 50 WG Fungicide  
EPA Reg. No. 264-1052

Section II - Continued

Adament 50 WG Fungicide is currently registered for use on a number of crops including pome fruit. With this 8570-1 form, one copy of annotated labeling, and two copies of final printed labeling; we are notifying the Agency that we are restricting the use of Adament 50 WG Fungicide on pome fruit from New York by adding "Not for use in New York." See page 8 of the enclosed annotated label.

This notification is consistent with the provision of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA, and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

b6  
b7C  
b7D  
b7E  
b7F  
b7G  
b7H  
b7I  
b7J  
b7K  
b7L  
b7M  
b7N  
b7O  
b7P  
b7Q  
b7R  
b7S  
b7T  
b7U  
b7V  
b7W  
b7X  
b7Y  
b7Z  
b7AA  
b7AB  
b7AC  
b7AD  
b7AE  
b7AF  
b7AG  
b7AH  
b7AI  
b7AJ  
b7AK  
b7AL  
b7AM  
b7AN  
b7AO  
b7AP  
b7AQ  
b7AR  
b7AS  
b7AT  
b7AU  
b7AV  
b7AW  
b7AX  
b7AY  
b7AZ  
b7BA  
b7BB  
b7BC  
b7BD  
b7BE  
b7BF  
b7BG  
b7BH  
b7BI  
b7BJ  
b7BK  
b7BL  
b7BM  
b7BN  
b7BO  
b7BP  
b7BQ  
b7BR  
b7BS  
b7BT  
b7BU  
b7BV  
b7BW  
b7BX  
b7BY  
b7BZ  
b7CA  
b7CB  
b7CC  
b7CD  
b7CE  
b7CF  
b7CG  
b7CH  
b7CI  
b7CJ  
b7CK  
b7CL  
b7CM  
b7CN  
b7CO  
b7CP  
b7CQ  
b7CR  
b7CS  
b7CT  
b7CU  
b7CV  
b7CW  
b7CX  
b7CY  
b7CZ  
b7DA  
b7DB  
b7DC  
b7DD  
b7DE  
b7DF  
b7DG  
b7DH  
b7DI  
b7DJ  
b7DK  
b7DL  
b7DM  
b7DN  
b7DO  
b7DP  
b7DQ  
b7DR  
b7DS  
b7DT  
b7DU  
b7DV  
b7DW  
b7DX  
b7DY  
b7DZ  
b7EA  
b7EB  
b7EC  
b7ED  
b7EE  
b7EF  
b7EG  
b7EH  
b7EI  
b7EJ  
b7EK  
b7EL  
b7EM  
b7EN  
b7EO  
b7EP  
b7EQ  
b7ER  
b7ES  
b7ET  
b7EU  
b7EV  
b7EW  
b7EX  
b7EY  
b7EZ  
b7FA  
b7FB  
b7FC  
b7FD  
b7FE  
b7FF  
b7FG  
b7FH  
b7FI  
b7FJ  
b7FK  
b7FL  
b7FM  
b7FN  
b7FO  
b7FP  
b7FQ  
b7FR  
b7FS  
b7FT  
b7FU  
b7FV  
b7FW  
b7FX  
b7FY  
b7FZ  
b7GA  
b7GB  
b7GC  
b7GD  
b7GE  
b7GF  
b7GG  
b7GH  
b7GI  
b7GJ  
b7GK  
b7GL  
b7GM  
b7GN  
b7GO  
b7GP  
b7GQ  
b7GR  
b7GS  
b7GT  
b7GU  
b7GV  
b7GW  
b7GX  
b7GY  
b7GZ  
b7HA  
b7HB  
b7HC  
b7HD  
b7HE  
b7HF  
b7HG  
b7HH  
b7HI  
b7HJ  
b7HK  
b7HL  
b7HM  
b7HN  
b7HO  
b7HP  
b7HQ  
b7HR  
b7HS  
b7HT  
b7HU  
b7HV  
b7HW  
b7HX  
b7HY  
b7HZ  
b7IA  
b7IB  
b7IC  
b7ID  
b7IE  
b7IF  
b7IG  
b7IH  
b7II  
b7IJ  
b7IK  
b7IL  
b7IM  
b7IN  
b7IO  
b7IP  
b7IQ  
b7IR  
b7IS  
b7IT  
b7IU  
b7IV  
b7IW  
b7IX  
b7IY  
b7IZ  
b7JA  
b7JB  
b7JC  
b7JD  
b7JE  
b7JF  
b7JG  
b7JH  
b7JI  
b7JJ  
b7JK  
b7JL  
b7JM  
b7JN  
b7JO  
b7JP  
b7JQ  
b7JR  
b7JS  
b7JT  
b7JU  
b7JV  
b7JW  
b7JX  
b7JY  
b7JZ  
b7KA  
b7KB  
b7KC  
b7KD  
b7KE  
b7KF  
b7KG  
b7KH  
b7KI  
b7KJ  
b7KK  
b7KL  
b7KM  
b7KN  
b7KO  
b7KP  
b7KQ  
b7KR  
b7KS  
b7KT  
b7KU  
b7KV  
b7KW  
b7KX  
b7KY  
b7KZ  
b7LA  
b7LB  
b7LC  
b7LD  
b7LE  
b7LF  
b7LG  
b7LH  
b7LI  
b7LJ  
b7LK  
b7LL  
b7LM  
b7LN  
b7LO  
b7LP  
b7LQ  
b7LR  
b7LS  
b7LT  
b7LU  
b7LV  
b7LW  
b7LX  
b7LY  
b7LZ  
b7MA  
b7MB  
b7MC  
b7MD  
b7ME  
b7MF  
b7MG  
b7MH  
b7MI  
b7MJ  
b7MK  
b7ML  
b7MN  
b7MO  
b7MP  
b7MQ  
b7MR  
b7MS  
b7MT  
b7MU  
b7MV  
b7MW  
b7MX  
b7MY  
b7MZ  
b7NA  
b7NB  
b7NC  
b7ND  
b7NE  
b7NF  
b7NG  
b7NH  
b7NI  
b7NJ  
b7NK  
b7NL  
b7NM  
b7NN  
b7NO  
b7NP  
b7NQ  
b7NR  
b7NS  
b7NT  
b7NU  
b7NV  
b7NW  
b7NX  
b7NY  
b7NZ  
b7OA  
b7OB  
b7OC  
b7OD  
b7OE  
b7OF  
b7OG  
b7OH  
b7OI  
b7OJ  
b7OK  
b7OL  
b7OM  
b7ON  
b7OO  
b7OP  
b7OQ  
b7OR  
b7OS  
b7OT  
b7OU  
b7OV  
b7OW  
b7OX  
b7OY  
b7OZ  
b7PA  
b7PB  
b7PC  
b7PD  
b7PE  
b7PF  
b7PG  
b7PH  
b7PI  
b7PJ  
b7PK  
b7PL  
b7PM  
b7PN  
b7PO  
b7PP  
b7PQ  
b7PR  
b7PS  
b7PT  
b7PU  
b7PV  
b7PW  
b7PX  
b7PY  
b7PZ  
b7QA  
b7QB  
b7QC  
b7QD  
b7QE  
b7QF  
b7QG  
b7QH  
b7QI  
b7QJ  
b7QK  
b7QL  
b7QM  
b7QN  
b7QO  
b7QP  
b7QQ  
b7QR  
b7QS  
b7QT  
b7QU  
b7QV  
b7QW  
b7QX  
b7QY  
b7QZ  
b7RA  
b7RB  
b7RC  
b7RD  
b7RE  
b7RF  
b7RG  
b7RH  
b7RI  
b7RJ  
b7RK  
b7RL  
b7RM  
b7RN  
b7RO  
b7RP  
b7RQ  
b7RR  
b7RS  
b7RT  
b7RU  
b7RV  
b7RW  
b7RX  
b7RY  
b7RZ  
b7SA  
b7SB  
b7SC  
b7SD  
b7SE  
b7SF  
b7SG  
b7SH  
b7SI  
b7SJ  
b7SK  
b7SL  
b7SM  
b7SN  
b7SO  
b7SP  
b7SQ  
b7SR  
b7SS  
b7ST  
b7SU  
b7SV  
b7SW  
b7SX  
b7SY  
b7SZ  
b7TA  
b7TB  
b7TC  
b7TD  
b7TE  
b7TF  
b7TG  
b7TH  
b7TI  
b7TJ  
b7TK  
b7TL  
b7TM  
b7TN  
b7TO  
b7TP  
b7TQ  
b7TR  
b7TS  
b7TT  
b7TU  
b7TV  
b7TW  
b7TX  
b7TY  
b7TZ  
b7UA  
b7UB  
b7UC  
b7UD  
b7UE  
b7UF  
b7UG  
b7UH  
b7UI  
b7UJ  
b7UK  
b7UL  
b7UM  
b7UN  
b7UO  
b7UP  
b7UQ  
b7UR  
b7US  
b7UT  
b7UU  
b7UV  
b7UW  
b7UX  
b7UY  
b7UZ  
b7VA  
b7VB  
b7VC  
b7VD  
b7VE  
b7VF  
b7VG  
b7VH  
b7VI  
b7VJ  
b7VK  
b7VL  
b7VM  
b7VN  
b7VO  
b7VP  
b7VQ  
b7VR  
b7VS  
b7VT  
b7VU  
b7VV  
b7VW  
b7VX  
b7VY  
b7VZ  
b7WA  
b7WB  
b7WC  
b7WD  
b7WE  
b7WF  
b7WG  
b7WH  
b7WI  
b7WJ  
b7WK  
b7WL  
b7WM  
b7WN  
b7WO  
b7WP  
b7WQ  
b7WR  
b7WS  
b7WT  
b7WU  
b7WV  
b7WW  
b7WX  
b7WY  
b7WZ  
b7XA  
b7XB  
b7XC  
b7XD  
b7XE  
b7XF  
b7XG  
b7XH  
b7XI  
b7XJ  
b7XK  
b7XL  
b7XM  
b7XN  
b7XO  
b7XP  
b7XQ  
b7XR  
b7XS  
b7XT  
b7XU  
b7XV  
b7XW  
b7XZ  
b7YA  
b7YB  
b7YC  
b7YD  
b7YE  
b7YF  
b7YG  
b7YH  
b7YI  
b7YJ  
b7YK  
b7YL  
b7YM  
b7YN  
b7YO  
b7YP  
b7YQ  
b7YR  
b7YS  
b7YT  
b7YU  
b7YV  
b7YW  
b7YX  
b7YY  
b7YZ  
b7ZA  
b7ZB  
b7ZC  
b7ZD  
b7ZE  
b7ZF  
b7ZG  
b7ZH  
b7ZI  
b7ZJ  
b7ZK  
b7ZL  
b7ZM  
b7ZN  
b7ZO  
b7ZP  
b7ZQ  
b7ZR  
b7ZS  
b7ZT  
b7ZU  
b7ZV  
b7ZW  
b7ZX  
b7ZY  
b7ZZ

40915

S 887501

# Bayer CropScience



Document Processing Desk (NOTIF)  
Office of Pesticide Programs (7504P)  
U.S. Environmental Protection Agency  
Room S4900  
One Potomac Yard  
2777 S. Crystal Drive  
Arlington, VA 22202

December 2, 2010

Bayer CropScience  
2 T.W. Alexander Drive  
P. O. Box 12014  
RTP, NC 27709  
Phone: (919) 549-2000

Subject: Adamaent 50 WG Fungicide, EPA Reg. No. 264-1052  
Notification of Other Revisions per PR Notice 98-10

Dear Sir or Madam:

Adament 50 WG Fungicide is currently registered for use on a number of crops including pome fruit. As described in Section II of the enclose 8570-1 form, we are notifying the Agency that we are restricting the use of Adament 50 WG Fungicide on pome fruit from New York by adding "Not for use in New York."

Since this is a notification, no PRIA fee is required.

If you have any questions or need additional information, please contact me by phone at (919) 549-2631 or by e-mail at [mel.tolliver@bayer.com](mailto:mel.tolliver@bayer.com).


Sincerely,

*Melvin K. Tolliver*

Melvin K. Tolliver  
Registration Product Manager, Fungicides

Enclosures:

1. EPA application form 8570-1
2. Draft annotated labeling
3. Final Printed labeling (two copies)



# ADAMENT® 50 WG Fungicide

For control of certain diseases on almonds, grapes, pistachio, pome fruit, stone fruit and tree nuts

Active Ingredients:	
Tebuconazole.....	25.0%
Trifloxystrobin.....	25.0%
Other Ingredients:.....	50.0%
	Total 100.0%

EPA Reg. No. 264-1052

EPA Est. No.

## KEEP OUT OF REACH OF CHILDREN CAUTION

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577  
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

### FIRST AID

IF IN EYES:	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
IF SWALLOWED:	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
IF ON SKIN:	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
IF INHALED:	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>
In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.	
NOTE TO PHYSICIAN: No specific antidote. Treat Symptomatically.	

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Causes moderate eye irritation. Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

#### Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical resistant gloves made of any waterproof material

### NOTIFICATION

FEB 15 2011

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**ENGINEERING CONTROL STATEMENTS**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water or rinsate.

**Ground Water Advisory:**

Several trifloxystrobin degradates have properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

**Surface Water Advisory:**

This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.  
Do not use this product until you have read the entire label.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

#### Pesticide Storage

Store in a cool, dry place and in such a manner as to prevent cross-contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Dispose of pesticide as directed. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer CropScience Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer CropScience Emergency Response telephone number is 1-800-334-7577.

#### Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

#### Container Disposal

Non-refillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Offer for recycling, if available. If not recycled, then puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### GENERAL INFORMATION

ADAMENT® 50 WG Fungicide is a broad spectrum fungicide for the control of certain diseases of almonds, grapes, pistachio, pome fruit, stone fruit and tree nuts. ADAMENT 50 WG Fungicide works by interfering with both energy and cell membrane production in plant pathogenic fungi.

UNDER CERTAIN CONDITIONS CONDUCIVE TO EXTENDED INFECTION PERIODS, ADDITIONAL FUNGICIDE APPLICATIONS BEYOND THE NUMBER ALLOWED BY THIS LABEL MAY BE NEEDED. UNDER THESE CONDITIONS, USE ANOTHER FUNGICIDE REGISTERED FOR THE CROP/DISEASE.

DO NOT APPLY ADAMENT TO CONCORD GRAPES OR CROP INJURY MAY OCCUR.

### RESISTANCE MANAGEMENT

The active ingredients in ADAMENT 50 WG Fungicide belong to two different chemistry classes.

Tebuconazole belongs to the DMI (Group 3) class of chemistry which exhibits no known cross-resistance to other chemical classes. Tebuconazole may exhibit cross-resistance to other Group 3 fungicides, such as propiconazole and myclobutanil.

Trifloxystrobin belongs to the QoI (Group 11) class of chemistry which exhibits no known cross-resistance to other chemical classes. Trifloxystrobin does exhibit cross-resistance to other Group 11 fungicides, such as azoxystrobin, pyraclostrobin, and kresoxim-methyl. The NA-QoI Working Group has established the following general guidelines for the maximum number of applications of a Group 11-containing fungicide. In addition to that, the maximum number of applications may be restricted to a specific limit on a particular crop (see crop-specific recommendations). Follow the specific crop directions that limit the total number of sprays on a crop and the required alternations with fungicides from other resistance management groups. In situations requiring multiple fungicide sprays, develop season-long spray programs for Group 11-containing fungicides. In programs in which pre-mixes of a Group 11 fungicide with a fungicide of another group are utilized, the number of Group 11-containing applications should be no more than about 1/2 of the total number of fungicide applications per season.

Fungal pathogens are known to develop resistance to products with the same mode of action when used repeatedly. Because resistance development cannot be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include rotating and/or tank mixing with products having different modes of action or limiting the total number of applications per season. Bayer CropScience encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

### SPRAY EQUIPMENT

Thorough coverage is necessary to provide good disease control. Applications using sufficient water volume to provide thorough and uniform coverage generally provide the most effective disease control. For ground application equipment, use a minimum of 10 gal./A. For aerial application equipment, use a minimum of 5 gal./A.

**Broadcast Ground Sprayers**

Equip sprayers with nozzles that provide accurate and uniform application. Be certain that nozzles are the same size and uniformly spaced across the boom. Calibrate the sprayer before use.

Use a pump with the capacity to: (1) maintain a minimum of 35 psi at nozzles, and (2) provide sufficient agitation in the tank to keep the mixture in suspension; this requires recirculation of 10% of the tank volume per minute. Use jet agitators or a liquid sparge tube for vigorous agitation.

Use screens to protect the pump and to prevent nozzles from clogging. Screens placed on the suction side of the pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh screens at the nozzles. Check nozzle manufacturer's recommendations.

For information on spray equipment and calibration, consult sprayer manufacturer's and/or state recommendations. For specific local spray schedules, consult the current state agricultural experiment station recommendations.

**AERIAL APPLICATION**

Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply directly to humans or animals. Not registered for aerial application in New York State.

**CHEMIGATION**

Do not apply this product through any type of irrigation system.

**MIXING PROCEDURES**

Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. Do not let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

**ADAMENT 50 WG Fungicide Alone:** Add approximately 1/2 of the required amount of water to the mix tank. With the agitator running, add the ADAMENT 50 WG Fungicide to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after the ADAMENT 50 WG Fungicide has completely and uniformly dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

**ADAMENT 50 WG Fungicide + Tank Mix Partners:** Add approximately 1/2 of the required amount of water to the mix tank. Start the agitator running before adding any tank mix partners. In general, tank mix partners should be added in this order: products packaged in watersoluble packaging\*, wettable powders, wettable granules (dry flowables), liquid flowables such as ADAMENT 50 WG Fungicide, liquids, and emulsifiable concentrates. Always allow each tank mix partner to become fully and uniformly dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all of the mixture has been applied.

\* Note: When using ADAMENT 50 WG Fungicide in tank mixtures, all products in water-soluble packaging should be added to the tank before any other tank mix partner, including ADAMENT 50 WG Fungicide. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank-mix partner to the tank. If using ADAMENT 50 WG Fungicide in a tank mixture, observe all directions for use, crop/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix partner product label. No label dosage rate may be exceeded, and the most restrictive label precautions and limitations must be followed. This product must not be mixed with any product which prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

ADAMENT 50 WG Fungicide is compatible with most insecticide, fungicide, and foliar nutrient products. However, the physical compatibility of ADAMENT 50 WG Fungicide with tank mix partners should be tested before use. To determine the physical compatibility of ADAMENT 50 WG Fungicide with other products, use a jar test, as described below.

Using a quart jar, add the proportionate amounts of the products to 1 quart of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

The crop safety of all potential tank mixes, including additives and other pesticides, on all crops has not been tested. Before applying any tank mixture not specifically recommended on this label, the safety to the target crop should be confirmed. To test for crop safety, apply the tank mixture to the target crop in a small area and in accordance with label instructions for the target crop.

**RECOMMENDATIONS TO AVOID SPRAY DRIFT**

Do not make applications when conditions favor drift beyond the target application area. When drift may be a problem, take measures to reduce drift, including:

1. Do not spray if wind speeds are or become excessive. Do not spray if wind speed is 15 mph or greater. If nontarget crops are located downwind, use caution when spraying if wind is present. Do not spray if winds are gusty.
2. Use caution when conditions are favorable for drift (high temperatures, drought, low relative humidity).



9/15

3. Do not apply when a temperature inversion exists. If inversion conditions are suspected, consult with local weather services before making an application.

**RAINFASTNESS**

ADAMENT 50 WG Fungicide is rainfast when dry, generally in 2 hours, and dependent on local weather conditions, such as humidity and wind.

**USE DIRECTIONS FOR SPECIFIC CROPS**

ADAMENT 50 WG Fungicide provides control or suppression of several important diseases of almonds, grapes, pistachio, pome fruit, stone fruit and tree nuts. When reference is made to disease suppression, suppression can mean either erratic control from good to fair, or consistent control at a level below that obtained with the best commercial disease control products.

<b>Almonds</b>		
<b>Disease Control</b>	<b>Rate (oz./acre)</b>	<b>Application Timing</b>
Brown Rot Blossom Blight <i>(Monilinia laxa, M. fructicola)</i>  Jacket Rot, Green Fruit Rot <i>(Botrytis cinerea)</i>  Alternaria <i>(Alternaria alternata)</i>  Anthracnose <i>(Colletotrichum acutatum)</i>  Shot Hole <i>(Wilsonomyces carpophilus)</i>  Powdery Mildew <i>(Podosphaera tridactyla,</i> <i>Sphaerotheca pannosa)</i>  Scab <i>(Cladosporium carpophilum)</i>  Rust <i>(Tranzschelia discolor)</i>	4 to 8	Begin applications when conditions are favorable for disease but before infection. Apply on a 7- to 14-day spray schedule.  Use higher rates and shorter intervals when disease pressure is severe.  Refer to University and/or extension guidelines for best application timings.
<b>Restrictions:</b> Do not apply more than 32 oz. of ADAMENT 50 WG FUNGICIDE per acre per season. Do not make more than 4 total applications of ADAMENT 50 WG FUNGICIDE per season. Do not apply ADAMENT 50 WG FUNGICIDE within 60 days of harvest or after hull split. Do not cut cover crops in treated areas for feed, or allow livestock to graze treated areas. ADAMENT 50 WG FUNGICIDE is a mixture of a Group 11 and a Group 3 fungicide. To limit the potential for the development of resistance, do not apply more than 2 sequential applications of ADAMENT 50 WG FUNGICIDE or other Group 11-containing fungicides.		

Grapes: DO NOT APPLY ADAMENT TO CONCORD GRAPES OR CROP INJURY MAY OCCUR.		
Disease Control	Rate (oz./acre)	Application Timing
Powdery Mildew ( <i>Uncinula necator</i> )	3	Begin applications when conditions are favorable for disease but before infection. Apply on a 14-day spray schedule.
	4	Begin applications when conditions are favorable for disease but before infection. Apply on a 21-day spray schedule.
Botrytis Bunch Rot ( <i>Botrytis cinerea</i> )	6	Refer to timings listed above for grape powdery mildew. Research data show a trend toward better control if fungicides are applied at bloom, pre-bunch closure, and veraison. Continue on a 14- to 21-day spray schedule.  Use shorter intervals when disease pressure is severe.
Phomopsis Cane & Leaf Spot ( <i>Phomopsis viticola</i> )	6	Begin applications at bud break and continue up to bunch closure on a 14- to 21-day spray schedule.  Use shorter intervals when disease pressure is severe.
Black Rot ( <i>Guignardia bidwellii</i> )	4	Begin applications when conditions are favorable for disease but before infection. Apply on a 10- to 14-day spray schedule.  Use shorter intervals when disease pressure is severe.
Disease Suppression	Rate (oz./acre)	Application Timing
Botrytis Bunch Rot ( <i>Botrytis cinerea</i> )  Phomopsis Cane & Leaf Spot ( <i>Phomopsis viticola</i> )	4	Refer to timings listed above for grape powdery mildew.  Use shorter intervals when disease pressure is severe.
Downy Mildew ( <i>Plasmopara viticola</i> )	7.2	Begin applications when conditions are favorable for disease but before infection. Apply on a 10- to 14-day spray schedule.  Use shorter intervals when disease pressure is severe.
<b>Restrictions:</b> Do not apply more than 48 oz. of ADAMENT 50 WG FUNGICIDE per acre per season. Do not make more than 6 total applications of ADAMENT 50 WG Fungicide per season. Do not apply ADAMENT 50 WG Fungicide within 14 days of harvest. ADAMENT 50 WG Fungicide is a mixture of a Group 11 and a Group 3 fungicide. To limit the potential for the development of resistance, do not apply more than 2 sequential applications of ADAMENT 50 WG FUNGICIDE or other Group 11-containing fungicides.		

Pistachios		
Disease Control	Rate (oz./acre)	Application Timing
Blossom & Shoot Blight <i>(Botrytis cinerea)</i>  Alternaria Late Blight <i>(Alternaria alternata)</i>	4 to 8	Begin applications when conditions are favorable for disease but before infection. Apply on a 14- to 21-day spray schedule.  Use higher rates and shorter intervals when disease pressure is severe.
Botryosphaeria Panicle & Shoot Blight <i>(Botryosphaeria dothidea)</i>  Septoria Leaf Spot <i>(Septoria pistaciarum)</i>	4 to 8	Begin applications when conditions are favorable for disease but before infection. Apply on a 14- to 21-day spray schedule.  Use higher rates and shorter intervals when disease pressure is severe.
<b>Restrictions:</b> Do not apply more than 24 oz. of ADAMENT 50 WG FUNGICIDE per acre per season. Do not make more than 4 total applications of ADAMENT 50 WG FUNGICIDE per season. Do not apply ADAMENT 50 WG FUNGICIDE within 35 days of harvest or after hull split. Do not cut cover crops in treated areas for feed, or allow livestock to graze treated areas. ADAMENT 50 WG FUNGICIDE is a mixture of a Group 11 and a Group 3 fungicide. To limit the potential for the development of resistance, do not apply more than 2 sequential applications of ADAMENT 50 WG FUNGICIDE or other Group 11-containing fungicides.		

Pome Fruits: (Apples, Pears, Crabapples, Loquat, Mayhaw, Quince)		Not for use in New York
Disease Control	Rate (oz./acre)	Application Timing
Scab ( <i>Venturia</i> spp.)	Preventive 4	Begin applications at green tip and continue as needed on a 7- to 10-day spray schedule.  Use higher rates and shorter intervals when disease pressure is severe.
	Post-Infection 5	Refer to University and/or extension guidelines for best application timings.
Cedar-Apple Rust ( <i>Gymnosporangium juniperi-virginianae</i> )	4 to 5	Begin applications at green tip and continue as needed on a 7- to 10-day spray schedule.  Use higher rates and shorter intervals when disease pressure is severe.
Powdery Mildew ( <i>Podosphaera leucotricha</i> )	4 to 5	Begin applications at green tip and continue as needed on a 10- to 14-day spray schedule.  Use higher rates and shorter intervals when disease pressure is severe.
Sooty Blotch ( <i>Gloeodes pomigena</i> )  Fly Speck ( <i>Schizothyrium pomi</i> )	4 to 5	Begin applications when conditions are favorable for disease but before infection. Apply on a 10- to 14-day spray schedule.  Use higher rates and shorter intervals when disease pressure is severe.
Disease Suppression	Rate (oz./acre)	Application Timing
Bitter Rot ( <i>Glomerella cingulata</i> )	6	Begin applications when conditions are favorable for disease but before infection. Apply ADAMENT 50 WG FUNGICIDE solo at the specified rate or use a tank mix of ADAMENT 50 WG FUNGICIDE with 1.2 lbs. active ingredient of Captan per acre.  Use higher rates and shorter intervals when disease pressure is severe.  Continue applications as needed on a 10- to 14-day spray schedule.
White Rot ( <i>Botryosphaeria dothidea</i> )	Tank mix with CAPTAN: 3	
<b>Restrictions:</b> Do not apply more than 22 oz. of ADAMENT 50 WG FUNGICIDE per acre per season. Do not make more than 4 total applications of ADAMENT 50 WG FUNGICIDE per season. Do not apply ADAMENT 50 WG FUNGICIDE within 75 days of harvest. ADAMENT 50 WG FUNGICIDE is a mixture of a Group 11 and a Group 3 fungicide. To limit the potential for the development of resistance, do not apply more than 2 sequential applications of ADAMENT 50 WG FUNGICIDE or other fungicide containing either a Group 3 or Group 11 fungicide.		

Stone Fruit (apricot, cherry, nectarine, peach, plum, plumcot, prune [fresh])		
Disease Control	Rate (oz./acre)	Application Timing
Brown Rot Blossom Blight <i>(Monilinia laxa, M. fructicola)</i>  Jacket Rot, Green Fruit Rot <i>(Botrytis cinerea)</i>  Shot Hole <i>(Wilsonomyces carpophilus)</i>  Alternaria <i>(Alternaria alternata)</i>  Anthracnose <i>(Colletotrichum acutatum)</i>  Powdery Mildew <i>(Podosphaera spp., Sphaerotheca pannosa)</i>  Rusty Spot <i>(Podosphaera leucotricha)</i>  Scab <i>(Cladosporium carpophilum)</i>  Cherry Leaf Spot <i>(Blumeriella jaapii)</i>  Fruit Rot <i>(Monilinia fructicola)</i>  Rust <i>(Tranzschelia discolor)</i>	4 to 8	Begin applications when conditions are favorable for disease but before infection. Apply on a 7- to 14-day spray schedule.  Use higher rates and shorter intervals when disease pressure is severe.  Refer to University and/or extension guidelines for best application timings.
<b>Restrictions:</b> Do not apply more than 32 oz. of ADAMENT 50 WG Fungicide per acre per season. Do not make more than 4 total applications of ADAMENT 50 WG Fungicide per season. Do not apply ADAMENT 50 WG Fungicide within 1 day of harvest. ADAMENT 50 WG Fungicide is a mixture of a Group 11 and a Group 3 fungicide. To limit the potential for the development of resistance, do not apply more than 2 sequential applications of ADAMENT 50 WG Fungicide or other Group 11 fungicides.		

Tree Nuts: beechnuts, brazil nuts, butternuts, cashews, chinquapins, filberts, hickory nuts, macadamia nuts, pecans, walnuts (See Specific Use Directions for almonds and pistachios)		
Disease Control	Rate (oz./acre)	Application Timing
Jacket Rot, Green Fruit Rot, Blossom & Shoot Blight ( <i>Botrytis cinerea</i> )  Alternaria ( <i>Alternaria alternata</i> )  Anthracnose ( <i>Colletotrichum acutatum</i> , <i>Glomerella cingulata</i> )  Shot Hole ( <i>Wilsonomyces carpophilus</i> )  Powdery Mildew ( <i>Podosphaera tridactyla</i> , <i>Sphaerotheca pannosa</i> )  Scab ( <i>Cladosporium carpophilum</i> , <i>Cladosporium caryigenum</i> )  Eastern Filbert Blight ( <i>Anisogramma anomala</i> )  Rust ( <i>Tranzschelia discolor</i> )	4 to 8	Begin applications when conditions are favorable for disease but before infection. Apply on a 7- to 14-day spray schedule.  Use higher rates and shorter intervals when disease pressure is severe.  Refer to University and/or extension guidelines for best application timings.
Botryosphaeria Panicle & Shoot Blight ( <i>Botryosphaeria dothidea</i> )  Septoria Leaf Spot ( <i>Septoria pistaciarum</i> )	4 to 8	Begin applications when conditions are favorable for disease but before infection. Apply on a 14- to 21-day spray schedule.  Use higher rates and shorter intervals when disease pressure is severe.
<b>Restrictions:</b> Do not apply more than 32 oz. of ADAMENT 50 WG FUNGICIDE per acre per season. Do not make more than 4 total applications of ADAMENT 50 WG FUNGICIDE per season. Do not apply ADAMENT 50 WG FUNGICIDE within 60 days of harvest or after hull split. Do not cut cover crops in treated areas for feed, or allow livestock to graze treated areas. ADAMENT 50 WG FUNGICIDE is a mixture of a Group 11 and a Group 3 fungicide. To limit the potential for the development of resistance, do not apply more than 2 sequential applications of ADAMENT 50 WG FUNGICIDE or other Group 11 -containing fungicides.		

**ROTATIONAL RESTRICTIONS**

Treated areas may be replanted immediately following harvest with any crop listed on this label. For crops not listed on this label, do not plant back within 120 days of last application.

15 of 15

**IMPORTANT: READ BEFORE USE**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

**NET CONTENTS: 40 OZ**

**PRODUCED FOR**



**Bayer CropScience**

Bayer CropScience LP  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
1-866-99BAYER (1-866-992-2937)

ADAMENT 50 WG Fungicide (MASTER) Approved 10/18/10, Notification 12/02/10