



**OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION**

WASHINGTON, D.C. 20460

October 09, 2025

**SENT BY EMAIL**

Megan Mader  
megan.mader1@bayer.com  
BAYER CROPSCIENCE LP

Subject: Labeling Notification per Pesticide Registration Notice (PRN) 98-10 - Revision to label  
regarding soybeans  
Product Name: LUNA PRIVILEGE  
Admin Number: 264-1078  
EPA Receipt Date: 06/19/2025  
Action Case Number: 00660544

Dear Megan Mader:

The U.S. Environmental Protection Agency is in receipt of your application for notification under Pesticide Registration Notice 98-10 for the above referenced product. The EPA has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The labeling submitted with this application has been stamped "Notification" and will be placed in our records.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

If you have questions, please contact Julie Kelleher via email at [kelleher.julie@epa.gov](mailto:kelleher.julie@epa.gov).

Sincerely,

*Kable Bo Davis*

Kable Bo Davis, Senior Advisor  
FHB, RD  
Office of Pesticide Programs

# LUNA<sup>®</sup> PRIVILEGE

**ABN: Velum<sup>®</sup> Prime, Velum<sup>®</sup> 500, Velum<sup>®</sup> One,  
FLUOPYRAM 500 ST; Velum<sup>®</sup>**

**Broad-spectrum fungicide and nematicide for use as a foliar or soil treatment for suppression of listed crop diseases and suppression of plant pathogenic nematodes.**

**For uses on:** Artichoke, (Globe); Brassica Head and Stem Vegetables (Group 5-16); Brassica Leafy Greens (Subgroup 4-16B) (except watercress); Bulb vegetables (Group 3-07); Carrot; Celtuce; Cereals (Groups 15 and 16) (except rice); Citrus (Group 10-10); Coffee; Corn; Cotton (subgroup 20C); Cucurbits (Group 9); Fennel; Florence (fresh leaves and stalks); Fruiting Vegetables (Group 8-10); Ginseng; Grapes and Small vines (except fuzzy kiwifruit) (Subgroup 13-07F); Herb (Subgroup 19A); Hops; Kohlrabi; Leafy greens (Subgroup 4-16A); Leaf petiole vegetables (Subgroup 22B); Legume Vegetables (Group 6-22) (except Dried Shelled Pulse Pea Legume Vegetables subgroup 6-22F); Mint (spearmint and peppermint, fresh and dried leaves); Papaya; Peanut; Pome fruit (Group 11-10); Potato and other root, tuberous and corm vegetables (except sugarbeet) (Subgroups 1B and 1C); Rapeseed (Subgroup 20A); Small Berries (canberries and bushberries) (Subgroups 13-07A and 13-07B); Sorghum; Soybean; Spice (Group 26); Stone Fruits (Group 12-12); Strawberry and other low-growing berries, except cranberry (Subgroup 13-07G); Sugarbeet; Sunflower (Subgroup 20B); Tobacco; Tree Nuts (Group 14-12).

## ACTIVE INGREDIENT:

Fluopyram: *N*-[2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl]-2-(trifluoromethyl)benzamide\* ..... 41.5%

**OTHER INGREDIENTS:**..... 58.5%

**TOTAL:** ..... **100.0%**

**Contains 4.16 lbs FLUOPYRAM per gallon**

\*(CAS Number 658066-35-4)

**EPA Reg. No. 264-1078**

**EPA Est. No.**

SUSPENSION CONCENTRATE

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

[See additional precautionary statements and directions for use on label.]

## NOTIFICATION

264-1078

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

10/09/2025

## FIRST AID

<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• <b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• <b>DO NOT</b> give anything by mouth to an unconscious person.</li></ul>
<b>IF ON SKIN:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<p><b>For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577.</b> In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	
<b>NOTE TO PHYSICIAN:</b> Treat Symptomatically.	

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

#### Personal Protective Equipment (PPE):

Applicators and other handlers must wear long-sleeved shirt and long pants, shoes plus socks, and chemical-resistant (such as natural rubber) gloves.

#### User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

## Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

#### Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- User should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

For terrestrial uses: **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment wash water or rinsate.

### Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of Fluopyram. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff.

### Ground Water Advisory

This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

## **DIRECTIONS FOR USE**

### **STOP - READ THE LABEL BEFORE USE**

**It is a violation of federal law to use this product in a manner inconsistent with its labeling.**

**Read entire label before using this product.**

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**Not for sale, distribution, or use in Nassau and Suffolk counties, New York except as permitted under FIFRA 24(c), Special Local Need registration.**

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the **restricted-entry interval (REI) of 12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls over long-sleeved shirt and long pants, socks and shoes, and chemical-resistant gloves made of any waterproof material such as natural rubber  $\geq$  14 mils.

### **PRODUCT INFORMATION**

LUNA® PRIVILEGE is a broad-spectrum fungicide and nematicide for use as a foliar or soil treatment for suppression of listed crop diseases and suppression of plant pathogenic nematodes.

### **LABELED USES**

**Foliar uses:** Artichoke, (Globe); Brassica Head and Stem Vegetables (Group 5-16); Brassica Leafy Greens (Subgroup 4-16B) (except watercress); Bulb vegetables (Group 3-07); Carrot; Celtuce; Cereals (Groups 15 and 16) (except rice); Citrus (Group 10-10); Corn; Cotton (subgroup 20C); Cucurbits (Group 9); Fennel; Florence (fresh leaves and stalks); Fruiting Vegetables (Group 8-10); Ginseng; Grapes and Small vines (except fuzzy kiwifruit) (Subgroup 13-07F); Herb (Subgroup 19A); Hops; Kohlrabi; Leafy greens (Subgroup 4-16A); Leaf petiole vegetables (Subgroup 22B); Legume Vegetables (Group 6-22) (except Dried Shelled Pulse Pea Legume Vegetables subgroup 6-22F); Mint (spearmint and peppermint, fresh and dried leaves); Papaya; Peanut; Pome fruit (Group 11-10); Potato and other root, tuberous and corm vegetables (except sugarbeet) (Subgroups 1B and 1C); Rapeseed (Subgroup 20A); Small Berries (canberries and bushberries) (Subgroups 13-07A and 13-07B); Sorghum; Soybean; Spice (Group 26); Stone Fruits (Group 12-12); Strawberry and other low-growing berries, except cranberry (Subgroup 13-07G); Sugarbeet; Sunflower (Subgroup 20B); Tobacco; Tree Nuts (Group 14-12).

**Soil uses:** Artichoke, (Globe); Brassica Head and Stem Vegetables (Group 5-16); Brassica Leafy Greens (Subgroup 4-16B) (except watercress); Bulb vegetables (Group 3-07); Carrot; Celtnce; Cereals (Groups 15 and 16) (except rice); Citrus (Group 10-10); Coffee; Corn; Cotton (subgroup 20C); Cucurbits (Group 9); Fennel; Florence (fresh leaves and stalks); Fruiting Vegetables (Group 8-10); Ginseng; Grapes and Small vines (except fuzzy kiwifruit) (Subgroup 13-07F); Herb (Subgroup 19A); Hops; Kohlrabi; Leafy greens (Subgroup 4-16A); Leaf petiole vegetables (Subgroup 22B); Legume Vegetables (Group 6-22) (except Dried Shelled Pulse Pea Legume Vegetables subgroup 6-22F); Mint (spearmint and peppermint, fresh and dried leaves); Peanut; Pome fruit (Group 11-10); Potato and other root, tuberous and corm vegetables (except sugarbeet) (Subgroups 1B and 1C); Rapeseed (Subgroup 20A); Small Berries (canberries and bushberries) (Subgroups 13-07A and 13-07B); Sorghum; Soybean; Spice (Group 26); Stone Fruits (Group 12-12); Strawberry and other low-growing berries, except cranberry (Subgroup 13-07G); Sugarbeet; Sunflower (Subgroup 20B); Tobacco; Tree Nuts (Group 14-12).

## RESISTANCE MANAGEMENT

For resistance management, please note that LUNA PRIVILEGE contains a Group 7 fungicide. Any fungal population may contain individuals naturally resistant to LUNA PRIVILEGE and other Group 7 fungicides. A gradual or total loss of pest control may occur over time if these (fungicides) are used repeatedly in the same fields. Appropriate resistance-management strategies must be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of LUNA PRIVILEGE or other Group 7 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal/bacterial populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Bayer CropScience at 1-866-99BAYER (1-866-992-2937). You can also contact your pesticide distributor or university extension specialist to report resistance.

IPM: It is advised that applications of fungicides be integrated into an overall disease and pest management program, following cultural practices known to reduce disease development. Consult your local extension specialist, certified crop advisor and/or manufacturer representative for additional IPM strategies established for your area. This product may be used in Agricultural Extension advisory (disease forecasting or risk assessment) programs, which specify application timing based on environmental factors favorable for disease development.

## **MANDATORY SPRAY DRIFT**

### **Aerial Applications**

- **DO NOT** release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- For all applications, applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- **DO NOT** apply when wind speeds exceed 10 mph at the application site.
- **DO NOT** apply during temperature inversions.

### **Ground Applications**

- For all applications, applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.
- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.

### **Boom-less Ground Applications**

- Applicators are required to use a medium or coarser droplet size (ASABE S572.1) for all applications.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- User must turn off outward pointing nozzles at row ends and when spraying outer row.
- **DO NOT** apply during temperature inversions.

## **SPRAY DRIFT ADVISORIES**

- THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
- BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

### **IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

### **Controlling Droplet Size - Ground Boom**

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

### **Controlling Droplet Size – Aircraft**

- Adjust Nozzles - Follow nozzle manufacturer's recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

### **BOOM HEIGHT – Ground Boom**

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

### **RELEASE HEIGHT – Aircraft**

Higher release heights increase the potential for spray drift. When applying aerially to crops, **DO NOT** release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.

### **SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

### **TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

### **TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversion.

### **WIND**

Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.** Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

### **BOOM-LESS GROUND APPLICATIONS**

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

### **HANDHELD TECHNOLOGY APPLICATIONS**

Take precautions to minimize spray drift.

## **APPLICATION INFORMATION**

Use sufficient water volume to provide thorough and uniform coverage to obtain the most effective disease control.

### **Ground Application**

For optimum disease control, apply in sufficient water to ensure thorough coverage of foliage, bloom, and fruit.

### **Aerial Application**

For aerial application equipment, a minimum of **10** gallons of water per acre for tree crops and **2** gallons of water per acre for field and vegetable crops is required.

### **Boom-less Ground Application**

Air-assisted or air-blast sprayers move spray droplets into the crop canopy using a forced-air system. The fan must be set up to deliver only enough air volume to penetrate the canopy and provide good coverage. Adjust deflectors or other aiming devices to direct spray only to the target area. Equip sprayers with nozzles that provide accurate and uniform application.



## Greenhouse Application

LUNA PRIVILEGE can be used on cucurbits, fruiting vegetables, leafy greens (Subgroup 4-16A); leaf petiole vegetable (subgroup 22B), strawberries and other low-growing berries (except cranberries) as foliar, soil, or soil-less mix applied treatment only in greenhouses with good resistance management practices. See crop-specific use directions. Crop safety has not been confirmed on all cultivars. Plant compatibility testing is advised when first using under your greenhouse conditions. Some mild yellowing of leaf margins may be observed on cucurbits. LUNA PRIVILEGE is physically compatible with most commonly used fungicide, herbicide, insecticide, and foliar nutrient products. However, the compatibility of LUNA PRIVILEGE with all potential tank-mix partners has not been fully investigated. Prevent intense sunlight exposure for several hours after application. Solutions must be used within one day of mixing.

## In-furrow at-plant applications

Where permitted by crop specific use directions apply in-furrow during planting operations. Direct applications into the open furrow and cover with soil.

## Chemigation Application

Apply this product only through center pivot, motorized-lateral move, traveling gun, solid set or portable (wheel move, side roll, end tow, or hand move) and drip irrigation systems. **DO NOT** apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, must shut the system down and make necessary adjustments should the need arise.

LUNA PRIVILEGE has not been sufficiently tested when applied through irrigation systems to assure consistent product performance for all labeled uses. The following application techniques are provided for user reference but do not constitute a warranty of fitness for application through sprinkler or drip irrigation equipment.

**DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label prescribed safety devices for public water systems are in place. 'Public water system' means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ), back flow preventer, or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an alternative to the RPZ, the water from the public water system must be discharged into a reservoir tank prior to pesticide introduction.

There must be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. Pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The systems must contain functional interlocking controls, to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

**DO NOT** apply when wind speed favors drift. Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur. Apply pesticide continuously for the duration of the

water application. For mixing instructions, please refer to directions in the “Spray mixing and compatibility” section.

This product can be used through two basic types of irrigation systems as outlined in **Sections A and B** below. The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. Determine which type of irrigation system is in place, then refer to the appropriate directions provided below for each type. See crops section on the label for required treatment rates and additional use information.

#### **A. Center Pivot, Motorized-Lateral Move and Traveling Gun Irrigation Equipment**

For injections of pesticides, these continuously moving systems must use a positive displacement injection pump of either diaphragm or piston type and be constructed of materials that are compatible with pesticides. They must also be capable of being fitted with a system interlock and capable of injection at pressures approximately 2-3 times those encountered within the irrigation water line. Venturi applicator units cannot be used on these systems. Thoroughly mix required amount of this product for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run. Mixture in the chemical supply tank must be continuously agitated during the injection run. Shut off injection equipment after one revolution or run, but continue to operate irrigation system until this product has been cleared from the last sprinkler head.

#### **B. Solid-Set, Portable (Wheel Move, Side Roll, End Tow, or Hand Move) and Drip Irrigation Equipment**

With stationary systems, an effectively designed in-line Venturi applicator unit is preferred to support even and quick distribution. However, a positive-displacement pump can also be used. For solid set systems, determine acreage covered by sprinkler. Fill the tank of injection equipment with water and adjust flow to use contents over 30 to 45 minutes. Mix desired amount of this product for acreage to be covered with water so that the total mixture of this product plus water in the injection tank is equal to the quantity of water used during calibration. Provide chemical supply tank agitation sufficient for mixing until chemigation is completed. Operate entire system at normal pressures as advised by the manufacturer of injection equipment used, for amount of time established during calibration. This product can be injected during the irrigation cycle or as a separate application. For drip irrigation systems, introduce fungicide into irrigation solution for a period sufficient to distribute the product uniformly in the crop. Stop injection equipment with any system after treatment is completed and continue to operate irrigation system until this product has been cleared from the last sprinkler head or drip irrigation line.

### **SPRAY MIXING AND COMPATIBILITY**

Begin with clean spray equipment and add one-half of the required amount of water to the spray or mixing tank and start agitation. Add the required quantity of fungicide and the tank-mix partner if applicable to the water and complete filling with water to the required total volume. Follow the advice of your State Cooperative Extension Service for tank mixing with other products. In general, follow the order beginning first with water conditioners, water soluble packaging (wait for it to completely dissolve), wettable powders and water-dispersible granular products, liquid flowables and suspension concentrates, emulsifiable concentrates, and adjuvants last. Maintain agitation throughout spraying. **DO NOT** allow spray mixture to remain in the tank overnight, or for long periods during the day without agitation.

LUNA PRIVILEGE is physically compatible with most commonly used fungicide, herbicide, insecticide, and foliar nutrient products. However, the compatibility of LUNA PRIVILEGE with all potential tank-mix partners has not been fully investigated. If tank mixing with other pesticides is desirable, conduct a jar test with the volumes and rates typically used in agricultural application. Using a small container of water, add the proportionate amounts of the products: wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 15 minutes. Look for signs of separation, globules, sludge, flakes, or other precipitates. Physical compatibility is indicated if the combination remains mixed or can be remixed readily.

The crop safety of all potential tank-mixes with LUNA PRIVILEGE has not been tested on all crops listed on the label. Before applying any tank-mixture not specified on this label, safety to the target crop must be confirmed on a small portion of the crop listed on the label to be treated to ensure an adverse response will not occur.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

### **PRODUCT RESTRICTIONS**

**DO NOT** apply more than the maximum yearly rate for each specific crop from any combination of products containing Fluopyram.

### **ROTATIONAL CROP RESTRICTIONS**

The following crops can be replanted immediately following the last application of LUNA PRIVILEGE: Artichoke, (Globe); Brassica Head and Stem Vegetables (Group 5-16); Brassica Leafy Greens (Subgroup 4-16B) (except watercress); Bulb vegetables (Group 3-07); Carrot; Celtnce; Cereals (Groups 15 and 16) (except rice); Citrus (Group 10-10); Coffee; Corn; Cotton (subgroup 20C); Cucurbits (Group 9); Fennel; Florence (fresh leaves and stalks); Fruiting Vegetables (Group 8-10); Ginseng; Grapes and Small vines (except fuzzy kiwifruit) (Subgroup 13-07F); Herb (Subgroup 19A); Hops; Kohlrabi; Leafy greens (Subgroup 4-16A); Leaf petiole vegetables (Subgroup 22B); Legume Vegetables (Group 6-22) (except Dried Shelled Pulse Pea Legume Vegetables subgroup 6-22F); Mint (spearmint and peppermint, fresh and dried leaves); Papaya; Peanut; Pome fruit (Group 11-10); Potato and other root, tuberous and corm vegetables (except sugarbeet) (Subgroups 1B and 1C); Rapeseed (Subgroup 20A); Small Berries (caneberries and bushberries) (Subgroups 13-07A and 13-07B); Sorghum; Soybean; Spice (Group 26); Stone Fruits (Group 12-12); Strawberry and other low-growing berries, except cranberry (Subgroup 13-07G); Sugarbeet; Sunflower (Subgroup 20B); Tobacco; Tree Nuts (Group 14-12).

Alfalfa can be planted after 14 days and sugarcane can be planted after 14 days in Region 3 (entire state of Florida; Mobile and Baldwin counties in Alabama).

**DO NOT** rotate to crops other than those listed above.

## USE DIRECTIONS FOR SPECIFIC CROPS

ARTICHOKE, (GLOBE)		
Disease/Pest Suppression	Application Rate	Application Instructions
<p>Nematodes</p> <p>Powdery mildew (<i>Leveillula taurica</i>) (<i>Erysiphe cichoracearum</i>)</p>	<p>3.2 to 6.84 fl oz/acre</p> <p>(0.104 – 0.222 lb/acre fluopyram)</p>	<p>[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]</p> <p>[Drip applications are effective for suppression of this disease.] [Use the high rate when applying via drip equipment.]</p>
<p>Gray mold (<i>Botrytis cinerea</i>)</p>	<p>6.84 fl oz/acre</p> <p>(0.222 lb/acre fluopyram)</p>	<p>[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]</p>
<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]</li> <li>• Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]</li> <li>• Maximum number of applications per year: [4 (at 3.42 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]</li> <li>• [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]</li> <li>• Minimum retreatment interval: 7 days</li> <li>• <b>DO NOT</b> apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).</li> <li>• Apply using [ground], [aerial], [or] chemigation equipment.</li> <li>• Can be applied the day of harvest.</li> <li>• To limit the potential for development of disease resistance to this fungicide class, <b>DO NOT</b> make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.</li> <li>• For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.</li> </ul>		

**BRASSICA HEAD AND STEM VEGETABLES (Group 5-16)**

Broccoli; Brussels sprouts; Cabbage; Cabbage, Chinese, Napa; Cauliflower; Cultivars, varieties, and hybrids of these commodities.

<b>Disease/Pest Suppression</b>	<b>Application Rate</b>	<b>Application Instructions</b>
Powdery mildew ( <i>Erysiphe polygoni</i> ) ( <i>Erysiphe cruciferarum</i> )	3.2 to 6.84 fl oz/acre  (0.104 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 5- to 7-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.] [Drip applications are effective for suppression of this disease.] [Use the high rate when applying via drip equipment.]
Alternaria leaf spot ( <i>Alternaria</i> spp.)	4.0 to 6.84 fl oz/acre  (0.130 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 5- to 7-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]
Botrytis gray mold ( <i>Botrytis cinerea</i> )  Sclerotinia stem rot ( <i>Sclerotinia sclerotiorum</i> ) ( <i>Sclerotinia minor</i> )	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]
Alternaria leaf spot ( <i>Alternaria</i> spp.)  Botrytis gray mold ( <i>Botrytis cinerea</i> )  Nematodes  Powdery mildew ( <i>Erysiphe polygoni</i> ) ( <i>Erysiphe cruciferarum</i> )  Sclerotinia stem rot ( <i>Sclerotinia sclerotiorum</i> ) ( <i>Sclerotinia minor</i> )	3.6 to 6.84 fl oz/acre  (0.117 – 0.222 lb/acre fluopyram)	<b>Soil Application</b> [Use the higher rate if pest infestations are expected to be severe.] Apply specified dosage [in the following methods]: <ul style="list-style-type: none"> <li>• Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.</li> <li>• [Post-planting drench, or hill drench.]</li> <li>• [Transplant water drench with mechanical planting.]</li> <li>• [Transplant water drench with hand planting.]</li> </ul> Minimum 5-day interval between soil applications.

## BRASSICA HEAD AND STEM VEGETABLES (Group 5-16) - Continued

### Restrictions:

- Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]
- Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]
- Maximum number of applications per year: [4 (at 3.42 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]
- [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]
- Minimum retreatment interval: 5 days
- **DO NOT** apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using [ground], [aerial], [or] chemigation equipment.
- Can be applied the day of harvest.
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.

**BRASSICA LEAFY GREENS (Subgroup 4-16B), EXCEPT WATERCRESS**

Arugula; broccoli, Chinese; broccoli raab; cabbage, abyssinian; cabbage, Chinese, bok choy; cabbage, seakale; collards; cress, garden; cress, upland; hanover salad; kale; maca, leaves; mizuna; mustard greens; radish, leaves; rape greens; rocket, wild; shepherd's purse; turnip greens; cultivars, varieties, and hybrids of these commodities.

Disease Suppression	Application Rate	Application Instructions
Powdery mildew ( <i>Erysiphe polygoni</i> ) ( <i>Erysiphe cruciferarum</i> )	3.2 to 6.84 fl oz/acre  (0.104 - 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 5- to 7-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.] [Drip applications are effective for suppression of this disease.] [Use the high rate when applying via drip equipment.]
Alternaria leaf spot ( <i>Alternaria</i> spp.)	4.0 to 6.84 fl oz/acre  (0.130 - 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 5- to 7-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]
Botrytis gray mold ( <i>Botrytis cinerea</i> )  Sclerotinia stem rot ( <i>Sclerotinia sclerotiorum</i> ) ( <i>Sclerotinia minor</i> )	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]
Pest Suppression	Application Rate	Application Instructions
Nematodes	3.6 to 6.84 fl oz/acre  (0.117 - 0.222 lb/acre fluopyram)	[Use the higher rate if pest infestations are expected to be severe.] Apply specified dosage [in the following methods]: <ul style="list-style-type: none"> <li>• Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.</li> <li>• [Post-planting drench, or hill drench.]</li> <li>• [Transplant water drench with mechanical planting.]</li> <li>• [Transplant water drench with hand planting.]</li> </ul> Minimum 5-day interval between soil applications.

**Restrictions:**

- Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]
- Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]
- Maximum number of applications per year: [4 (at 3.42 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]
- [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]

<b>BRASSICA LEAFY GREENS (Subgroup 4-16B) - Continued</b>
<ul style="list-style-type: none"><li>• <b>DO NOT</b> apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).</li><li>• Apply using [ground], [aerial], [or] chemigation equipment.</li><li>• Can be applied the day of harvest.</li><li>• To limit the potential for development of disease resistance to this fungicide class, <b>DO NOT</b> make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.</li><li>• For soil application, to limit the potential for development of disease resistance to this chemical class, the first foliar fungicide application after LUNA PRIVILEGE should be a product from a different FRAC group.</li></ul>



## BULB VEGETABLES (group 3-07)

**Onion, bulb subgroup:** Bulb Daylily; Bulb Fritillaria; Bulb Garlic; Bulb Great-Headed Garlic; Bulb Serpent Garlic; Bulb Lily; Bulb Onion; Bulb Chinese Onion; Pearl Onion; Bulb Potato Onion; Bulb Shallot; cultivars, varieties, and/or hybrids of these.

**Onion, green subgroup:** Fresh Leaves Chive; Fresh Leaves Chinese Chive; Elegans Hosta; Leaves Fritillaria; Kurrat; Lady's Leek; Leek; Wild Leek; Beltsville Bunching Onion; Fresh Onion; Green Onion; Macrostem Onion; Tree Onion Tops; Welsh Onion Tops; Fresh Leaves Shallot; cultivars, varieties, and/or hybrids of these.

Disease/Pest Suppression	Application Rate	Application Instructions
Purple blotch ( <i>Alternaria porri</i> )	4.0 to 6.84 fl oz/acre  (0.130 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 5- to 7-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]
Botrytis leaf blight; neck rot ( <i>Botrytis squamosa</i> ) ( <i>Botrytis allii</i> ) ( <i>Botrytis cinerea</i> )	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 5- to 7-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]
White rot ( <i>Sclerotium cepivorum</i> )	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 5- to 7-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Botrytis leaf blight, neck rot <sup>[1]</sup> ( <i>Botrytis squamosa</i> ) ( <i>Botrytis allii</i> ) ( <i>Botrytis cinerea</i> )  Nematodes <sup>[1]</sup>  Purple blotch <sup>[1]</sup> ( <i>Alternaria porri</i> )  Pink root <sup>[1]</sup> ( <i>Phoma terrestris</i> )  White rot <sup>[1]</sup> ( <i>Sclerotium cepivorum</i> )	3.0 to 6.84 fl oz/acre  (0.098 – 0.222 lb/acre fluopyram)	<b>Soil Application</b> Apply specified dosage using any of the following methods: <ul style="list-style-type: none"> <li>• [Pre-plant banded or broadcast spray directed to the soil and incorporated into the planting bed during planting.]</li> <li>• [In-furrow drench.]</li> <li>• [In-furrow spray during planting directed on or below seed.]</li> <li>• [Chemigation into the root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.]</li> <li>• [Chemigation using overhead irrigation equipment.]</li> <li>• [Post-plant drench or hill drench.]</li> </ul> Minimum 14-day interval between applications.

## ***BULB VEGETABLES (group 3-07) - Continued***

### **Restrictions:**

- Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]
- Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]
- Maximum number of applications per year: [4 (at 3.42 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]
- [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]
- Minimum foliar retreatment interval: 5 days.
- Minimum soil retreatment interval: 14 days.
- **DO NOT** apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using [ground,] [aerial] [or] chemigation equipment.
- For foliar applications, LUNA PRIVILEGE can be applied the day of harvest.
- For soil applications, **DO NOT** apply LUNA PRIVILEGE within 30 days of harvest.
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.

[<sup>1</sup>Not for use in CA [without a supplemental label.]]

CARROT		
Disease/Pest Suppression	Application Rate	Application Instructions
Powdery mildew ( <i>Erysiphe</i> spp.) ( <i>Leveillula taurica</i> )	4.8 to 6.84 fl oz/acre  (0.156 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 5- to 7-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]
Cottony rot ( <i>Sclerotinia sclerotiorum</i> )  Leaf blight ( <i>Alternaria dauci</i> )  Leaf spot / Early blight ( <i>Cercospora carotae</i> )  Nematodes	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 5- to 7-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]</li> <li>• Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]</li> <li>• Maximum number of applications per year: [2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]</li> <li>• [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]</li> <li>• Minimum retreatment interval: 5 days</li> <li>• <b>DO NOT</b> apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (foliar only).</li> <li>• Apply using [ground,] [aerial,] [or] overhead chemigation equipment.</li> <li>• Can be applied the day of harvest.</li> <li>• To limit the potential for development of disease resistance to this fungicide class, <b>DO NOT</b> make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.</li> <li>• Tops or greens from this crop can be utilized for food or feed.</li> </ul>		

CELTUCE AND FENNEL, FLORENCE (FRESH LEAVES AND STALK) <sup>[1]</sup>		
Disease Suppression	Application Rate	Application Instructions
Powdery mildew ( <i>Erysiphe cichoracearum</i> )	3.2 to 6.84 fl oz/acre  (0.104 - 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.] [Drip applications are effective for suppression of this disease.] [Use the high rate when applying via drip equipment. ]
Lettuce drop ( <i>Sclerotinia minor</i> ) ( <i>Sclerotinia sclerotiorum</i> )	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7 to 10-day interval. [Can be applied as a banded spray treatment for celtuce (limited by BBCH15 stage, 5 leaf stage).]
Botrytis rot ( <i>Botrytis cinerea</i> )  Late blight ( <i>Septoria apiicola</i> )	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval.
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]</li> <li>Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]</li> <li>Maximum number of applications per year: [4 (at 3.42 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]</li> <li>[Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]</li> <li><b>DO NOT</b> apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).</li> <li>Apply using [ground,] [aerial,] [or] chemigation equipment.</li> <li>Can be applied broadcast the day of harvest (field use).</li> <li>For greenhouse uses, the PHI is 3 days.</li> <li>For banded application, PHI is 20 days.</li> <li>To limit the potential for development of disease resistance to this fungicide class, <b>DO NOT</b> make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.</li> </ul>		
[ <sup>1</sup> Not for use in CA [without a supplemental label.]]		

**CEREALS (Groups 15 and 16), EXCEPT RICE**

See separate instructions for: Corn and Sorghum.

Barley; Buckwheat; Pearl Millet; Proso Millet; Oats; Rye; Teosinte; Triticale; Wheat.

Disease/Pest Suppression	Application Rate	Application Instructions
Powdery mildew ( <i>Erysiphe graminis</i> )  Septoria leaf and glume blotch ( <i>Septoria</i> spp.)	3.42 fl oz/acre  (0.111 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval. [Can be applied in furrow.]
Nematodes	2.0 to 3.42 fl oz/acre  (0.065 – 0.111 lb/acre fluopyram)	<b>Soil Application</b> Apply specified dosage using any of the following methods: <ul style="list-style-type: none"> <li>• In-furrow spray during planting directed on or below seed.</li> <li>• Chemigation into root-zone through low-pressure drip or trickle irrigation.</li> <li>• Overhead chemigation in sufficient water volume to move LUNA PRIVILEGE into the root zone</li> </ul> Minimum 14-day interval between soil applications.

**Restrictions:**

- Maximum single application rate: [3.42 fl oz/acre of LUNA PRIVILEGE (0.111 lb/acre fluopyram)]
- Maximum annual application rate: [6.84 fl oz of LUNA PRIVILEGE per acre (0.222 lb/acre fluopyram) per year.]
- Maximum number of applications per year: [3 (at 2.28 fl oz/acre of LUNA PRIVILEGE) or 2 (at 3.42 fl oz/acre of LUNA PRIVILEGE)]
- [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 3.42 fl oz/acre and the maximum annual rate must not exceed 6.84 fl oz/acre of LUNA PRIVILEGE.]
- Minimum retreatment interval: 14 days
- **DO NOT** apply more than 0.222 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using [ground,] [aerial,] [or] chemigation equipment.
- **DO NOT** apply LUNA PRIVILEGE within 30 days of grain or straw harvest.
- **DO NOT** harvest hay or allow livestock to graze treated area for 14 days after application.
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.

## CITRUS FRUIT (Group 10-10)

Australian Desert Lime; Australian Finger Lime; Australian Round Lime; Brown River Finger Lime; Calamondin; Citron; Citrus Hybrids; Grapefruit; Japanese Summer Grapefruit; Kumquat; Lemon; Lime; Mediterranean Mandarin; Mount White Lime; New Guinea Wild Lime; Sour Orange; Sweet Orange; Pummelo; Russell River Lime; Satsuma Mandarin; Sweet Lime; Tachibana Orange; Tahiti Lime; Tangelo; Tangerine (Mandarin); Tangor; Trifoliate Orange; Uniq Fruit; cultivars, varieties and/or hybrids of these.

Disease/Pest Suppression	Application Rate	Application Instructions
Alternaria brown spot ( <i>Alternaria alternata</i> )	4.8 to 6.84 fl oz/acre	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 21-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]
Scab ( <i>Elsinoe fawcettii</i> )	(0.156 – 0.222 lb/acre fluopyram)	
Alternaria ( <i>Alternaria alternata</i> )	6.84 fl oz/acre	<b>Soil Applications</b> - Apply specified dosage by chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment. For optimum results, apply to newly planted trees or those previously trained to drip, trickle or micro-sprinkler irrigation. Soil must be lightly pre-wetted to break soil surface tension prior to applications.  Minimum 30-day interval between soil applications.
Greasy Spot ( <i>Mycosphaerella citri</i> )	(0.222 lb/acre fluopyram)	
Nematodes		
Melanose ( <i>Diaporthe citri</i> )		
Scab ( <i>Elsinoe fawcettii</i> )		

### Restrictions:

- Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]
- Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]
- Maximum number of applications per year: [2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]
- [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]
- Minimum foliar retreatment interval: 7 days.
- Minimum soil retreatment interval: 30 days.
- **DO NOT** apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using [ground,] [aerial,] [or] chemigation equipment.
- **DO NOT** apply LUNA PRIVILEGE within 7 days of harvest.
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.

COFFEE <sup>[1]</sup>		
Pest / Disease Suppression	Application Rate	Application Instructions
Nematodes	13.7 fl oz/acre  (0.445 lb/acre fluopyram)	Apply at the critical timings for pest/disease suppression as a drench application. Refer to University and/or extension guidelines for best application timings.
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• Maximum single application rate: 13.7 fl oz/acre of LUNA PRIVILEGE (0.445 lb/acre fluopyram)</li> <li>• Maximum annual application rate: 13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.</li> <li>• Maximum number of applications per year: 1</li> <li>• <b>DO NOT</b> apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).</li> <li>• Apply using ground or chemigation equipment.</li> <li>• <b>DO NOT</b> apply LUNA PRIVILEGE within 90 days of harvest.</li> <li>• To limit the potential for development of disease resistance to this fungicide class, <b>DO NOT</b> make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.</li> <li>• For soil application, to limit the potential for development of disease resistance to this chemical class, the first foliar fungicide application after LUNA PRIVILEGE should be a product from a different FRAC group.</li> </ul>		
[ <sup>1</sup> Not for use in CA [without a supplemental label.]]		

## CORN

Field Corn; Field Corn Grown For Seed; Popcorn; Sweet Corn.

Disease/Pest Suppression	Application Rate	Application Instructions
Powdery mildew ( <i>Erysiphe graminis</i> )  Septoria leaf and glume blotch ( <i>Septoria</i> spp.)	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval. [Can be applied in furrow.]
Nematodes	2.0 to 6.84 fl oz/acre  (0.065 – 0.222 lb/acre fluopyram)	<b>Soil Applications</b> Apply specified dosage using any of the following methods: <ul style="list-style-type: none"><li>• In-furrow spray during planting directed on or below seed.</li><li>• Chemigation into root-zone through low-pressure drip or trickle irrigation.</li><li>• Overhead chemigation in sufficient water volume to move LUNA PRIVILEGE into the root zone.</li></ul> Minimum 14-day interval between soil applications.

### Restrictions:

- Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]
- Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]
- Maximum number of applications per year: [6 (at 2.28 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]
- [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]
- Minimum retreatment interval: 14 days
- **DO NOT** apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using ground, aerial, or chemigation equipment.
- **DO NOT** apply LUNA PRIVILEGE within 14 days of harvest.
- **DO NOT** allow livestock to graze treated area for 14 days after application.
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.

## COTTON (Subgroup 20C)



Cottonseed; cultivars, varieties and other hybrids of these.		
Disease/Pest Suppression	Application Rate	Application Instructions
Hardlock ( <i>Fusarium</i> spp.)	4.8 to 6.84 fl oz/acre  (0.156 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings.
Target spot ( <i>Corenespora cassiicola</i> )	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. [Can be applied in furrow.]
<i>Fusarium</i> spp.  Target spot ( <i>Corenespora cassiicola</i> )  Nematodes  Fusarium wilt ( <i>Fusarium</i> spp.)	2.0 to 6.84 fl oz/acre  (0.065 – 0.222 lb/acre fluopyram)	<b>Soil Application</b> Apply specified dosage using any of the following methods: <ul style="list-style-type: none"> <li>• In-furrow spray during planting directed on or below seed.</li> <li>• Chemigation into root-zone through low-pressure drip or trickle irrigation.</li> <li>• Overhead chemigation in sufficient water volume to move LUNA PRIVILEGE into the root zone.</li> </ul>
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]</li> <li>• Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]</li> <li>• Maximum number of applications per year: [6 (at 2.28 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]</li> <li>• [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]</li> <li>• <b>DO NOT</b> apply more than one application of 6.84 fl oz per acre as a foliar treatment.</li> <li>• <b>DO NOT</b> apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (seed treatment, soil or foliar).</li> <li>• Apply using [ground,] [aerial,] [or] chemigation equipment.</li> <li>• <b>DO NOT</b> apply LUNA PRIVILEGE within 30 days of harvest.</li> <li>• To limit the potential for development of disease resistance to this fungicide class, <b>DO NOT</b> make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.</li> <li>• For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.</li> </ul>		

## CUCURBITS (Group 9)

**Melon subgroup:** Citron Melon; Muskmelon (hybrids and/or cultivars of *Cucumis Melo* including True Cantaloupe, Cantaloupe, Casaba, Crenshaw Melon, Golden Pershaw Melon, Honeydew Melon, Honey Balls, Mango Melon, Persian Melon, Pineapple Melon, Santa Claus Melon, Snake Melon); Watermelon.

**Squash/Cucumber subgroup:** Chayote (Fruit); Chinese Waxgourd; Cucumber; Gherkin; Gourd, Edible; *Momordica* spp.; Pumpkin; Squash, Summer; Squash, Winter.

Disease/Pest Suppression	Application Rate	Application Instructions
Powdery mildew ( <i>Sphaerotheca fuliginea</i> / <i>Podosphaera xanthii</i> ) ( <i>Erysiphe cichoracearum</i> )	3.2 to 6.84 fl oz/acre  (0.104 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 5- to 14-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals. ] [Drip applications are effective for suppression of this disease.] [Use the high rate when applying via drip equipment.] [Can be applied in furrow.]
Alternaria leaf spot ( <i>Alternaria</i> spp.)  Gray mold ( <i>Botrytis cinerea</i> )  Gummy stem blight ( <i>Didymella bryoniae</i> )	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval. [When disease pressure is severe, use the shorter intervals.]
Alternaria leaf spot ( <i>Alternaria</i> spp.)  Fusarium wilt ( <i>Fusarium</i> spp.)  Nematodes  Powdery mildew ( <i>Sphaerotheca fuliginea</i> / <i>Podosphaera xanthii</i> ) ( <i>Erysiphe cichoracearum</i> )	3.0 to 6.84 fl oz/acre  (0.098 – 0.222 lb/acre fluopyram)	<b>Soil Application</b> [Use the higher rate if pest infestations are expected to be severe.] Apply specified dosage [in the following methods]: <ul style="list-style-type: none"> <li>• Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.</li> <li>• [Post-planting drench, or hill drench.]</li> <li>• [Transplant water drench with mechanical planting.]</li> <li>• [Transplant water drench with hand planting.]</li> </ul> [Minimum 5-day interval between soil applications.]

### ***CUCURBITS (Group 9) - Continued***

**Note:** A mild yellowing on leaf margins is sometimes seen following application of LUNA PRIVILEGE in cucurbits.

**Restrictions:**

- Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]
- Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]
- Maximum number of applications per year: [4 (at 3.42 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]
- [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]
- Minimum retreatment interval: 5 days
- **DO NOT** apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using [ground,] [aerial,] [or] chemigation equipment.
- Can be applied broadcast the day of harvest (field use).
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.

## FRUITING VEGETABLES (Group 8-10)

**Tomato subgroup:** Bush Tomato; Cocona; Currant Tomato; Garden Huckleberry; Goji Berry; Groundcherry; Naranjilla; Sunberry; Tomatillo; Tomato; Tree Tomato; cultivars, varieties, and/or hybrids of these.

**Pepper/Eggplant subgroup:** African Eggplant; Bell Pepper; Eggplant; Martynia; Nonbell Pepper; Okra; Pea Eggplant; Pepino; Roselle; Scarlet Eggplant; cultivars, varieties, and/or hybrids of these.

Disease/Pest Suppression	Application Rate	Application Instructions
Powdery mildew ( <i>Oidiopsis taurica</i> / <i>Leveillula taurica</i> ) ( <i>Sphaerotheca</i> spp.)	4.8 to 6.84 fl oz/acre  (0.156 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7-day interval. [Drip applications are effective for suppression of this disease.] [Use the high rate when applying via drip equipment. ] [Can be applied in furrow.]
Early blight ( <i>Alternaria solani</i> )  Septoria blight ( <i>Septoria lycopersici</i> )	4.0 to 6.84 fl oz/acre  (0.130 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7-day interval.
Gray mold ( <i>Botrytis cinerea</i> )	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]
Early blight ( <i>Alternaria solani</i> )  Fusarium wilt ( <i>Fusarium oxysporum</i> )  Fusarium stem rot and vine decline ( <i>Fusarium falciforme</i> )  Nematodes  Powdery mildew ( <i>Oidiopsis taurica</i> / <i>Leveillula taurica</i> ) ( <i>Sphaerotheca</i> spp.)  Septoria blight ( <i>Septoria lycopersici</i> )	3.0 to 6.84 fl oz/acre  (0.098 – 0.222 lb/acre fluopyram)	<b>Soil Applications:</b> Apply specified dosage [in the following methods]: <ul style="list-style-type: none"> <li>• Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.</li> <li>• [In-furrow spray directed on or below seed.]</li> <li>• [Post-planting drench, or hill drench.]</li> <li>• [Transplant water drench with mechanical planting.]</li> <li>• [Transplant water drench with hand planting.]</li> </ul> Minimum 7-day interval between soil applications.

## **FRUITING VEGETABLES (Group 8-10) - Continued**

### **Restrictions:**

- Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]
- Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]
- Maximum number of applications per year: [4 (at 3.42 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]
- [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]
- Minimum retreatment interval: 7 days
- **DO NOT** apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using [ground,] [aerial,] [or] chemigation equipment.
- Can be applied broadcast the day of harvest (field use).
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.

GINSENG		
Disease Suppression	Application Rate	Application Instructions
Alternaria blight <i>(Alternaria panax)</i>  Botrytis blight <i>(Botrytis cinerea)</i>  Powdery mildew <i>(Erysiphe spp.)</i>	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 5- to 7-day interval.
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• Maximum single application rate: 6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)</li> <li>• Maximum annual application rate: 13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.</li> <li>• Maximum number of applications per year: 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)</li> <li>• Minimum retreatment interval: 5 days</li> <li>• <b>DO NOT</b> apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).</li> <li>• Apply using [ground,] [aerial,] [or] chemigation equipment.</li> <li>• <b>DO NOT</b> apply LUNA PRIVILEGE within 7 days of harvest.</li> <li>• To limit the potential for development of disease resistance to this fungicide class, <b>DO NOT</b> make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.</li> <li>• For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.</li> </ul>		

**GRAPES AND SMALL VINES (except fuzzy kiwifruit) (Subgroup 13-07F)**

Amur River Grape; Gooseberry; Grape; Hardy Kiwifruit; Maypop; Schisandra Berry; cultivars varieties, and/or hybrids of these.

Disease Suppression	Application Rate	Application Instructions
Powdery mildew ( <i>Uncinula necator</i> )	3.2 to 6.84 fl oz/acre  (0.104 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 12- to 21-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.] [Drip applications are effective for suppression of this disease.] [Use the high rate when applying via drip equipment.]
Black rot ( <i>Guignardia bidwellii</i> )  Botrytis bunch rot / Gray mold ( <i>Botrytis cinerea</i> )	6.0 to 6.84 fl oz/acre  (0.195 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 12- to 21-day interval. [When disease pressure is severe, use the shorter intervals.]

**Restrictions:**

- Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]
- Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]
- Maximum number of applications per year: [4 (at 3.42 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]
- [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]
- Minimum retreatment interval: 12 days
- **DO NOT** apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application.
- Apply using [ground,] [aerial,] [or] chemigation equipment.
- **DO NOT** use LUNA PRIVILEGE after BBCH 73 stage (when bunches begin to hang) or within 45 days of harvest.
- For foliar application, to limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.

## HERB (Subgroup 19A)

Angelica; Balm; Basil; Borage; Burnet; Camomile; Catnip; Chervil (Dried); Chive; Chinese Chive; Chive; Clary; Coriander (Leaf); Costmary; Culantro (Leaf); Curry (Leaf); Dillweed; Horehound; Hyssop; Lavender; Lemongrass; Lovage (Leaf); Marigold; Marjoram (*Origanum* Spp.); Nasturtium; Parsley (Dried); Pennyroyal; Rosemary; Rue; Sage; Savory, Summer And Winter; Sweet Bay; Tansy; Tarragon; Thyme; Wintergreen; Woodruff; And Wormwood.

Disease Suppression	Application Rate	Application Instructions
Powdery mildew ( <i>Erysiphe</i> spp.)	4.0 to 6.84 fl oz/acre  (0.130 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.] [Drip applications are effective for suppression of this disease.] [Use the high rate when applying via drip equipment.]
Gray mold ( <i>Botrytis cinerea</i> )  Leaf blight ( <i>Alternaria</i> spp.)	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval.

### Restrictions:

- Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]
- Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]
- Maximum number of applications per year: [3 (at 4.56 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]
- [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]
- Minimum retreatment interval: 7 days
- **DO NOT** apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using [ground,] [aerial,] [or] chemigation equipment.
- **DO NOT** apply LUNA PRIVILEGE within 14 days of harvest.
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.



<b>HOPS</b>		
<b>Disease/Pest Suppression</b>	<b>Application Rate</b>	<b>Application Instructions</b>
Powdery mildew ( <i>Sphaerotheca humuli</i> ) ( <i>Sphaerotheca macularis</i> )	3.2 to 6.84 fl oz/acre  (0.104 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval. [Make applications in sufficient water volume (just to the point of runoff). ][Thorough coverage is needed for disease suppression.] Apply increasing product rate as plant size increases.  3.2 fl oz with 15 - 30 gals./acre 4.8 fl oz with 31 - 60 gals./acre 6.4 fl oz with 61 - 90 gals./acre 6.84 fl oz with 91 - 200 gals./acre  [Drip applications are effective for suppression of this disease.] [Use the high rate when applying via drip equipment.]
Gray mold ( <i>Botrytis cinerea</i> )	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval.
Nematodes  Powdery mildew ( <i>Sphaerotheca humuli</i> ) ( <i>Sphaerotheca macularis</i> )	4.0 to 6.84 fl oz/acre  (0.130 – 0.222 lb/acre fluopyram)	<b>Soil Application</b> Apply specified dosage by soil drench at planting, or by chemigation into the root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment. For optimum results, apply to newly planted vines or those previously trained to drip, trickle or micro-sprinkler irrigation. Soil must be lightly pre-wetted to break surface tension prior to application.  Minimum 14-day interval between soil applications.

## **HOPS - Continued**

### **Restrictions:**

- Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]
- Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]
- Maximum number of applications per year: [4 (at 3.42 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]
- [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]
- Minimum retreatment interval: 14 days
- **DO NOT** apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using [ground,] [aerial,] [or] chemigation equipment.
- **DO NOT** apply LUNA PRIVILEGE within 7 days of harvest.
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.

KOHLRABI <sup>[1]</sup>		
Disease Suppression	Application Rate	Application Instructions
Powdery mildew ( <i>Erysiphe polygoni</i> ) ( <i>Erysiphe cruciferarum</i> )	3.2 to 6.84 fl oz/acre  (0.104 - 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 5- to 7-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.] [Drip applications are effective for suppression of this disease.] [Use the high rate when applying via drip equipment.]
Alternaria leaf spot ( <i>Alternaria</i> spp.)	4.0 to 6.84 fl oz/acre  (0.130 - 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 5- to 7-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]
Botrytis gray mold ( <i>Botrytis cinerea</i> )  Sclerotinia stem rot ( <i>Sclerotinia sclerotiorum</i> ) ( <i>Sclerotinia minor</i> )	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]
Pest Suppression	Application Rate	Application Instructions
Nematodes	3.6 to 6.84 fl oz/acre  (0.117 - 0.222 lb/acre fluopyram)	[Use the higher rate if pest infestations are expected to be severe.] Apply specified dosage [in the following methods]: <ul style="list-style-type: none"> <li>• Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.</li> <li>• [Post-planting drench, or hill drench.]</li> <li>• [Transplant water drench with mechanical planting.]</li> <li>• [Transplant water drench with hand planting.]</li> </ul> Minimum 5-day interval between soil applications.
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]</li> <li>• Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]</li> <li>• Maximum number of applications per year: [4 (at 3.42 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]</li> <li>• [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]</li> <li>• <b>DO NOT</b> apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).</li> <li>• Apply using [ground], [aerial], [or] chemigation equipment.</li> <li>• Can be applied the day of harvest.</li> </ul>		

<b>KOHLRABI - <i>Continued</i></b>
<ul style="list-style-type: none"><li>• To limit the potential for development of disease resistance to this fungicide class, DO NOT make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.</li><li>• For soil application, to limit the potential for development of disease resistance to this chemical class, the first foliar fungicide application after LUNA PRIVILEGE should be a product from a different FRAC group.</li></ul> <p>[<sup>1</sup>Not for use in CA [without a supplemental label.]]</p>

**LEAFY GREENS (Subgroup 4-16A)**

Amaranth, Chinese; amaranth, leafy; aster, Indian; blackjack; cat's whiskers; cham-chwi; cham-na-mul; chervil, fresh leaves; chipilin; chrysanthemum, garland; cilantro, fresh leaves; corn salad; cosmos; dandelion, leaves; danggwi, leaves; dillweed; dock; dol-nam-mul; ebolo; endive; escarole; fameflower; feather cockscomb; Good King Henry; huauzontle; jute, leaves; lettuce, bitter; lettuce, head; lettuce, leaf; orach; parsley, fresh leaves; plantain, buckhorn; primrose, English; purslane, garden; purslane, winter; radicchio; spinach; spinach, Malabar; spinach, New Zealand; spinach, tanier; Swiss chard; violet, Chinese, leaves; cultivars, varieties, and hybrids of these commodities.

Disease Suppression	Application Rate	Application Instructions
Fusarium wilt ( <i>Fusarium</i> spp.)	3.2 to 6.84 fl oz/acre	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.] [Drip applications are effective for suppression of this disease.] [Use the high rate when applying via drip equipment. ]
Powdery mildew ( <i>Erysiphe cichoracearum</i> )	(0.104 – 0.222 lb/acre fluopyram)	
Lettuce drop ( <i>Sclerotinia minor</i> ) ( <i>Sclerotinia sclerotiorum</i> )	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7 to 10-day interval. [Can be applied as a banded spray treatment for lettuce (limited by BBCH15 stage, 5 leaf stage).]
Botrytis rot ( <i>Botrytis cinerea</i> )	6.84 fl oz/acre	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval.
Late blight ( <i>Septoria apiicola</i> )	(0.222 lb/acre fluopyram)	

**Restrictions:**

- Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]
- Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]
- Maximum number of applications per year: [4 (at 3.42 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]
- [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]
- Minimum retreatment interval: 7 days
- **DO NOT** apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using [ground,] [aerial,] [or] chemigation equipment.
- Can be applied the day of harvest (field use).
- For greenhouse uses, the PHI is 3 days.
- [For banded application on lettuce, PHI is 20 days.]
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.

**LEAF PETIOLE VEGETABLES (Subgroup 22B)**

Cardoon; celery; Chinese; fuki; rhubarb; udo; zuiki; cultivars, varieties, and hybrids of these commodities.

Disease Suppression	Application Rate	Application Instructions
Powdery mildew ( <i>Erysiphe cichoracearum</i> )	3.2 to 6.84 fl oz/acre  (0.104 - 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.] [Drip applications are effective for suppression of this disease.] [Use the high rate when applying via drip equipment. ]
Sclerotinia root rot ( <i>Sclerotinia minor</i> ) ( <i>Sclerotinia sclerotiorum</i> )	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7 to 10-day interval. [Can be applied as a banded spray treatment for lettuce (limited by BBCH15 stage, 5 leaf stage).]
Botrytis rot ( <i>Botrytis cinerea</i> )  Late blight ( <i>Septoria apiicola</i> )	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval.

**Restrictions:**

- Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]
- Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]
- Maximum number of applications per year: [4 (at 3.42 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]
- [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]
- **DO NOT** apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using [ground,] [aerial,] [or] chemigation equipment.
- Can be applied broadcast the day of harvest (field use).
- For greenhouse uses, the PHI is 3 days.
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.

## LEGUME VEGETABLES

(Group 6-22) (except Dried Shelled Pulse Pea Legume Vegetable Subgroup 6-22F)

### SOYBEAN

**Edible podded bean legume vegetable (Subgroup 6-22A):** Asparagus bean, catjang bean, Chinese longbean, cowpea, French bean, garden bean, goa bean, green bean, guar bean, jackbean, kidney bean, lablab bean, moth bean, mung bean, navy bean, rice bean, scarlet runner bean, snap bean, sword bean, urd bean, vegetable soybean, velvet bean, wax bean, winged pea, yardlong bean

**Edible podded pea legume vegetable (Subgroup 6-22B):** Chickpea, dwarf pea, edible podded pea, grass-pea, green pea, lentil, pigeon pea, snap pea, snow pea, sugar snap pea

**Succulent shelled bean legume vegetable (Subgroup 6-22C):** Andean lupin, blackeyed pea, blue lupin, broad bean (fava bean), catjang bean, cowpea, crowder pea, goa bean, grain lupin, jackbean, lablab bean, lima bean, moth bean, scarlet runner bean, southern pea, sweet lupin, vegetable soybean (edamame), velvet bean, wax bean, white lupin, white sweet lupin, yellow lupin

**Succulent shelled pea legume vegetable (Subgroup 6-22D):** Chickpea, English pea, garden pea, green pea, lentil, pigeon pea

**Dried shelled pulse bean legume vegetable (except soybean) (Subgroup 6-22E):** Adzuki bean, African yam bean, American potato bean, Andean lupin, asparagus bean, black bean, blackeyed pea, blue lupin, broad bean, catjang bean, Chinese longbean, cowpea, cranberry bean, crowder pea, dry bean, field bean, French bean, garden bean, goa bean, grain lupin, great northern bean, green bean, guar bean, horse gram, jackbean, kidney bean, lablab bean, lima bean, morama bean, moth bean, mung bean, navy bean, pink bean, pinto bean, red bean, rice bean, scarlet runner bean, southern pea, sweet lupin, sword bean, tepary bean, urd bean, velvet bean, white lupin, white sweet lupin, winged pea, yardlong bean, yellow bean, yellow lupin

Disease/Pest Suppression	Application Rate	Application Instructions
Fusarium wilt ( <i>Fusarium</i> spp.)	4.8 to 6.84 fl oz/acre	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.] [Can be applied in furrow.]
Powdery mildew ( <i>Erysiphe pisi</i> )	(0.156 – 0.222 lb/acre fluopyram)	
Alternaria blight ( <i>Alternaria</i> spp.)	6.84 fl oz/acre	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]
White mold ( <i>Sclerotinia sclerotiorum</i> )	(0.222 lb/acre fluopyram)	
Gray mold ( <i>Botrytis cinerea</i> )		
Fusarium wilt ( <i>Fusarium</i> spp.)	2.0 to 6.84 fl oz/acre	<b>Soil Application</b> Apply in seed furrow and cover with soil, or chemigation into root-zone through low-pressure drip or trickle irrigation.  Minimum 7 day interval between soil applications.
Nematodes	(0.065 – 0.222 lb/acre fluopyram)	
Sudden Death Syndrome (SDS)		

( <i>Fusarium virguliforme</i> )		
<b>LEGUME VEGETABLES (Group 6-22) (except Dried Shelled Pulse Pea Legume Vegetable Subgroup 6-22F) - Continued</b>		
<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]</li> <li>• Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]</li> <li>• Maximum number of applications per year: [6 (at 2.28 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]</li> <li>• [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]</li> <li>• Minimum retreatment interval: 7 days</li> <li>• <b>DO NOT</b> apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).</li> <li>• Apply using [ground,] [aerial,] [or] chemigation equipment.</li> <li>• <b>DO NOT</b> apply LUNA PRIVILEGE within 14 days of harvest.</li> <li>• To limit the potential for development of disease resistance to this fungicide class, <b>DO NOT</b> make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.</li> <li>• For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.</li> <li>• <b>DO NOT</b> allow livestock to graze forage or hay and <b>DO NOT</b> harvest forage or hay for food or feed.</li> </ul>		



**MINT (Spearmint and Peppermint, fresh and dried leaves)<sup>[1]</sup>**

Pest / Disease Suppression	Application Rate	Application Instructions
Nematodes	6.0 to 6.84 fl oz/acre  (0.195 - 0.222 lb/acre fluopyram)	<p>Apply at the critical timings for pest/disease suppression. Refer to University and/or extension guidelines for best application timings. Continue as needed on a minimum 21-day interval.</p> <p>Apply specified dosage via overhead chemigation or as a foliar broadcast application in a minimum of 20 gallons per acre followed with ½ to ¾ inch of overhead irrigation within one hour of the application to move the product into the soil profile.</p>

**Restrictions:**

- Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]
- Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]
- Maximum number of applications per year: [2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]
- [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]
- **DO NOT** apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using ground or chemigation equipment.
- **DO NOT** apply LUNA PRIVILEGE within 28 days of harvest.
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, the first foliar fungicide application after LUNA PRIVILEGE should be a product from a different FRAC group.

[<sup>1</sup>Not for use in CA [without a supplemental label.]]

PAPAYA <sup>[1]</sup>		
Pest / Disease Suppression	Application Rate	Application Instructions
Black spot ( <i>Asperisporium caricae</i> )	3.8 fl oz/acre  (0.123 lb/acre fluopyram)	Apply at the critical timings for pest/disease suppression. Refer to University and/or extension guidelines for best application timings. Continue as needed on a minimum 7-day interval.
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• Maximum single application rate: 3.8 fl oz/acre of LUNA PRIVILEGE (0.123 lb/acre fluopyram)</li> <li>• Maximum annual application rate: 11.4 fl oz of LUNA PRIVILEGE per acre (0.369 lb/acre fluopyram) per year.</li> <li>• Maximum number of applications per year: 3</li> <li>• <b>DO NOT</b> apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).</li> <li>• Apply using ground, aerial, or chemigation equipment.</li> <li>• May be applied the day of harvest.</li> <li>• <b>DO NOT</b> use an organo-silicone surfactant.</li> <li>• To limit the potential for development of disease resistance to this fungicide class, <b>DO NOT</b> make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.</li> </ul>		
[ <sup>1</sup> Not for use in CA [without a supplemental label.]]		

PEANUT		
Disease/Pest Suppression	Application Rate	Application Instructions
Early leaf spot ( <i>Cercospora arachidicola</i> )  Late leaf spot ( <i>Cercosporidium personatum</i> )	5.6 to 6.84 fl oz/acre  (0.182 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval. [When disease pressure is severe, use the higher rates.] [Can be applied in furrow.]
Aspergillus crown rot ( <i>Aspergillus niger</i> )  Early leaf spot ( <i>Cercospora arachidicola</i> )  Late leaf spot ( <i>Cercosporidium personatum</i> )  Nematodes  White mold ( <i>Sclerotium rolfsii</i> )	4.0 to 6.84 fl oz/acre  (0.130 – 0.222 lb/acre fluopyram)	<b>Soil Application</b> Apply specified dosage using any of the following methods: <ul style="list-style-type: none"> <li>• [In-furrow spray during planting directed on or below seed.]</li> <li>• [Chemigation into root-zone through low-pressure drip or trickle irrigation.]</li> <li>• [Overhead chemigation in sufficient water volume to move LUNA PRIVILEGE into the root zone.]</li> </ul> Minimum 14-day interval between soil applications.
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]</li> <li>• Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]</li> <li>• Maximum number of applications per year: [3 (at 4.56 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]</li> <li>• [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]</li> <li>• Minimum retreatment interval: 14 days</li> <li>• <b>DO NOT</b> apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (seed treatment, soil or foliar).</li> <li>• Apply using [ground;] [aerial] [or] chemigation equipment.</li> <li>• <b>DO NOT</b> apply LUNA PRIVILEGE within 7 days of harvest (spray and soil applications).</li> <li>• To limit the potential for development of disease resistance to this fungicide class, <b>DO NOT</b> make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.</li> <li>• For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.</li> <li>• <b>DO NOT</b> allow livestock to graze peanut hay.</li> </ul>		

## POME FRUIT (Group 11-10)

Apple; Azarole; Crabapple; Loquat; Mayhaw; Medlar; Pear; Asian Pear; Quince; Chinese Quince; Japanese Quince; Tejocote; cultivars, varieties and/or hybrids of these.

Disease/Pest Suppression	Application Rate	Application Instructions
Powdery mildew ( <i>Podosphaera leucotricha</i> )	2.4 to 6.84 fl oz/acre  (0.078 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.] [Drip applications are effective for suppression of this disease.] [Use the high rate when applying via drip equipment.]
Scab, leaf ( <i>Venturia</i> spp.)	4.0 to 6.84 fl oz/acre  (0.130 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]
Scab, fruit ( <i>Venturia</i> spp.)	5.6 to 6.84 fl oz/acre  (0.182 – 0.222 lb/acre fluopyram)	[Make applications at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]
Sooty blotch ( <i>Gloeodes pomigena</i> )  Fly speck ( <i>Schizothyrium pomi</i> )	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]
Alternaria leaf spot ( <i>Alternaria</i> spp.)  Nematodes  Powdery mildew ( <i>Podosphaera leucotricha</i> )	4.0 to 6.84 fl oz/acre  (0.130 – 0.222 lb/acre fluopyram)	<b>Soil Applications:</b> Apply specified dosage by chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment. For optimum results, apply to newly planted trees or those previously trained to drip, trickle or micro-sprinkler irrigation. Soil must be lightly pre-wetted to break soil surface tension prior to applications.  Minimum 30-day interval between soil applications.

**POME FRUIT (Group 11-10) - Continued**

**Restrictions:**

- Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]
- Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]
- Maximum number of applications per year: [5 (at 2.74 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]
- [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]
- Minimum foliar retreatment interval: 7 days.
- Minimum soil retreatment interval: 30 days.
- **DO NOT** apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using [ground,] [aerial,] [or] chemigation equipment.
- **DO NOT** apply LUNA PRIVILEGE within 7 days of harvest.
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.

# **POTATO AND OTHER ROOT, TUBEROUS AND CORM VEGETABLES (Except Sugarbeet) (Subgroups 1B and 1C)**

(See separate use directions for: Artichoke, Globe; Carrot; Ginseng; and Sugarbeet)

**Root vegetables (except sugar beet) subgroup:** Beet, Garden; Burdock, Edible; Celeriac; Chervil, Turnip-Rooted; Chicory; Horseradish; Parsley, Turnip-Rooted; Parsnip; Radish; Radish, Oriental; Rutabaga; Salsify; Salsify, Black; Salsify, Spanish; Skirret; Turnip.

**Tuberous and corm vegetables subgroup:** Arracacha; Arrowroot; Chinese Artichoke; Artichoke, Jerusalem; Canna, Edible; Cassava, Bitter And Sweet; Chayote (Root); Chufa; Dasheen; Ginger; Leren; Potato; Sweet Potato; Tanier; Turmeric; Yam Bean; Yam, True.

Disease/Pest Suppression	Application Rate	Application Instructions
Brown spot <i>(Alternaria alternata)</i>  Early blight <i>(Alternaria solani)</i>  Black dot <i>(Colletotrichum coccodes)</i>  Botrytis leaf spot <i>(Botrytis cinerea)</i>  Powdery mildew <i>(Erysiphe polygoni)</i> <i>(Erysiphe betae)</i> <i>(Erysiphe spp.)</i>	4.0 to 6.84 fl oz/acre  (0.130 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 5- to 7-day interval. [When disease pressure is severe; use the higher rates and/or shorter intervals.] [Can be applied in furrow.]
White mold <i>(Sclerotinia sclerotiorum)</i>  Ascochyta leaf spot <i>(Ascochyta cynarae)</i>  Cercospora leaf spot <i>(Cercospora beticola)</i>	6.0 to 6.84 fl oz/acre  (0.195 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 5- to 7-day interval.

**POTATO AND OTHER ROOT, TUBEROUS AND CORM VEGETABLES (Except Sugarbeet) (Subgroups 1B and 1C) - Continued**

<b>Disease/Pest Suppression</b>	<b>Application Rate</b>	<b>Application Instructions</b>
Black dot ( <i>Colletotrichum coccodes</i> )  Early blight ( <i>Alternaria solani</i> )  Early dying ( <i>Verticillium</i> spp.)  Nematodes  White mold ( <i>Sclerotinia sclerotiorum</i> )	4.0 to 6.84 fl oz/acre  (0.130 – 0.222 lb/acre fluopyram)	<b>Soil Applications:</b> [Apply specified dosage as an in-furrow spray during planting directed on or below seed.] Apply specified dosage using overhead chemigation equipment. For transplanted crops including sweet potatoes: <ul style="list-style-type: none"> <li>• Post-planting drench, or hill drench.</li> <li>• Transplant water drench with mechanical planting.</li> <li>• Transplant water drench with hand planting.</li> </ul> [Despite suppression of root-knot nematode, tuber quality may not be adequately protected. ][If root-knot nematode is a severe economic pest, the use of other suppression measures should also be employed.]  Minimum 5-day interval between soil applications.

**Restrictions:**

- Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]
- Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]
- Maximum number of applications per year: [3 (at 4.56 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]
- [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]
- Minimum retreatment interval: 5 days
- **DO NOT** apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using [ground,] [aerial,] [or] chemigation equipment.
- **DO NOT** apply LUNA PRIVILEGE within 7 days of harvest.
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.
- Tops or greens can be utilized for food or feed.
- The grazing of livestock in treated areas within 7 days of application is prohibited.

## RAPESEED (Subgroup 20A)

Borage; Crambe; Cuphea; Echium; Flax Seed; Gold Of Pleasure; Hare's Ear Mustard; Lesquerella; Lunaria; Meadowfoam; Milkweed; Mustard Seed; Oil Radish; Poppy Seed; Rapeseed; Sesame; Sweet Rocket cultivars, varieties, and/or hybrids of these.

Disease Suppression	Application Rate	Application Instructions
Alternaria blackspot ( <i>Alternaria</i> spp.)  Sclerotinia stem rot ( <i>Sclerotinia sclerotiorum</i> )	4.1 fl oz/acre  (0.133 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on an 8- to 14-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.] [Good spray coverage of the plants is essential.]
Powdery mildew ( <i>Erysiphe cruciferarum</i> )	4.1 fl oz/acre  (0.133 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on an 8- to 14-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.] [Good spray coverage of the plants is essential.] [In furrow treatment is effective for suppression of this disease.]

### Restrictions:

- Maximum single application rate: 4.1 fl oz/acre of LUNA PRIVILEGE (0.133 lb/acre fluopyram)
- Maximum annual application rate: 8.22 fl oz of LUNA PRIVILEGE per acre (0.267 lb/acre fluopyram) per year.
- Maximum number of applications per year: 2 (at 4.1 fl oz/acre of LUNA PRIVILEGE)
- Minimum retreatment interval: 8 days
- **DO NOT** apply more than 0.267 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using [ground,] [aerial,] [or] chemigation equipment.
- **DO NOT** apply LUNA PRIVILEGE within 30 days of harvest.
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.



**SMALL BERRIES (Caneberries and Bushberries) (Subgroups 13-07A and 13-07B)**

**Caneberry subgroup:** Blackberry; Loganberry; Raspberry, Red And Black; Wild Raspberry; cultivars, varieties, and/or hybrids of these.

**Bushberry subgroup:** Aronia Berry; Blueberry, Highbush; Blueberry, Lowbush; Buffalo Currant; Chilean Guava; Currant, Black; Currant, Red; Elderberry; European, Barberry; Gooseberry; Cranberry, Highbush; Honeysuckle, Edible; Huckleberry; Jostaberry; Juneberry; Lingonberry; Native Currant; Salal; Sea Buckthorn; cultivars, varieties, and/or hybrids of these.

Disease/Pest Suppression	Application Rate	Application Instructions
Monilinia blight and mummy berry ( <i>Monilinia</i> spp.)	4.8 to 6.84 fl oz/acre  (0.156 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval.
Powdery mildew ( <i>Microsphaera</i> spp.)	4.8 to 6.84 fl oz/acre  (0.156 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval. [Drip applications are effective for suppression of this disease.][Use the high rate when applying via drip equipment.]
Alternaria leaf spot and fruit rot ( <i>Alternaria</i> spp.)  Gray mold ( <i>Botrytis cinerea</i> )  Leaf spot and blotch ( <i>Mycosphaerella</i> spp.) ( <i>Septoria</i> spp.)	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval.
Alternaria leaf spot ( <i>Alternaria</i> spp.)  Leaf spot and blotch ( <i>Mycosphaerella</i> spp.) ( <i>Septoria</i> spp.)  Nematodes  Powdery mildew ( <i>Microsphaera</i> spp.)	4.0 to 6.84 fl oz/acre  (0.130 – 0.222 lb/acre fluopyram)	<b>Soil Application</b> Apply specified dosage by soil drench at planting, or by chemigation into the root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment. For optimum results, apply to newly planted trees or those previously trained to drip, trickle or micro-sprinkler irrigation. Soil must be lightly pre-wetted to break surface tension prior to application.  Minimum 7-day interval between soil applications.

***SMALL BERRIES (Caneberries and Bushberries) (Subgroups 13-07A and 13-07B) - Continued***

**Restrictions:**

- Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]
- Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]
- Maximum number of applications per year: [3 (at 4.56 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]
- [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]
- Minimum retreatment interval: 7 days
- **DO NOT** apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using [ground], [aerial], [or] chemigation equipment.
- Can be applied the day of harvest.
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.

## SORGHUM

Disease/Pest Suppression	Application Rate	Application Instructions
Alternaria head blight ( <i>Alternaria</i> spp.)	5.47 fl oz/acre  (0.178 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. [Can be applied in furrow.]
Nematodes	2.0 to 5.47 fl oz/acre  (0.065 – 0.178 lb/acre fluopyram)	<b>Soil Application</b> Apply specified dosage using any of the following methods: <ul style="list-style-type: none"> <li>• In-furrow spray during planting directed on or below seed.</li> <li>• Chemigation into root-zone through low-pressure drip or trickle irrigation.</li> <li>• Overhead chemigation in sufficient water volume to move LUNA PRIVILEGE into the root zone</li> </ul>

### Restrictions:

- Maximum single application rate: [5.47 fl oz/acre of LUNA PRIVILEGE (0.178 lb/acre fluopyram)]
- Maximum annual application rate: [5.47 fl oz of LUNA PRIVILEGE per acre (0.178 lb/acre fluopyram) per year.]
- Maximum number of applications per year: [2 (at 2.73 fl oz/acre of LUNA PRIVILEGE) or 1 (at 5.47 fl oz/acre of LUNA PRIVILEGE)]
- [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 5.47 fl oz/acre and the maximum annual rate must not exceed 5.47 fl oz/acre of LUNA PRIVILEGE.]
- **DO NOT** apply more than one application of 5.47 fl oz per acre as a foliar treatment.
- **DO NOT** apply more than 0.178 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using [ground,] [aerial,] [or] chemigation equipment.
- **DO NOT** apply LUNA PRIVILEGE within 30 days of harvest of grain or stover.
- **DO NOT** allow livestock to graze treated area for 14 days after application.
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.

SOYBEAN		
Disease/Pest Suppression	Application Rate	Application Instructions
Fusarium wilt ( <i>Fusarium</i> spp.)  Powdery mildew ( <i>Erysiphe pisi</i> )	4.8 to 6.84 fl oz/acre  (0.156 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.] [Can be applied in furrow.]
Alternaria blight ( <i>Alternaria</i> spp.)  White mold ( <i>Sclerotinia sclerotiorum</i> )  Gray mold ( <i>Botrytis cinerea</i> )	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]
Fusarium wilt ( <i>Fusarium</i> spp.)  Nematodes  Sudden Death Syndrome (SDS) ( <i>Fusarium virguliforme</i> )	2.0 to 6.84 fl oz/acre  (0.065 – 0.222 lb/acre fluopyram)	<b>Soil Application</b> Apply in seed furrow and cover with soil, or chemigation into root-zone through low-pressure drip or trickle irrigation.  Minimum 7 day interval between soil applications.
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]</li> <li>Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]</li> <li>Maximum number of applications per year: [6 (at 2.28 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]</li> <li>[Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]</li> <li>Minimum retreatment interval: 7 days</li> <li><b>DO NOT</b> apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).</li> <li>Apply using [ground,] [aerial,] [or] chemigation equipment.</li> <li><b>DO NOT</b> apply LUNA PRIVILEGE within 14 days of harvest.</li> <li>To limit the potential for development of disease resistance to this fungicide class, <b>DO NOT</b> make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.</li> <li>For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.</li> <li><b>DO NOT</b> allow livestock to graze forage or hay and <b>DO NOT</b> harvest forage or hay for food or feed.</li> </ul>		

**SPICE (Group 26)**

Ajowan, seed; Alder buckhorn; Allspice; Ambrette, seed; Amla, seed; Angelica, dahurian, seed; Angelica, seed; Angostura, bark; Anise pepper; Anise, seed; Anise, star; Annatto, seed; Asafoetida; Ashwagandha, fruit; Autumn crocus; Balsam, Peruvian; Barberry, bark; Batavia-cassia, bark; Batavia-cassia, fruit; Belleric myrobalan; Betel vine; Birch, bark; Bisnaga, seed; Bitterwood; Black bread weed; Bloodroot; Blue mallee; Blushwood, seed; Boldo, leaf; Buchu; Calamus root; Candlebush; Canella, bark; Caper buds; Caper spurge, seed; Caraway, black; Caraway, fruit; Cardamom, black; Cardamom, Ethiopian; Cardamom, green; Cardamom, Nepal; Cardamom-amomum; Cascara sagrada; Cassia, bark; Cassia, Chinese, bark; Cassia, Chinese, fruit; Cassia, fruit; Cat's claw, bark; Catechu, bark; Celery, seed; Chaste tree, berry; Chaste tree, Chinese, roots; Chervil, seed; Chinese hawthorn; Chinese nutmeg tree; Chinese wineberry, fruit; Chinese-pepper; Cinnamon, bark; Cinnamon, fruit; Cinnamon, Saigon, bark; Cinnamon, Saigon, fruit; Clove buds; Clusterleaf; Comfrey; Copaiba; Coptis; Coriander, fruit; Coriander, seed; Cotton, bark; Crampbark; Cubeb, seed; Culantro, seed; Culvers root; Cumin; Cumin, black; Dill, seed; Dorrigio pepper, berry; Dorrigio pepper, leaf; Dragon blood; Echinacea, seed; Epimedium; Eucalyptus; Eucommia, bark; European beech; Felty germander; Fennel flower, seed; Fennel, common, fruit; Fennel, common, seed; Fennel, Florence, fruit; Fennel, Florence, seed; Fenugreek, seed; Fingerroot; Flame lily, seed; Frankincense; Frankincense, Indian; Fringetree, bark; Galbanum, resin; Gambooge; Grains of paradise; Grains of Selim; Guaiac; Guarana; Guggul; Gum Arabic; Gum ghatti; Gum karaya; Gum tragacanth; Haw, black; Honewort, seed; Imperatoria; Indian tobacco, seed; Iva; Jalap; Jamaica dogwood, bark; Juniper berry; Kaffir lime, leaf; Kewra; Kokam; Linden, leaf; Lovage, seed; Mace; Magnolia, bark; Mahaleb; Malabar cardamom; Malabar-tamarind; Malabathrum; Mastic; Micromeria, white; Milk thistle; Mioga; Miracle fruit; Mistletoe; Mojave yucca; Muira puama; Mustard, black; Mustard, brown; Mustard, seed; Mustard, white; Myrrh; Myrrh, bisabol; Myrtle, anise; Myrtle, leaf; Myrtle, lemon; Nasturtium, bush, pods; Nasturtium, garden, pods; Nasturtium, pods; Nettle, stinging, seed; Nutmeg; Osha; Pepper, black; Pepper, Indian long; Pepper, Javanese long; Pepper, leaf; Pepper, pink; Pepper, Sichuan; Pepper, white; Pepperbush, berry; Pepperbush, leaf; Peppercorn, green; Peppertree; Peppertree, Peruvian; Perilla, seed; Phellodendron; Pine, maritime; Poppy, seed; Prickly ash, Chinese; Prickly ash, Southern, bark; Pygeum; Qing hua jiao; Quassia, bark; Quebracho, bark; Quillaja; Quinine; Rauwolfia, bark; Resin spurge; Rue; Saffron crocus; Sandalwood, seed; Sassafras, bark; Sassafras, leaf; Saunders, red; Saw palmetto; Sesame, seed; Silktree, bark; Simaruba, bark; Skunk cabbage, root; Slippery elm; Stemona, root; Suma; Sumac, fragrant; Sumac, smooth, leaf; Taheebo, bark; Tamarind, seed; Tasmanian pepper, berry; Tasmanian pepper, leaf; Threelaf caper; Tsaoko; Vanilla; Wattleseed; White willow; Willow; Witch hazel; Yaw root; Yellow gentian, roots; Yohimbe; Cultivars, varieties, and hybrids of these commodities.

Disease Suppression	Application Rate	Application Instructions
Powdery mildew ( <i>Erysiphe</i> spp.)	4.0 to 6.84 fl oz/acre  (0.130 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.] [Drip applications are effective for suppression of this disease.] [Use the high rate when applying via drip equipment.]
Gray mold ( <i>Botrytis cinerea</i> )  Leaf blight ( <i>Alternaria</i> spp.)	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval.

**Restrictions:**

- Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]
- Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]

### **SPICE (Group 26) – Continued**

- Maximum number of applications per year: [3 (at 4.56 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]
- [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]
- Minimum retreatment interval: 7 days
- **DO NOT** apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using [ground,] [aerial,] [or] chemigation equipment.
- **DO NOT** apply LUNA PRIVILEGE within 14 days of harvest.
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.

For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.

## STONE FRUITS (Group 12-12)

**Cherry subgroup:** Capulin; Black Cherry; Nanking Cherry; Sweet Cherry; Tart Cherry; cultivars, varieties, and/or hybrids of these.

**Peach subgroup:** Peach; Nectarine; cultivars, varieties, and/or hybrids of these.

**Plum subgroup:** Apricot; Japanese Apricot; Chinese Jujube; Plum; American Plum; Beach Plum; Canada Plum; Cherry Plum; Chickasaw Plum; Damson Plum; Japanese Plum; Klamath Plum; Plumcot; Prune Plum; Sloe; cultivars, varieties, and/or hybrids of these.

Disease/Pest Suppression	Application Rate	Application Instructions
Powdery mildew Rusty spot ( <i>Podosphaera</i> spp.) ( <i>Sphaerotheca pannosa</i> )	4.0 to 6.84 fl oz/acre  (0.130 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 5- to 7-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.] [Drip applications are effective for suppression of this disease.] [Use the high rate when applying via drip equipment.]
Brown rot blossom blight Fruit rot ( <i>Monilinia</i> spp.)  Shot hole ( <i>Wilsonomyces carpophilus</i> )	4.0 to 6.84 fl oz/acre  (0.130 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 5- to 7-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]
Scab ( <i>Cladosporium carpophilum</i> )	4.8 to 6.84 fl oz/acre  (0.156 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 5- to 7-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]
Cherry leaf spot ( <i>Blumeriella jaapii</i> )  Jacket rot Green fruit rot ( <i>Botrytis cinerea</i> )	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 5- to 7-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]
Nematodes  Powdery mildew  Rusty spot ( <i>Podosphaera</i> spp.) ( <i>Sphaerotheca pannosa</i> )	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	<b>Soil Applications</b> - Apply specified dosage by chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment. For optimum results, apply to newly planted trees or those previously trained to drip, trickle or micro-sprinkler irrigation. Soil must be lightly pre-wetted to break soil surface tension prior to applications.  Minimum 30-day interval between soil applications.

## **STONE FRUITS (Group 12-12) - Continued**

### **Restrictions:**

- Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]
- Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]
- Maximum number of applications per year: [3 (at 4.56 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]
- [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]
- Minimum foliar retreatment interval: 5 days.
- Minimum soil retreatment interval: 30 days.
- **DO NOT** apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using [ground,] [aerial,] [or] chemigation equipment.
- Can be applied the day of harvest.
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.



## STRAWBERRY AND OTHER LOW-GROWING BERRIES (Subgroup 13-07G) (EXCEPT CRANBERRY)

Bearberry; Bilberry; Blueberry, Lowbush; Cloudberry; Lingonberry; Muntries; Partridgeberry; Strawberry; cultivars, varieties, and/or hybrids of these.

Disease/Pest Suppression	Application Rate	Application Instructions
Gray mold ( <i>Botrytis cinerea</i> )  Leaf spot ( <i>Mycosphaerella fragariae</i> )  Rhizopus fruit rot ( <i>Rhizopus</i> spp.)	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7-day interval.
Powdery mildew ( <i>Sphaerotheca macularis</i> ) ( <i>Podosphaera aphanis</i> )	3.2 to 6.84 fl oz/acre  (0.104 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7-day interval. [When disease pressure is severe, use the higher rates.]  [Drip applications are effective for suppression of this disease.]  [Use the high rate when applying via drip equipment.]
Crown rot/ Charcoal rot ( <i>Macrophomina</i> spp.)  Nematodes  Powdery mildew ( <i>Sphaerotheca macularis</i> ) ( <i>Podosphaera aphanis</i> )	4.0 to 6.84 fl oz/acre  (0.130 – 0.222 lb/acre fluopyram)	<b>Soil Applications</b> Apply specified dosage [in the following methods]: <ul style="list-style-type: none"> <li>• Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.</li> <li>• [Post-planting drench, or hill drench.]</li> <li>• [Transplant water drench with mechanical planting.]</li> <li>• [Transplant water drench with hand planting.]</li> </ul> Minimum 7-day interval between soil applications.

**STRAWBERRY AND OTHER LOW-GROWING BERRIES (Subgroup 13-07G) (EXCEPT CRANBERRY) - Continued**

**Restrictions:**

- Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]
- Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]
- Maximum number of applications per year: [4 (at 3.42 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]
- [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]
- Minimum retreatment interval: 7 days
- **DO NOT** apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using [ground], [aerial], [or] chemigation equipment.
- Can be applied broadcast the day of harvest (field use).
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.

SUGARBEET		
Disease/Pest Suppression	Application Rate	Application Instructions
Cercospora leaf spot ( <i>Cercospora beticola</i> )  Powdery mildew ( <i>Erysiphe polygoni</i> )	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 5- to 7-day interval. [When disease pressure is severe, use the higher rates and/or shorter intervals.]
Cercospora leaf spot ( <i>Cercospora beticola</i> )  Powdery mildew ( <i>Erysiphe polygoni</i> )  Nematodes	3.0 to 6.84 fl oz/acre  (0.098 – 0.222 lb/acre fluopyram)	<b>Soil Application</b> Apply specified dosage using any of the following methods: <ul style="list-style-type: none"> <li>[In-furrow spray during planting directed on or below seed.]</li> <li>[Chemigation using overhead irrigation equipment.]</li> </ul> Minimum 14-day interval between applications.
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]</li> <li>Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]</li> <li>Maximum number of applications per year: [4 (at 3.42 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]</li> <li>[Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]</li> <li>Minimum foliar retreatment interval: 5 days.</li> <li>Minimum soil retreatment interval: 14 days.</li> <li><b>DO NOT</b> apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).</li> <li>Apply using [ground,] [aerial] [or] chemigation equipment.</li> <li>For foliar uses: <b>DO NOT</b> apply LUNA PRIVILEGE within 7 days of harvest.</li> <li>For soil uses: <b>DO NOT</b> apply LUNA PRIVILEGE within 30 days of harvest.</li> <li>To limit the potential for development of disease resistance to this fungicide class, <b>DO NOT</b> make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.</li> <li>For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC Group.</li> <li>Tops or greens from this crop can be utilized for food or feed.</li> </ul>		

## SUNFLOWER (Subgroup 20B)

Calendula; Castor Oil Plant; Chinese Tallowtree; Euphorbia; Evening Primrose; Jojoba; Niger Seed; Rose Hip; Safflower; Stokes Aster; Sunflower; Tallowwood; Tea Oil Plant; Vernonia; cultivars, varieties, and/or hybrids of these.

Disease/Pest Suppression	Application Rate	Application Instructions
Alternaria leaf spot ( <i>Alternaria</i> spp.)	6.84 fl oz/acre	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval. [Can be applied in furrow.]
Nematodes	(0.222 lb/acre fluopyram)	
Powdery mildew ( <i>Erysiphe cichoracearum</i> )		
Sclerotinia wilt and head rot ( <i>Sclerotinia sclerotiorum</i> )		

### Restrictions:

- Maximum single application rate: 6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)
- Maximum annual application rate: 13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.
- Maximum number of applications per year: 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)
- Minimum retreatment interval: 14 days
- **DO NOT** apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using [ground,] [aerial,] [or] chemigation equipment.
- **DO NOT** apply LUNA PRIVILEGE within 14 days of harvest.
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.

TOBACCO		
Pest Suppression	Application Rate	Application Instructions
Nematodes	3.0 to 6.84 fl oz/acre  (0.098 – 0.222 lb/acre fluopyram)	<b>Soil Application</b> Apply specified dosage using any of the following methods: <ul style="list-style-type: none"> <li>• [Pre-plant banded or broadcast spray directed to the soil and incorporated during the bedding operation.]</li> <li>• [In-furrow drench.]</li> <li>• [In-furrow spray or in-furrow banded spray.]</li> <li>• [Transplant water drench with mechanical planting.]</li> <li>• [Post-plant banded spray directed to the soil at the base of plants and incorporated during cultivation operations.]</li> <li>• [Transplant water drench with hand planting.]</li> </ul>
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]</li> <li>• Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]</li> <li>• Maximum number of applications per year: [4 (at 3.42 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]</li> <li>• [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]</li> <li>• <b>DO NOT</b> apply more than 0.446 lbs fluopyram per acre per year, regardless of formulation or method of application.</li> <li>• Apply using [ground,] [aerial,] [or] chemigation equipment.</li> <li>• <b>DO NOT</b> apply LUNA PRIVILEGE within 30 days of harvest.</li> <li>• To limit the potential for development of disease resistance to this fungicide class, <b>DO NOT</b> make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.</li> <li>• For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.</li> </ul>		

## TREE NUTS (Group 14-12)

African Nut-Tree; Almond; Beechnut; Brazil Nut; Brazilian Pine; Bunya; Bur Oak; Butternut; Cajou Nut; Candlenut; Cashew; Chestnut; Chinquapin; Coconut; Coquito Nut; Dika Nut; Ginkgo; Guiana Chestnut; Hazelnut; Heartnut; Hickory Nut; Japanese Horse-Chestnut; Macadamia Nut; Mongongo Nut; Monkey-Pot; Monkey Puzzle Nut; Okari Nut; Pachira Nut; Peach Palm Nut; Pecan; Pequi; Pili Nut; Pine Nut; Pistachio; Sapucaia Nut; Tropical Almond; Walnut, Black; Walnut, English; Yellowhorn; Cultivars, varieties, and/or hybrids of these.

Disease/Pest Suppression	Application Rate	Application Instructions
Brown rot blossom blight <i>(Monilinia laxa)</i> <i>(Monilinia fructicola)</i>  Scab <i>(Cladosporium spp.)</i>  Shot hole <i>(Wilsonomyces carpophilus)</i>	3.2 to 6.84 fl oz/acre  (0.104 – 0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval.[ When disease pressure is severe, use the higher rates and/or shorter intervals.]
Powdery mildew <i>(Sphaeotheca pannosa)</i> <i>(Podosphaera tridactyla)</i> <i>(Microsphaera spp.)</i>	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. [When disease pressure is severe, use the shorter intervals.]  [Drip applications are effective for suppression of this disease.] [Use the high rate when applying via drip equipment.]
Alternaria <i>(Alternaria alternata)</i>  Anthracnose <i>(Colletotrichum acutatum)</i>  Blossom and shoot blight <i>(Botrytis cinerea)</i>  Botryosphaeria panicle and shoot blight <i>(Botryosphaeria dothidea)</i>  Jacket rot <i>(Botrytis cinerea)</i>  Septoria leaf spot <i>(Septoria pistaciarum)</i>	6.84 fl oz/acre  (0.222 lb/acre fluopyram)	[Apply at the critical timings for disease suppression.] Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. [When disease pressure is severe, use the shorter intervals.]

**TREE NUTS (Group 14-12) - Continued**

<b>Disease/Pest Suppression</b>	<b>Application Rate</b>	<b>Application Instructions</b>
<p>Nematodes</p> <p>Powdery mildew (<i>Sphaeotheca pannosa</i>) (<i>Podosphaera tridactyla</i>) (<i>Microsphaera</i> spp.)</p> <p>Septoria leaf spot (<i>Septoria pistaciarum</i>)</p>	<p>4.0 to 6.84 fl oz/acre</p> <p>(0.130 – 0.222 lb/acre fluopyram)</p>	<p><b>Soil Applications</b> - Apply specified dosage by chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment. For optimum results, apply to newly planted trees or those previously trained to drip, trickle or micro-sprinkler irrigation. Soil must be lightly pre-wetted to break soil surface tension prior to applications.</p> <p>Minimum 30-day interval between soil applications.</p>

**Restrictions:**

- Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]
- Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]
- Maximum number of applications per year: [4 (at 3.42 fl oz/acre of LUNA PRIVILEGE) or 2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]
- [Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]
- Minimum foliar retreatment interval: 7 days.
- Minimum soil retreatment interval: 30 days.
- **DO NOT** apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using [ground,] [aerial,] [or] chemigation equipment.
- **DO NOT** apply LUNA PRIVILEGE within 14 days of harvest.
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after LUNA PRIVILEGE be a product from a different FRAC group.

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in original container and keep tightly closed when not in use. Store in a cool dry place. Avoid cross-contamination with other pesticides.

**Pesticide Disposal:** Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of federal law. If these wastes cannot be disposed of by use according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

**Container Handling:**

### **Products in Non-Refillable Containers**

#### **Rigid, Non-refillable containers (equal to or less than 5 gallons)**

Non-refillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

#### **Dilutable Products in Rigid Non-refillable Containers that are Too Large to Shake (i.e., with capacities greater than 5 gallons or 50 lbs)**

Non-refillable container. **DO NOT** reuse or refill this container. Refer to Bottom Discharge Intermediate Bulk Container (IBC) or Top Discharge IBC, Drums, Kegs information as follows.

#### **Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)**

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

#### **Top Discharge IBC, Drums, Kegs (e.g.– Snyder 120 Next Gen, Bonar B120, Drums, Kegs)**

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. To triple rinse the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

#### **Products in Non-Refillable Fiber Drums with Liners**

Non-refillable container. **DO NOT** reuse or refill this container. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment, then offer for recycling if available or dispose of in a sanitary landfill or by incineration. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner.

#### **Products in Non-Rigid, Non-refillable Containers**

Nonrefillable container. **DO NOT** reuse or refill this container. Completely empty container into application equipment. Then offer for recycling if available or dispose of in a sanitary landfill or by other procedures approved by state and local authorities."



## **Products in Refillable Containers**

Refillable container. Refer to Bottom Discharge IBC or Top Discharge IBC, Drums, Kegs information as follows. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Contact your Ag retailer or Bayer CropScience for container return, disposal and recycling information.

### **Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)**

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

### **Top Discharge IBC, Drums, Kegs (e.g.– Snyder 120 Next Gen, Bonar B120, Drums, Kegs)**

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To triple rinse the containers before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by or by other procedures approved by state and local authorities.

End users are authorized to remove tamper evident cables as required to remove the product from the container unless the container is equipped with one way valves and refilling or returning is planned. If this is the case, end users are not authorized to remove tamper evident cables, one way valves or clean container.

## IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience LP. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSOURCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSOURCE LP SHALL NOT BE LIABLE FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSOURCE'S LP ELECTION, THE REPLACEMENT OF PRODUCT.

LUNA PRIVILEGE is specially formulated and sold by Bayer Group for the control of various pathogens according to the directions on this label. The purchase price of LUNA PRIVILEGE includes a prepaid license under which purchaser agrees to employ the purchased quantity of LUNA PRIVILEGE only for the above-specified uses and to provide notice of the terms and conditions of this license to any subsequent purchaser. Uses of LUNA PRIVILEGE other than those specified on this label are not licensed through the purchase of this product.

### NET CONTENTS:

### PRODUCED FOR



Bayer CropScience LP  
800 N. Lindbergh Blvd.  
St. Louis, MO 63167  
1-866-99BAYER (1-866-992-2937)

LUNA PRIVILEGE (PENDING) 06/12/2025



Bayer CropScience LP  
800 N. Lindbergh Blvd.  
St. Louis, MO 63167  
1-866-99BAYER (1-866-992-2937)

**LUNA® PRIVILEGE**  
[ABN: Velum® Prime, Velum® One, Velum®]

EPA Reg. No. 264-1078

For Use On: Coffee and Mint

*This supplemental label expires on 01/31/2026 and must not be used or distributed after this date.*

### Supplemental Label

## KEEP OUT OF REACH OF CHILDREN CAUTION

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Read this label and the product package label before using this product. The labeling must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the product label for LUNA® PRIVILEGE attached to the container. Use of LUNA® PRIVILEGE according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for LUNA® PRIVILEGE.

#### COFFEE

Pest Suppression	Application Rate	Application Instructions
Nematodes	13.7 fl oz/acre  (0.445 lb/acre fluopyram)	Apply at the critical timings for pest/disease suppression as a drench application. Refer to University and/or extension guidelines for best application timings.

#### Restrictions:

- Maximum single application rate: 13.7 fl oz/acre of LUNA PRIVILEGE (0.445 lb/acre fluopyram)
- Maximum annual application rate: 13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.
- Maximum number of applications per year: 1
- **DO NOT** apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using ground or chemigation equipment.
- **DO NOT** apply LUNA PRIVILEGE within 90 days of harvest.
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, the first foliar fungicide application after LUNA PRIVILEGE should be a product from a different FRAC group.

LUNA® and Velum® are Registered trademarks of Bayer.

Created on 11/18/2022

MINT (Spearmint and Peppermint, fresh and dried leaves)		
Pest Suppression	Application Rate	Application Instructions
Nematodes	6.0 to 6.84 fl oz/acre  (0.195 - 0.222 lb/acre fluopyram)	<p>Apply at the critical timings for pest/disease suppression. Refer to University and/or extension guidelines for best application timings. Continue as needed on a minimum 21-day interval.</p> <p>Apply specified dosage via overhead chemigation or as a foliar broadcast application in a minimum of 20 gallons per acre followed with ½ to ¾ inch of overhead irrigation within one hour of the application to move the product into the soil profile.</p>
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>Maximum single application rate: [6.84 fl oz/acre of LUNA PRIVILEGE (0.222 lb/acre fluopyram)]</li> <li>Maximum annual application rate: [13.7 fl oz of LUNA PRIVILEGE per acre (0.445 lb/acre fluopyram) per year.]</li> <li>Maximum number of applications per year: [2 (at 6.84 fl oz/acre of LUNA PRIVILEGE)]</li> <li>[Note to reviewer: the rates and number of applications listed in the above three bullets must be consistent with the rate(s) listed under 'Application Rate' above. The maximum single rate must not exceed 6.84 fl oz/acre and the maximum annual rate must not exceed 13.7 fl oz/acre of LUNA PRIVILEGE.]</li> <li><b>DO NOT</b> apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).</li> <li>Apply using ground or chemigation equipment.</li> <li><b>DO NOT</b> apply LUNA PRIVILEGE within 28 days of harvest.</li> <li>To limit the potential for development of disease resistance to this fungicide class, <b>DO NOT</b> make more than 2 sequential applications of LUNA PRIVILEGE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.</li> <li>For soil application, to limit the potential for development of disease resistance to this chemical class, the first foliar fungicide application after LUNA PRIVILEGE should be a product from a different FRAC group.</li> </ul>		

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

*As with any crop-protection product, always read and follow label directions.*

*For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*

LUNA® and Velum® are Registered trademarks of Bayer.

Created on 11/18/2022