



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

September 15, 2021

Marian Bleeke
Registration Manager
Bayer CropScience LP
800 Lindbergh Blvd.
St. Louis, MO 63167

Subject: Label Amendment – Increase preharvest interval to subgroup 13-07F
Product Name: Luna Experience
EPA Registration Number: 264-1091
Application Date: 08/03/2021
Decision Number: 577749

Dear Marian:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 264-1091
Decision No. 577749

with FIFRA section 6. If you have any questions, please contact Jennifer Drobish by phone at 703-347-8480, or via email at Drobish.jennifer@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Shaja B. Joyner". The signature is written in a cursive style with a large, looping initial "S".

Shaja B. Joyner, Product Manager 20
Fungicide-Herbicide Branch
Registration Division 7505P

Enclosure

FLUOPYRAM	GROUP	7	FUNGICIDE
TEBUCONAZOLE	GROUP	3	FUNGICIDE

LUNA[®] EXPERIENCE

Broad spectrum fungicide for control of plant diseases.

Almond; Barley; Brassica leafy greens (subgroup 5B); Bulb vegetables (group 3-07); Corn, field, grain; Corn, pop, grain; Corn, sweet; Cotton; Cucurbits (group 9); Fruiting Vegetables (group 8-10); Grapes; Hops; Dry and Succulent Beans; Peanut; Pistachio; Pome fruit (group 11); Soybean; Stone Fruits (group 12); Sunflower; Tree Nuts (group 14) and Wheat.

ACTIVE INGREDIENTS:

FLUOPYRAM*..... 17.6%

TEBUCONAZOLE*..... 17.6%

OTHER INGREDIENTS:..... 64.8%

TOTAL:..... **100.0%**

Contains 1.67 lbs FLUOPYRAM and 1.67 lbs TEBUCONAZOLE per gallon

*(CAS Numbers 658066-35-4 and 107534-96-3)

EPA Reg. No. 264-1091

EPA Est. No.

SUSPENSION CONCENTRATE

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

[See additional precautionary statements and directions for use on label.]

<p>ACCEPTED</p> <p>09/15/2021</p> <p>Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 264-1091</p>

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• DO NOT induce vomiting unless told to do so by a poison control center or doctor.• Have person sip a glass of water if able to swallow.• DO NOT give anything by mouth to an unconscious person.
IF ON SKIN:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
<p style="text-align: center;">For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577. In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	
NOTE TO PHYSICIAN: Treat Symptomatically.	

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, absorbed through skin or inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE):

Applicators and other handlers must wear long-sleeved shirt and long pants, shoes plus socks, and chemical-resistant (such as nitrile or butyl) gloves.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- User should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses: This pesticide is toxic to mammals, fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when disposing of equipment wash water or rinsate. This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of Fluopyram. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff.

This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

DIRECTIONS FOR USE
STOP - READ THE LABEL BEFORE USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Not for sale, distribution, or use in Nassau and Suffolk counties, New York except as permitted under FIFRA 24(c), Special Local Need registration.

Aerial Application Prohibited in New York State.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during **the restricted-entry interval (REI) of 12 hours for all crops except when performing cane tying, turning, or girdling on grapes.**
The REI is 5 days for treated grapes when conducting cane tying, turning, or girdling on grapes. The REI is 12 hours for all other activities in grapes.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- coveralls over long-sleeved shirt and long pants
- socks and shoes
- chemical-resistant gloves made of any waterproof material such as natural rubber \geq 14 mils

PRODUCT INFORMATION

LUNA® EXPERIENCE is a broad-spectrum fungicide with preventative, systemic and curative properties labeled for the control or suppression of certain crop diseases.

LABELED USES

Almond; Barley; Brassica leafy greens (subgroup 5B); Bulb vegetables (group 3-07); Corn, field, grain; Corn, pop, grain; Corn, sweet; Cotton; Cucurbits (group 9); Fruiting Vegetables (group 8-10); Grapes; Hops; Dry and Succulent Beans; Peanut; Pistachio; Pome fruit (group 11); Soybean; Stone Fruits (group 12); Sunflower; Tree Nuts (group 14) and Wheat.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, please note that LUNA EXPERIENCE contains both a Group 7 and Group 3 fungicide. Any fungal population may contain individuals naturally resistant to LUNA EXPERIENCE and other Group 7 or Group 3 fungicides. A gradual or total loss of pest control may occur over time if these (fungicides) are used repeatedly in the same fields. Appropriate resistance-management strategies must be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of LUNA EXPERIENCE or other Group 7 or Group 3 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal/bacterial populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Bayer CropScience at 1-866-99BAYER (1-866-992-2937). You can also contact your pesticide distributor or university extension specialist to report resistance.

MANDATORY SPRAY DRIFT

Aerial Applications

- For aerial applications, **DO NOT** apply when wind speeds exceed 10 mph at the application site.
- For all applications, applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- **DO NOT** release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply during temperature inversions.

Airblast applications

- Sprays must be directed into the canopy.
- For all applications, applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- User must turn off outward pointing nozzles at row ends and when spraying outer row.
- **DO NOT** apply during temperature inversions.

Ground Boom Applications

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- For all applications, applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

- THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
- BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles - Follow nozzle manufacturer's recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT – Aircraft

Higher release heights increase the potential for spray drift. When applying aerially to crops, do not release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversion.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

BOOM-LESS GROUND APPLICATIONS

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

HANDHELD TECHNOLOGY APPLICATIONS

Take precautions to minimize spray drift.

APPLICATION INFORMATION

Use sufficient water volume to provide thorough and uniform coverage to obtain the most effective disease control.

Ground Application

For optimum disease control, apply in sufficient water to ensure thorough coverage of foliage, bloom, and fruit.

Aerial Application

For aerial application equipment, a minimum of **10** gallons of water per acre for tree crops and **2** gallons of water per acre for field and vegetable crops is required. Aerial Application Prohibited in New York State.

Air-Blast Application

Air-assisted or air-blast sprayers move spray droplets into the crop canopy using a forced-air system. The fan should be set up to deliver only enough air volume to penetrate the canopy and provide good coverage. Adjust deflectors or other aiming devices to direct spray only to the target area. Equip sprayers with nozzles that provide accurate and uniform application.

Chemigation Application

Apply this product only through center pivot, motorized-lateral move, traveling gun, solid set or portable (wheelmove, side roll, end tow, or hand move) irrigation systems. **DO NOT** apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, must shut the system down and make necessary adjustments should the need arise.

LUNA EXPERIENCE has not been sufficiently tested when applied through irrigation systems to assure consistent product performance for all labeled uses. The following application techniques are provided for user reference but do not constitute a warranty of fitness for application through sprinkler irrigation equipment.

DO NOT connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label prescribed safety devices for public water systems are in place. 'Public water system' means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ), back flow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an alternative to the RPZ, the water from the public water system must be discharged into a reservoir tank prior to pesticide introduction.

There must be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. Pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The systems must contain functional interlocking controls, to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed

and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

DO NOT apply when wind speed favors drift. Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur. Apply pesticide continuously for the duration of the water application. For mixing instructions, please refer to specific use directions in the “Spray mixing and compatibility” section.

This product may be used through two basic types of irrigation systems as outlined in **Sections A and B** below. The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. Determine which type of irrigation system is in place, then refer to the appropriate directions provided below for each type. See crops section on the label for required treatment rates and additional use information.

A. Center Pivot, Motorized-Lateral Move and Traveling Gun Irrigation Equipment

For injections of pesticides, these continuously moving systems must use a positive displacement injection pump of either diaphragm or piston type and be constructed of materials that are compatible with pesticides. They must also be capable of being fitted with a system interlock and capable of injection at pressures approximately 2-3 times those encountered within the irrigation water line. Venturi applicator units cannot be used on these systems. Thoroughly mix required amount of this product for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run. Mixture in the chemical supply tank must be continuously agitated during the injection run. Shut off injection equipment after one revolution or run, but continue to operate irrigation system until this product has been cleared from the last sprinkler head.

B. Solid-Set, Portable (Wheel Move, Side Roll, End Tow, or Hand Move) Equipment

With stationary systems, an effectively designed in-line Venturi applicator unit is preferred to support even and quick distribution. However, a positive-displacement pump can also be used. For solid set systems, determine acreage covered by sprinkler. Fill the tank of injection equipment with water and adjust flow to use contents over 30 to 45 minutes. Mix desired amount of this product for acreage to be covered with water so that the total mixture of this product plus water in the injection tank is equal to the quantity of water used during calibration. Provide chemical supply tank agitation sufficient for mixing until chemigation is completed. Operate entire system at normal pressures recommended by the manufacturer of injection equipment used, for amount of time established during calibration. This product can be injected during the irrigation cycle or as a separate application. Stop injection equipment with any system after treatment is completed and continue to operate irrigation system until this product has been cleared from the last sprinkler head.

SPRAY MIXING AND COMPATIBILITY

Begin with clean spray equipment and add one-half of the required amount of water to the spray or mixing tank and start agitation. Add the required quantity of fungicide and the tank-mix partner if applicable to the water and complete filling with water to the required total volume. Follow the recommendations of your State Cooperative Extension Service for tank-mixing with other products. In general, follow the order beginning first with water conditioners, water soluble packaging (wait for it to completely dissolve), wettable powders and water-dispersible granular products, liquid flowables and suspension concentrates, emulsifiable concentrates, and adjuvants last. Maintain agitation throughout spraying. **DO NOT** allow spray mixture to remain in the tank overnight, or for long periods during the day without agitation.

LUNA EXPERIENCE is physically compatible with most commonly used fungicide, herbicide, insecticide, and foliar nutrient products. However, the compatibility of LUNA EXPERIENCE with all potential tank-mix partners has not been fully investigated. If tank mixing with other pesticides is desirable, conduct a jar test with the

volumes and rates typically used in agricultural application. Using a small container of water, add the proportionate amounts of the products: wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 15 minutes. Look for signs of separation, globules, sludge, flakes, or other precipitates. Physical compatibility is indicated if the combination remains mixed or can be remixed readily.

The crop safety of all potential tank-mixes with LUNA EXPERIENCE has not been tested on all crops listed on the label. Before applying any tank-mixture not specified on this label, safety to the target crop must be confirmed on a small portion of the crop listed on the label to be treated to ensure an adverse response will not occur.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

PRODUCT RESTRICTIONS AND LIMITATIONS

DO NOT apply more than the maximum yearly rate for each specific crop from any combination of products containing FLUOPYRAM or TEBUCONAZOLE.

ROTATIONAL CROP RESTRICTIONS

Areas treated with this product can be replanted immediately following harvest with any crop for which both a FLUOPYRAM and a TEBUCONAZOLE tolerance exist. This includes: Almond; Barley; Brassica leafy greens (subgroup 5B); Bulb vegetables (group 3-07); Corn, field, grain; Corn, pop, grain; Corn, sweet; Cotton; Cucurbits (group 9); Dry and Succulent beans; Fruiting Vegetables (group 8-10); Grapes; Hops; Peanut; Pecan; Pistachio; Pome fruit (group 11); Soybean; Stone Fruits (group 12); Sunflower; Tree Nuts (group 14) and Wheat.

The following crops can be planted into treated areas 120 days after the last application of LUNA EXPERIENCE:

Alfalfa; Artichoke,(Globe); Brassica leafy greens subgroup 5A; Carrot; Cereals (except corn and rice) (group 15) [which include: buckwheat, millet (pearl and proso), oats, rye, sorghum, teosinte, and triticale]; Citrus (group 10-10); Dill seed; Ginseng; Herb (subgroup 19A); Leafy vegetables (except watercress) (group 4); Potato and other root, tuberous and corm vegetables (except sugarbeet) (subgroups 1B and 1C); Rapeseed (subgroup 20A); Small Berries (caneberries and bushberries) (subgroups 13-07A and 13-07B); Strawberry and other low-growing berries, except cranberry (subgroup 13-07G); Succulent Shelled Pea and Bean subgroup 6B; Sugarcane (in Region 3); and Sugarbeet.

DO NOT rotate to crops other than those listed above.

USE DIRECTIONS FOR SPECIFIC CROPS

ALMOND		
Disease Control	Application Rate	Product Instructions
Brown rot blossom blight <i>(Monilinia laxa)</i> <i>(Monilinia fructicola)</i> Shot hole <i>(Wilsonomyces carpophilus)</i> Powdery mildew <i>(Sphaerotheca pannosa)</i> <i>(Podosphaera tridactyla)</i> Jacket rot ^[1] Green fruit rot <i>(Botrytis cinerea)</i> Alternaria leaf spot ^[1] <i>(Alternaria alternata)</i> Anthracnose ^[1] <i>(Colletotrichum acutatum)</i> Scab ^[1] <i>(Cladosporium carpophilum)</i> Rust ^[1] <i>(Tranzschelia discolor)</i>	6.0 to 17 fl oz/acre (0.078 – 0.222 lb/acre fluopyram) (0.078 – 0.222 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Disease Suppression	Application Rate	Product Instructions
Hull rot ^[1] <i>(Rhizopus spp.)</i> <i>(Monilinia spp.)</i>	6.0 to 17 fl oz/acre (0.078 – 0.222 lb/acre fluopyram) (0.078 – 0.222 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Restrictions: <ul style="list-style-type: none"> Maximum single application rate: 17 fl oz/acre of LUNA EXPERIENCE (0.222 lb/acre fluopyram and 0.222 lb/acre tebuconazole). 		

- Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 5 (at 6.0 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 7 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.900 lbs tebuconazole per acre per year, including soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE within 35 days of harvest.
- **DO NOT** cut cover crops in treated areas for feed, or allow livestock to graze treated areas.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

[¹Not for use in CA [without a supplemental label.]]

BARLEY		
Disease Control	Application Rate	Product Instructions
Powdery mildew (<i>Erysiphe graminis</i>)	8.6 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings.
Septoria leaf and glume blotch (<i>Septoria</i> spp.)	(0.112 lb/acre fluopyram)	
Rusts (<i>Puccinia</i> spp.)	(0.112 lb/acre tebuconazole)	

Restrictions:

- Maximum single application rate: 8.6 fl oz/acre of LUNA EXPERIENCE (0.112 lb/acre fluopyram and 0.112 lb/acre tebuconazole).
- Maximum annual application rate: 8.6 fl oz/acre of LUNA EXPERIENCE (0.112 lb/acre fluopyram and 0.112 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 1
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.112 lbs tebuconazole per acre per year, including soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE within 30 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

BRASSICA LEAFY GREENS (Subgroup 5B)

Broccoli Raab; Cabbage, Chinese (Bok Choy); Collards; Kale; Mizuna; Mustard Greens; Mustard Spinach; Rape Greens.

Disease Control	Application Rate	Product Instructions
Powdery mildew ^[1] (<i>Erysiphe cruciferarum</i>) (<i>Erysiphe polygoni</i>) Cercospora leaf spot ^[1] (<i>Cercospora brassicola</i>) Alternaria blight / Black spot ¹ (<i>Alternaria</i> spp.)	6.0 to 8.6 fl oz/acre (0.078 – 0.112 lb/acre fluopyram) (0.078 – 0.112 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Botrytis gray mold (<i>Botrytis cinerea</i>) Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i>) (<i>Sclerotinia minor</i>)	8.6 fl oz/acre (0.112 lb/acre fluopyram) (0.112 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the shorter intervals.
Disease Suppression	Application Rate	Product Instructions
Rhizoctonia bottom rot (<i>Rhizoctonia solani</i>)	8.6 fl oz/acre (0.112 lb/acre fluopyram) (0.112 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the shorter intervals.

Restrictions:

- Maximum single application rate: 8.6 fl oz/acre of LUNA EXPERIENCE (0.112 lb/acre fluopyram and 0.112 lb/acre tebuconazole).
- Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 5 (at 6.0 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 7 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.450 lbs tebuconazole per acre per year, including soil and foliar uses.

- **DO NOT** apply LUNA EXPERIENCE within 7 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.
- Application to Turnip greens is limited to east of the Rockies.

[¹Not for use in CA [without a supplemental label.]]

BULB VEGETABLES (group 3-07)

Onion, bulb subgroup: Daylily, Bulb; Fritillaria, Bulb; Garlic, Bulb; Garlic, Great-Headed, Bulb; Garlic, Serpent, Bulb; Lily, Bulb; Onion, Bulb; Onion, Chinese, Bulb; Onion, Pearl; Onion, Potato, Bulb; Shallot, Bulb; cultivars, varieties, and/or hybrids of these.

Onion, green subgroup: Chive, Fresh Leaves; Chive, Chinese, Fresh Leaves; Elegans Hosta; Fritillaria, Leaves; Kurrat; Lady's Leek; Leek; Leek, Wild; Onion, Beltsville Bunching; Onion, Fresh; Onion, Green; Onion, Macrostem; Onion, Tree, Tops; Onion, Welsh, Tops; Shallot, Fresh Leaves; cultivars, varieties, and/or hybrids of these.

Disease Control	Application Rate	Product Instructions
Botrytis leaf blight, neck rot <i>(Botrytis squamosa)</i> <i>(Botrytis allii)</i> Purple blotch <i>(Alternaria porri)</i> Rust <i>(Puccinia allii)</i> <i>(Puccinia porri)</i>	8.0 to 12.8 fl oz/acre (0.104 – 0.167 lb/acre fluopyram) (0.104 – 0.167 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Stemphylium leaf blight / stalk rot <i>(Stemphylium vesicarium)</i> Cladosporium leaf blotch <i>(Cladosporium allii)</i>	12.8 fl oz/acre (0.167 lb/acre fluopyram) (0.167 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Disease Suppression	Application Rate	Product Instructions

White rot (<i>Sclerotium cepivorum</i>)	12.8 fl oz/acre (0.167 lb/acre fluopyram) (0.167 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
--	---	--

Restrictions:

- Maximum single application rate: 12.8 fl oz/acre of LUNA EXPERIENCE (0.167 lb/acre fluopyram and 0.167 lb/acre tebuconazole).
- Maximum annual application rate: 25.6 fl oz/acre of LUNA EXPERIENCE (0.334 lb/acre fluopyram and 0.334 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 3 (at 8.0 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 10 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.335 lbs tebuconazole per acre per year, including soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE within 7 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

CORN, FIELD, GRAIN; CORN, POP, GRAIN; CORN, SWEET

Sweet Corn; Field Corn; Field Corn Grown For Seed, and Popcorn

Disease Control	Application Rate	Product Instructions
Northern leaf blight <i>(Helminthosporium turcicum)</i> Southern leaf blight <i>(Helminthosporium maydis)</i> Rusts <i>(Puccinia spp.)</i>	12.8 fl oz/acre (0.167 lb/acre fluopyram) (0.167 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval.

Restrictions:

- Maximum single application rate: 12.8 fl oz/acre of LUNA EXPERIENCE (0.167 lb/acre fluopyram and 0.167 lb/acre tebuconazole).
- Maximum annual application rate: 25.6 fl oz/acre of LUNA EXPERIENCE (0.334 lb/acre fluopyram and 0.334 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 2
- Minimum retreatment interval: 14 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.691 lbs tebuconazole per acre per year, including soil and foliar uses.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.
- **Sweet corn:** may be applied up to 7 days before the harvest of ears or forage, and 49 days before the harvest of fodder.
- **Field, seed or popcorn:** may be applied up to 21 days before the harvest of forage, and 36 days before the harvest of grain or fodder.

COTTON		
Disease Control	Application Rate	Product Instructions
Southwestern cotton rust ^[1] (<i>Puccinia cacabata</i>)	17 fl oz/acre (0.222 lb/acre fluopyram) (0.222 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval.
<p>Restrictions:</p> <ul style="list-style-type: none"> • Maximum single application rate: 17 fl oz/acre of LUNA EXPERIENCE (0.222 lb/acre fluopyram and 0.222 lb/acre tebuconazole). • Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year. • Maximum number of applications per year: 2 • Minimum retreatment interval: 7 days • Apply using ground, aerial, or chemigation equipment. • Regardless of formulation or method of application, DO NOT apply more than 0.446 lbs Fluopyram or 0.691 lbs tebuconazole per acre per year, including seed treatment, soil and foliar uses. • DO NOT apply LUNA EXPERIENCE within 30 days of harvest. • To limit the potential for development of disease resistance to these fungicide classes, DO NOT make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group. <p>[¹Not for use in CA [without a supplemental label.]]</p>		

CUCURBITS (Group 9)

Melon subgroup: Citron Melon; Muskmelon; Watermelon.

Squash/Cucumber subgroup: Chayote (Fruit); Chinese Waxgourd; Cucumber; Gherkin; Gourd, Edible; *Momordica* spp.; Pumpkin; Squash, Summer; Squash, Winter.

Disease Control	Application Rate	Product Instructions
Powdery mildew <i>(Sphaerotheca fuliginea / Podosphaera xanthii)</i> <i>(Erysiphe cichoracearum)</i> Alternaria leaf spot <i>(Alternaria cucumerina)</i>	6.0 to 17 fl oz/acre (0.078 – 0.222 lb/acre fluopyram) (0.078 – 0.222 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Gummy stem blight ^[1] <i>(Didymella bryonae)</i> Belly rot ^[1] <i>(Rhizoctonia solani)</i> Anthracnose ^[1] <i>(Colletotrichum spp.)</i>	10.0 to 17 fl oz/acre (0.130 - 0.222 lb/acre fluopyram) (0.130 - 0.222 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.

Note: A mild yellowing on leaf margins is sometimes seen following application of LUNA EXPERIENCE in cucurbits.

Restrictions:

- Maximum single application rate: 17 fl oz/acre of LUNA EXPERIENCE (0.222 lb/acre fluopyram and 0.222 lb/acre tebuconazole).
- Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 5 (at 6.0 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 10 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.691 lbs tebuconazole per acre per year, including soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE within 7 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing

fungicide before rotating with a fungicide from a different Group.

[¹Not for use in CA [without a supplemental label.]]

FRUITING VEGETABLES (Group 8-10)

Tomato subgroup: Bush Tomato; Cocona; Currant Tomato; Garden Huckleberry; Goji Berry; Groundcherry; Naranjilla; Sunberry; Tomatillo; Tomato; Tree Tomato; cultivars, varieties, and/or hybrids of these.

Pepper/Eggplant subgroup: African Eggplant; Bell Pepper; Eggplant; Martynia; Nonbell Pepper; Okra; Pea Eggplant; Pepino; Roselle; Scarlet Eggplant; cultivars, varieties, and/or hybrids of these.

Disease Control	Application Rate	Product Instructions
Early blight (<i>Alternaria solani</i>)	6.0 to 17 fl oz/acre (0.078 – 0.222 lb/acre fluopyram) (0.078 – 0.222 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.

Restrictions:

- Maximum single application rate: 17 fl oz/acre of LUNA EXPERIENCE (0.222 lb/acre fluopyram and 0.222 lb/acre tebuconazole).
- Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 5 (at 6.0 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 7 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 1.35 lbs tebuconazole per acre per year, including soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE within 7 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

GRAPES		
Disease Control	Application Rate	Product Instructions
Powdery mildew (<i>Uncinula necator</i>)	6.0 to 8.6 fl oz/acre (0.078 – 0.112 lb/acre fluopyram) (0.078 – 0.112 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 12- to 21-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Botrytis bunch rot / Gray mold (<i>Botrytis cinerea</i>) Black rot (<i>Guignardia bidwellii</i>) Sour rot ^[1] (<i>various pathogens</i>)	8.0 to 8.6 fl oz/acre (0.104 – 0.112 lb/acre fluopyram) (0.104 – 0.112 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 12- to 21-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Disease Suppression	Application Rate	Product Instructions
Phomopsis cane and leaf spot (<i>Phomopsis viticola</i>)	8.6 fl oz/acre (0.112 lb/acre fluopyram) (0.112 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 12- to 21-day interval. When disease pressure is severe, use the shorter intervals.
Restrictions:		
<ul style="list-style-type: none"> • Maximum single application rate: 8.6 fl oz/acre of LUNA EXPERIENCE (0.112 lb/acre fluopyram and 0.112 lb/acre tebuconazole). • Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year. • Maximum number of applications per year: 5 (at 6.0 fl oz/acre of LUNA EXPERIENCE) • Minimum retreatment interval: 12 days • Apply using ground, aerial, or chemigation equipment. • Regardless of formulation or method of application, DO NOT apply more than 0.446 lbs Fluopyram or 0.900 lbs tebuconazole per acre per year, including soil and foliar uses. • DO NOT apply LUNA EXPERIENCE after BBCH 73 stage (when bunches begin to hang) or within 45 days of harvest. 		

- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

[¹Not for use in CA [without a supplemental label.]]

HOPS		
Disease Control	Application Rate	Product Instructions
Powdery mildew (<i>Sphaerotheca humuli</i>) (<i>Sphaerotheca macularis</i>)	8.0 to 17 fl oz/acre (0.104 - 0.222 lb/acre fluopyram) (0.104 - 0.222 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval. Make applications in sufficient water volume (just to the point of runoff). Thorough coverage is needed for disease control. Apply increasing product rate as plant size increases. 8.0 fl oz with 15 - 30 gals./acre 10.0 fl oz with 31 - 60 gals./acre 12.0 fl oz with 61 - 90 gals./acre 14.0 fl oz with 91 - 200 gals./acre
Disease Suppression	Application Rate	Product Instructions
Gray mold (<i>Botrytis</i> spp.)	17 fl oz/acre (0.222 lb/acre fluopyram) (0.222 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval.

Restrictions:

- Maximum single application rate: 17 fl oz/acre of LUNA EXPERIENCE (0.222 lb/acre fluopyram and 0.222 lb/acre tebuconazole).
- Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 4 (at 8.0 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 14 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.900 lbs tebuconazole per acre per year, including soil and foliar uses.
- DO NOT** apply LUNA EXPERIENCE within 14 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing

fungicide before rotating with a fungicide from a different Group.

DRY AND SUCCULENT BEANS^[1]

Disease Control	Application Rate	Product Instructions
Powdery mildew (<i>Erysiphe graminis</i>)	8.0 to 12.8 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval.
Septoria leaf and glume blotch (<i>Septoria</i> spp.)	(0.104 – 0.167 lb/acre fluopyram)	When disease pressure is severe, use the higher rates and/or shorter intervals.
Rusts (<i>Puccinia</i> spp.)	(0.104 – 0.167 lb/acre tebuconazole)	

Restrictions:

- Maximum single application rate: 12.8 fl oz/acre of LUNA EXPERIENCE (0.167 lb/acre fluopyram and 0.167 lb/acre tebuconazole).
- Maximum annual application rate: 25.6 fl oz/acre of LUNA EXPERIENCE (0.334 lb/acre fluopyram and 0.334 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 3 (at 8.0 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 10 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.672 lbs tebuconazole per acre per year, including soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE within 14 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.
- **DO NOT** allow livestock to graze treated area for 14 days and **DO NOT** harvest for food or feed within 14 days of application.

[¹Not for use in CA [without a supplemental label.]]

OKRA		
Disease Control	Application Rate	Product Instructions
Cercospora leaf spot (<i>Cercospora</i> spp.)	12.8 fl oz/acre (0.167 lb/acre fluopyram) (0.167 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval.
<p>Restrictions:</p> <ul style="list-style-type: none"> • Maximum single application rate: 12.8 fl oz/acre of LUNA EXPERIENCE (0.167 lb/acre fluopyram and 0.167 lb/acre tebuconazole). • Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year. • Maximum number of applications per year: 2 • Minimum retreatment interval: 14 days • Apply using ground, aerial, or chemigation equipment. • Regardless of formulation or method of application, DO NOT apply more than 0.446 lbs Fluopyram or 0.675 lbs tebuconazole per acre per year including soil and foliar uses. • DO NOT apply LUNA EXPERIENCE within 3 days of harvest. • To limit the potential for development of disease resistance to these fungicide classes, DO NOT make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group. 		

PEANUT		
Disease Control	Application Rate	Product Instructions
Sclerotium rot, White mold, Southern blight, Southern stem rot <i>(Sclerotium rolfsii)</i> Rhizoctonia limb rot, Peg rot, Pod rot <i>(Rhizoctonia solani)</i>	15.7 fl oz/acre (0.205 lb/acre fluopyram) (0.205 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval.
Early leaf spot <i>(Cercospora arachidicola)</i> Late leaf spot <i>(Cercosporidium personatum)</i> Leaf rust <i>(Puccinia arachidis)</i> Web blotch <i>(Phoma arachidicola)</i> Leaf scorch and Pepper spot <i>(Leptosphaerulina crassiasca)</i>	12 to 15.7 fl oz/acre (0.157 – 0.205 lb/acre fluopyram) (0.157 – 0.205 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval.
Disease Suppression	Application Rate	Product Instructions
Sclerotinia blight <i>(Sclerotinia minor)</i> Cylindrocadium black rot <i>(Cylindrocladium crotalariae)</i>	15.7 fl oz/acre (0.205 lb/acre fluopyram) (0.205 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval.
Restrictions: <ul style="list-style-type: none"> • Maximum single application rate: 15.7 fl oz/acre of LUNA EXPERIENCE (0.205 lb/acre fluopyram and 0.205 lb/acre tebuconazole). • Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year. • Maximum number of applications per year: 2 (at 15.7 fl oz/acre of LUNA EXPERIENCE) 		

- Minimum retreatment interval: 14 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.810 lbs tebuconazole per acre per year, including seed treatment, soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE within 14 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.
- **DO NOT** feed hay or threshings or allow livestock to graze in treated areas.

PECAN		
Disease Control	Application Rate	Product Instructions
Scab (<i>Cladosporium caryigenum</i>)	6.0 to 17 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval. When disease pressure is severe, use the higher rates.
Brown leaf spot ^[1] (<i>Sirosporium diffusium</i>)	(0.078 – 0.222 lb/acre fluopyram)	
Downy spot ^[1] (<i>Mycosphaerella caryigena</i>)	(0.078 – 0.222 lb/acre tebuconazole)	
Liver spot ^[1] (<i>Gnomonia caryae</i>)		
Vein spot ^[1] (<i>Gnomonia nerviseda</i>)		
Zonate leaf spot ^[1] (<i>Grovesinia pyramidalis</i>)		
Powdery mildew ^[1] (<i>Microsphaera</i> spp.)		
Restrictions:		
<ul style="list-style-type: none"> • Maximum single application rate: 17 fl oz/acre of LUNA EXPERIENCE (0.222 lb/acre fluopyram and 0.222 lb/acre tebuconazole). • Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year. • Maximum number of applications per year: 5 (at 6.0 fl oz/acre of LUNA EXPERIENCE) • Minimum retreatment interval: 14 days • Apply using ground, aerial, or chemigation equipment. 		

- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.900 lbs tebuconazole per acre per year, including soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE after shucks begin to split.
- **DO NOT** cut cover crops in treated areas for feed or allow livestock to graze treated areas.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

[¹Not for use in CA [without a supplemental label.]]

PISTACHIO

Disease Control	Application Rate	Product Instructions
Botryosphaeria panicle and shoot blight (<i>Botryosphaeria dothidea</i>)	6.0 to 17 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Septoria leaf spot (<i>Septoria pistaciarum</i>)	(0.078 – 0.222 lb/acre fluopyram)	
Blossom and shoot blight (<i>Botrytis cinerea</i>)	(0.078 – 0.222 lb/acre tebuconazole)	
Alternaria late blight (<i>Alternaria alternata</i>)		

Restrictions:

- Maximum single application rate: 17 fl oz/acre of LUNA EXPERIENCE (0.222 lb/acre fluopyram and 0.222 lb/acre tebuconazole).
- Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 5 (at 6.0 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 10 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.900 lbs tebuconazole per acre per year, including soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE within 35 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.
- **DO NOT** cut cover crops in treated areas for feed, or allow livestock to graze treated areas.

POME FRUIT (Group 11)

Apple, Crabapple, Loquat, Mayhaw, Pear, Oriental Pear, and Quince.

Disease Control	Application Rate	Product Instructions
Powdery mildew (<i>Podosphaera leucotricha</i>)	5.0 to 10 fl oz/acre (0.065 – 0.130 lb/acre fluopyram) (0.065 – 0.130 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Scab (<i>Venturia</i> spp.) Cedar apple rust (<i>Gymnosporangium juniperi-virginianae</i>)	8.0 to 10 fl oz/acre (0.104 – 0.130 lb/acre fluopyram) (0.104 – 0.130 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Disease Suppression	Application Rate	Product Instructions
Bitter rot (<i>Glomerella cingulata</i>) Sooty blotch (<i>Gloeodes pomigena</i>) Fly speck (<i>Schizothyrium pomi</i>)	8.0 to 10 fl oz/acre (0.104 – 0.130 lb/acre fluopyram) (0.104 – 0.130 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.

Restrictions:

- Maximum single application rate: 10 fl oz/acre of LUNA EXPERIENCE (0.130 lb/acre fluopyram and 0.130 lb/acre tebuconazole).
- Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 6 (at 5.0 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 7 days
- Apply using ground, aerial, or chemigation equipment.

- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 1.35 lbs tebuconazole per acre per year, including soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE within 75 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

SOYBEAN		
Disease Control	Application Rate	Product Instructions
Rust (<i>Uromyces appendiculatus</i>) Powdery mildew (<i>Microsphaera diffusa</i>)	8.6 fl oz/acre (0.112 lb/acre fluopyram) (0.112 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the shorter intervals.
Disease Suppression	Application Rate	Product Instructions
Pod and stem blight (<i>Diaporthe phaseolorum</i>)	8.6 fl oz/acre (0.112 lb/acre fluopyram) (0.112 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the shorter intervals.

Restrictions:

- Maximum single application rate: 8.6 fl oz/acre of LUNA EXPERIENCE (0.167 lb/acre fluopyram and 0.167 lb/acre tebuconazole).
- Maximum annual application rate: 25.8 fl oz/acre of LUNA EXPERIENCE (0.337 lb/acre fluopyram and 0.337 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 3
- Minimum retreatment interval: 10 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.337 lbs tebuconazole per acre per year, including seed treatment, soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE within 21 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing

fungicide before rotating with a fungicide from a different Group.

- **DO NOT** allow livestock to graze soybean and **DO NOT** harvest soybean hay for food or feed.

STONE FRUITS (Group 12)

Apricot; Cherry, sweet; Cherry, tart; Nectarine; Peach; Plum ; Plum, Chickasaw; Plum, Damson; Plum, Japanese; Plumcot; Prune (fresh).

Disease Control	Application Rate	Product Instructions
Brown rot blossom blight Fruit rot (<i>Monilinia</i> spp.) Powdery mildew Rusty spot (<i>Podosphaera</i> spp.) (<i>Sphaerotheca pannosa</i>) Cherry leaf spot (<i>Blumeriella jaapii</i>) Shot hole ^[1] (<i>Wilsonomyces carpophilus</i>) Jacket rot, Green fruit rot ^[1] (<i>Botrytis cinerea</i>) Peach leaf curl ^[1] (<i>Taphrina</i> spp.)	6.0 to 10 fl oz/acre (0.078 – 0.130 lb/acre fluopyram) (0.078 – 0.130 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Scab (<i>Cladosporium carpophilum</i>) Rust (<i>Tranzschelia discolor</i>)	8.0 to 10 fl oz/acre (0.104 – 0.130 lb/acre fluopyram) (0.104 – 0.130 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.

Restrictions:

- Maximum single application rate: 10 fl oz/acre of LUNA EXPERIENCE (0.130 lb/acre fluopyram and 0.130 lb/acre tebuconazole).
- Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole).

0.444 lb/acre tebuconazole) per year.

- Maximum number of applications per year: 5 (at 6.0 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 7 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 1.35 lbs tebuconazole per acre per year, including soil and foliar uses.
- May be applied the day of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

[¹Not for use in CA [without a supplemental label.]]

SUNFLOWER

Disease Control	Application Rate	Product Instructions
Rust (<i>Puccinia helianthi</i>)	9.0 to 12.8 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval. When disease pressure is severe, use the higher rates.
Alternaria leaf spot (<i>Alternaria</i> spp.)	(0.117 – 0.167 lb/acre fluopyram)	
Sclerotinia wilt and head rot (<i>Sclerotinia sclerotiorum</i>)	(0.117 – 0.167 lb/acre tebuconazole)	
Powdery mildew (<i>Erysiphe cichoracearum</i>)		

Restrictions:

- Maximum single application rate: 12.8 fl oz/acre of LUNA EXPERIENCE (0.167 lb/acre fluopyram and 0.167 lb/acre tebuconazole).
- Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 3 (at 9.0 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 14 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.450 lbs tebuconazole per acre per year, including soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE within 50 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing

fungicide before rotating with a fungicide from a different Group.

TREE NUTS (Group 14)

(See separate use directions for: Almonds, Pecan, and Pistachio)

Beech nut; Brazil nut; Butternut; Cashew; Chestnut; Chinquapin; Filbert (hazelnut); Hickory nut; Macadamia nut (bush nut); Walnut, black and English (Persian).

Disease Control	Application Rate	Product Instructions
Botryosphaeria panicle and shoot blight <i>(Botryosphaeria dothidea)</i> Brown rot blossom blight <i>(Monilinia laxa)</i> <i>(Monilinia fructicola)</i> Alternaria leaf and fruit spot <i>(Alternaria alternata)</i> Alternaria late blight <i>(Alternaria alternata)</i> Scab <i>(Cladosporium carpophilum)</i> <i>(Cladosporium caryigenum)</i> Shothole <i>(Wilsonomyces carpophilus)</i> Anthracnose ^[1] <i>(Colletotrichum acutatum)</i> <i>(Glomerella cingulata)</i>	10.0 to 17 fl oz/acre (0.130 - 0.222 lb/acre fluopyram) (0.130 - 0.222 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14- to 21-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Disease Suppression	Application Rate	Product Instructions
Eastern filbert blight <i>(Anisogramma anomala)</i>	10.0 to 17 fl oz/acre (0.130 - 0.222 lb/acre fluopyram)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.

	(0.130 - 0.222 lb/acre tebuconazole)	
--	--	--

Restrictions:

- Maximum single application rate: 17 fl oz/acre of LUNA EXPERIENCE (0.222 lb/acre fluopyram and 0.222 lb/acre tebuconazole).
- Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 3 (at 10 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 7 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.900 lbs tebuconazole per acre per year, including soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE within 35 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.
- **DO NOT** cut cover crops in treated areas for feed or allow livestock to graze treated areas.

[¹Not for use in CA [without a supplemental label.]]

WHEAT		
Disease Control	Application Rate	Product Instructions
Rusts (<i>Puccinia</i> spp.)	8.6 fl oz/acre (0.112 lb/acre fluopyram) (0.112 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings.
Restrictions: <ul style="list-style-type: none"> • Maximum single application rate: 8.6 fl oz/acre of LUNA EXPERIENCE (0.112 lb/acre fluopyram and 0.112 lb/acre tebuconazole). • Maximum annual application rate: 8.6 fl oz/acre of LUNA EXPERIENCE (0.112 lb/acre fluopyram and 0.112 lb/acre tebuconazole) per year. • Maximum number of applications per year: 1 • Apply using ground, aerial, or chemigation equipment. • Regardless of formulation or method of application, DO NOT apply more than 0.446 lbs Fluopyram or 0.112 lbs tebuconazole per acre per year, including soil and foliar uses. • DO NOT apply LUNA EXPERIENCE within 30 days of harvest. • To limit the potential for development of disease resistance to these fungicide classes, DO NOT make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group. 		

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container and keep tightly closed when not in use. Store in a cool dry place. Avoid cross-contamination with other pesticides.

Pesticide Disposal: Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of federal law. If these wastes cannot be disposed of by use according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling:

Non-Seed Treatment Products in Non-Refillable Containers

Rigid, Non-refillable containers (equal to or less than 5 gallons)

Non-refillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill.

Dilutable Non-Seed Treatment Products in Rigid Non-refillable Containers that are Too Large to Shake (i.e., with capacities greater than 5 gallons or 50 lbs)

Non-refillable container. **DO NOT** reuse or refill this container. Refer to Bottom Discharge IBC or Top Discharge IBC, Drums, Kegs information as follows.

Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Top Discharge IBC, Drums, Kegs (e.g.– Snyder 120 Next Gen, Bonar B120, Drums, Kegs)

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. To triple rinse the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

Non-Seed Treatment Products in Non-Refillable Fiber Drums with Liners

Non-refillable container. **DO NOT** reuse or refill this container. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment, then offer for recycling if available or dispose of in a sanitary landfill or by incineration. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner.

Non-Seed Treatment Products in Non-Rigid, Non-refillable Containers

Nonrefillable container. **DO NOT** reuse or refill this container. Completely empty container into application equipment. Then offer for recycling if available or dispose of in a sanitary landfill or by other procedures approved by state and local authorities."

Non-Seed Treatment Products in Refillable Containers

Refillable container. Refer to Bottom Discharge IBC or Top Discharge IBC, Drums, Kegs information as follows. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Contact your Ag retailer or Bayer CropScience for container return, disposal and recycling information.

Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Top Discharge IBC, Drums, Kegs (e.g.– Snyder 120 Next Gen, Bonar B120, Drums, Kegs)

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To triple rinse the containers before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

End users are authorized to remove tamper evident cables as required to remove the product from the container unless the container is equipped with one way valves and refilling or returning is planned. If this is the case, end users are not authorized to remove tamper evident cables, one way valves or clean container.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE SHALL NOT BE LIABLE FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

LUNA EXPERIENCE is specially formulated and sold by Bayer CropScience LP for the control of various pathogens according to the directions on this label. The purchase price of LUNA EXPERIENCE includes a prepaid license under which purchaser agrees to employ the purchased quantity of LUNA EXPERIENCE only for the above-specified uses and to provide notice of the terms and conditions of this license to any subsequent purchaser. Uses of LUNA EXPERIENCE other than those specified on this label are not licensed through the purchase of this product.

NET CONTENTS:

PRODUCED FOR



Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167
1-866-99BAYER (1-866-992-2937)

LUNA EXPERIENCE (PENDING) 07/22/2021, 08/20/2021