



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

August 19, 2020

Ona Maune
Federal Registration Manager
Bayer CropScience LP
800 N. Blvd.
St. Louis, MO 63167

Subject: Amendment to Add a Supplemental Label – Grasses Grown for Seed
Product Name: Indaziflam 200 SC Herbicide
EPA Registration Number: 264-1106
Application Date: July 8, 2020
Decision Number: 564772

Dear Ms. Maune:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval is only for a supplemental label that is an addendum to the master label. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

This supplemental labeling contains some new and/or revised uses and/or directions which may be additional to the uses and/or directions found on the label on or attached to the container, but this supplemental labeling does not by itself constitute the complete set of use directions. The complete set of use directions is set forth on the container label as combined with this supplemental labeling.

A stamped copy of your labeling is enclosed for your records. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Kable Bo Davis by phone at 703-306-0415, or via email at davis.kable@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "ES", is written over a horizontal line.

Emily Schmid
Product Manager 25
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure



Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167
1-866-99BAYER (1-866-992-2937))

ACCEPTED

08/19/2020

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 264-1106

Indaziflam 200SC Herbicide
ABN: Alion® Herbicide

EPA Reg. No: 264-1106

For Use On: Grasses Grown for Seed

*This supplemental label expires on 06/29/2023 and
must not be used or distributed after this date.*

Supplemental Label

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Read this label and the product package label before using this product. This Supplemental Label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the product label for Indaziflam 200SC Herbicide attached to the container.

APPLICATION DIRECTIONS FOR USE ON GRASS GROWN FOR SEED IN HIGH RAINFALL AREAS OF WESTERN OREGON WEST OF THE CASCADE MOUNTAINS

USE ONE of the following Indaziflam 200SC Herbicide use patterns **only once** over the duration of the grass seed stand life due to potential crop response:

CARBON-SEEDED GRASSES:

Apply Indaziflam 200SC Herbicide following planting using a carbon-seeding system to perennial ryegrass and tall fescue. Use an application rate of 1.0 oz/A. **Do not** apply Indaziflam 200SC Herbicide in any of the subsequent production years.

It is **recommended** that strict adherence to the carbon-planting guidelines be followed or crop injury and reduced stand can occur. Apply only to fields with well-prepared, firmly packed seedbeds with an adequate carbon band that will absorb the applied Indaziflam 200SC Herbicide.

It is **recommended** that the carbon band be a minimum of 1" wide or reduced stand can occur. Apply Indaziflam 200SC Herbicide following planting but before irrigation or rainfall.

ESTABLISHED GRASSES:

Apply Indaziflam 200SC Herbicide to established perennial ryegrass or tall fescue which has had at least one seed harvest and which **did not** have Indaziflam 200SC Herbicide applied in the establishment (1st year) of the crop. Indaziflam 200 SC Herbicide may also be applied at a rate of 2.0 to 2.0 fl oz/A to timothy grown for seed or hay that has been established at least one year and through a harvest for seed or hay. Apply at a rate of 1.0 to 2.0 fl oz/A. Apply in the fall prior to the onset of fall rains and emergence of winter annual grass weeds and volunteer crop. Check grass variety tolerance on a small scale before treating entire fields. Indaziflam 200SC Herbicide may be tankmixed with some registered post-emerge herbicides to control already emerged *Poa spp.* and annual ryegrass and volunteer crop. Refer to the individual product labels for specific use rates, application timings and/or precautions and restrictions. Always follow the most restrictive label.

USE RESTRICTIONS

- **Do not** apply Indaziflam 200SC Herbicide in any of the subsequent production years.
- **Do not apply** more than a total of 2.0 fl oz product/A (0.025 lb ai/A) per year when used in any grass grown for seed planting.
- **Do not apply** within two years of rotation to another crop (refer to Rotational Crop Restrictions)
- **Do not apply** Indaziflam 200SC Herbicide where there are exposed roots, open channels or depressions in the soil that would allow the product to move into the root zone or injury may occur.

APPLICATION DIRECTIONS FOR USE ON GRASS GROWN FOR SEED IN LOW RAINFALL AREAS EAST OF THE CASCADE MOUNTAINS IN OREGON AND WASHINGTON AND IDAHO

ESTABLISHED GRASSES ONLY:

Apply Indaziflam 200SC Herbicide at a rate of 2.0 to 3.0 fl oz/A to established perennial ryegrass, tall fescue, smooth brome grass, and wheatgrass seed stands which have been established at least one year and have been through a seed harvest. Apply in the fall prior to rainfall and winter annual weed germination in established grasses that are dormant but beginning to regrow. Check grass variety tolerance on a small scale before treating entire fields. Indaziflam 200 SC Herbicide may also be applied at a rate of 2.0 to 3.0 fl oz/A to timothy grown for seed or hay that has been established at least one year and through a harvest for seed or hay.

USE RESTRICTIONS

- **Do not apply** more than a total of 3.0 fl oz product/A (0.039 lb ai/A) per year when used in any grass grown for seed planting in eastern OR, eastern WA, and in Idaho.
- **Do not apply** within two years of rotation to another crop (refer to Rotational Crop Restrictions)
- **Do not apply** Indaziflam 200SC Herbicide where there are exposed roots, open channels or depressions in the soil that would allow the product to move into the root zone or injury may occur.
- Do not cut hay within 30 days of a single application.
- Do not apply Indaziflam 200SC Herbicide more than once per year.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).