264-1109

03-16-2010





U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs Registration Division (7505C) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460

264-1109

Date of Issuance:

MAR 1 6 2010

NOTICE OF PESTICIDE:

X Registration

Reregistration

(under FIFRA, as amended)

Term of Issuance:

EPA Reg. Number:

Unconditional

Name of Pesticide Product:

Poncho Votivo

Name and Address of Registrant (include ZIP Code):

Bayer CropScience 2 T.W. Alexander Drive P.O. Box 12014

Research Triangle Park, NC 27709

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

- 1. Submit and/or cite all data required for registration/registration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
- 2. Make the following label changes before you release the product for shipment:
 - a. Revise the EPA Registration Number to read, "EPA Reg. No. 264-1109".
- 3. The data requirements for storage stability (830-6317) and corrosion characteristics (830-6320) have not been satisfied, and must be submitted within eighteen months of the date of this letter.

Signature of Approving Official:

Date:

MAR 1 6 2010

Venus Eagle, Product Manager (01)

Insecticide-Rodenticide Branch, Registration Division (7505P)

Page 2 EPA Reg. No. 264-1109

4. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records. If you have any questions regarding this notice, please contact Kable Bo Davis at (703) 306-0415.

Venus Eagle Product Manager (01) Insecticide-Rodenticide Branch Registration Division (7505P)

Enclosure-Stamped Labeling

PONCHO® VOTIVO™

A systemic insecticide and biological seed treatment for use on corn and sorghum for the control of listed insect pests and listed soil plant pathogenic nematodes

ACTIVE INGREDIENTS: Clothianidin

Bacillus firmus I-1582

OTHER INGREDIENTS: 51.6% TOTAL:100.0%

Contains 4.17 pounds clothianidin per U.S. gallon

Contains 0.84 pounds Bacillus firmus per U.S. gallon contains a minimum of 2 X109 cfu/ml

EPA Reg. No. 264-RRNO

EPA Est.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577 For PRODUCT USE Information Call 1-800-331-2867

FIRST AID

IF SWALLOWED:	Call a poison control center or doctor immediately for treatment advice.		
	Have person sip a glass of water if able to swallow.		
	Do not induce vomiting unless told to do so by a poison control center or doctor.		
	Do not give anything by mouth to an unconscious person.		
IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15-20 minutes.		
	Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.		
	Call a poison control center or doctor for treatment advice.		

In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

NOTE TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically.

ACCEPTED with COMMENTS Im EPA Letter Dated:

PRECAUTIONARY STATEMENTS HAZARDS TO HUMAN AND DOMESTIC ANIMALS

MAR 16 2010 Under the Federal Insecticide.

Fungicide, and Redenticide Act,

CAUTION

Example of the positive registered under EPA Reg. No.

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical resistance category selection chart.

Workers involved with treating the seed (e.g. connecting and disconnecting hoses and transfer pumps, mixing, equipment calibration, etc.) and others exposed to the concentrate, and cleaners/repairers of seed treatment equipment must wear a long-sleeve shirt and long pants, shoes plus socks, chemical-resistant gloves (such as nitrile, butyl, neoprene, barrier laminate, polyvinyl chloride or Viton), and a dust/mist filtering respirator meeting NIOSH standards of at least N-95, R-95 or P-95.

Baggers and bag sewers must wear a long-sleeve shirt and long pants, shoes plus socks, and a dust/mist filtering respirator meeting NIOSH standards of at least N-95, R-95 or P-95.



User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as
 possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Contain any product spills or equipment leaks and dispose of wastes according to disposal instructions on this label. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

For use only in commercial seed treatment equipment. Not for use in hopper box, planter box, slurry box, or other on-farm seed treatment applications.

Read the entire label and observe all label directions and precautions before use. Do not use this product except as directed on this label or on other Bayer CropScience supplemental labeling for this product.

USE INFORMATION

PONCHO VOTiVO is a combination insecticide and biological seed treatment that, when applied to seed, protects the seed and seedling against certain early season insects and provides early season protection from plant pathogenic nematodes that attack the root system. As a result of the dual protection, there is an improvement in plant vigor, which often results in more uniform plants and greater yields. In areas of high nematode infestation additional control measures may be warranted.

COMPATIBILITY

It is essential that before using PONCHO VOTiVO seed treatment in any tank mixture the compatibility of the mixture be established. Add PONCHO VOTiVO at the labeled rate to a clean quart jar containing approximately one-half the amount of water intended for a final slurry application rate. Next, follow with all other tank mix components that will be used in the total slurry application. Add last the remaining balance of water. The total amount of volume is determined by the seed size and how much is necessary to ensure complete and uniform coverage and distribution on the seed, as well as the type of commercial seed treating application equipment that will be used.

DO NOT USE MIXTURES THAT CURDLE, PRECIPITATE, OR GEL. USE TANK MIXTURES IMMEDIATELY AFTER MIXING WITH ADEQUATE AGITATION.

)
4	1
,	J

ROTATIONAL CROP PLANT-BACK INTERVALS							
Crop Planted	Immediate Plant-back	30-Day Plant-back	4-Month Plant- back	8-Month Plant-back	12-Month Plant-back		
Field corn, popcorn	Cereal grains, Field corn, popcorn, sweet corn, rapeseed, canola, soybeans, root and tuber vegetables	Grasses, non-grass animal feeds, dry beans,		Sugar cane	Any crops without an earlier plant- back interval		
Sweet corn, sorghum	Cereal grains, Field corn, popcorn, sweet corn, rapeseed, soybeans, canola, sorghum root and tuber vegetables	Grasses, non-grass animal feeds, dried beans	Any crop without an earlier plant- back interval				

DIRECTIONS FOR SEED TREATMENT

All seed treated with this product must be colored with an EPA-approved dye or colorant that imparts an unnatural color to the seed to help prevent the inadvertent use of treated seed for food, feed or oil purposes.

FOR USE AS A SEED TREATMENT

Apply PONCHO VOTiVO as a water-based slurry either alone or with other registered seed treatment insecticides and fungicides through standard slurry or mist commercial seed treatment equipment. Treated seed must be labeled in accordance with the requirements of the Federal Seed Act. Observe the most restrictive of the labeling limitations and precautions of all products used in mixtures.

MIXING INSTRUCTIONS

When applied with seed treatment application equipment designed to treat seed utilizing a treatment slurry mixture, add approximately ½ to ¾ of the required water to the treatment slurry tank followed by the accompanying seed treatment products and PONCHO VOTiVO. The order of product addition is dependant on the accompanying seed treatment products and application equipment used. Allow each slurry component to disperse completely prior to the next addition until a uniform suspension is obtained. After all slurry components have been added and dispersed completely add the remaining ½ to ⅓ of required water for proper slurry volume and mix until uniform. Do not store mixed slurries for greater than 72 hours. Maintain adequate agitation until treatment slurry is applied. For additional instructions for use, slurry component mixing order, and other application suggestions contact your Bayer CropScience representative.



APPLICATION RATES

May be used in combination with other registered Bayer CropScience seed treatment fungicides and insecticides. Pre-test for compatibility with other seed treatment products.

CROP	INSECT PESTS	NEMATODE PESTS	RATE PONCHO VOTIVO
Corn (including field, sweet and pop)	Chinch bug Corn flea beetle Corn leaf aphid Cutworm, black Grape colaspis Seedcorn maggot Southern corn leaf beetle Southern corn rootworm Southern green stinkbug Storage pests¹ (including Indian meal moth, Lesser grain borer, Red flour beetle, Rice weevil, Rusty grain beetle) Sugarcane beetle White grub (including European, Chafer larvae, May/June beetle larvae, Japanese beetle larvae) Thrips Wireworm	Soil Nematodes including Dagger Nematode Lance Nematode Needle Nematode Pin Nematode Ring Nematode Root Knot Nematode Root Lesion Nematode Spiral Nematode Sting Nematode Stubby Root Nematode Stunt Nematode	To deliver 0.25 mg ai/seed of clothianidin: Use PONCHO VOTiVO at 1.35 fl oz / 80,000 seeds. To deliver 0.50 mg ai/seed of clothianidin: Use PONCHO VOTiVO at 2.7 fl oz / 80,000 seeds Do not apply more than 0.5 mg ai/seed of clothianidin
Sorghum	Cutworm, black Aphid (including Greenbug, Yellow sugarcane aphid, Corn leaf aphid, and English grain aphid) Chinch bugs White grub Wireworms	Ring Nematode Root Knot Nematode Root Lesion Nematode Sting Nematode Stubby Root Nematode	6.13 fl oz/cwt (100 pounds of seed)

¹ It is recommended that seed with existing populations of storage pests be fumigated prior to treating and bagging seed.

SEED BAG LABEL REQUIREMENTS

The Federal Seed Act requires that bags containing treated seeds shall be labeled with the following statements:

- This seed has been treated with Bacillus firmus and clothianidin.
- Do not use treated seed for food, feed or oil purposes.
- Store away from feeds and other foodstuffs.
- Wear long-sleeved shirt, long pants and waterproof gloves when handling treated seed.
- Dispose of all excess treated seed. Left over treated seed may be doublesown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- For Field Corn, Popcorn: Immediate Plant-back: Cereal grains, Field corn, popcorn, sweet corn, rapeseed, canola, soybeans, root and tuber, 30-Day Plant-back: Grasses, non-grass animal feeds, dry beans, 8-Month Plant-back: Sugar cane, 12-Month Plant-back: Any crops without an earlier plant-back interval.
- For Sweet corn, Sorghum: Immediate Plant-back: Cereal grains, Field corn, popcorn, sweet corn, rapeseed, soybeans, canola, sorghum root and tuber vegetables, 30-Day Plant-back: Grasses, non-grass animal feeds, dried beans, 4-Month Plant-back: Any crop without an earlier plant-back interval

² Follow up sprays of a locally registered insecticide may be needed under heavy pest pressure. Consult your local Extension office for recommendations.



STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE

Store in cool, dry conditions.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Containers with capacities equal to or less than 5 gallons

Non-refillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned stay out of smoke.

Containers that are too large to shake (i.e., with capacities greater than 5 gallons)

Non-refillable containers - Do not reuse or refill this container. After emptying product from container, either return container to Bayer CropScience per instructions from Bayer CropScience Customer Service Center (1-800-527-4781), or rinse and either recycle or dispose of the container as follows:

Bottom Discharge IBC (e.g. - Schuetz Caged IBC or Snyder Square Stackable

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inch on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Top Discharge IBC, Drums, Kegs (e.g.- Snyder 120 Next Gen, Bonar B120, Drums, and Kegs).

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. To triple rinse the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill



Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Bayer CropScience, no claims are made to guarantee germination of carry-over seed.

CONDITIONS: The directions for use of this product are believed to be adequate and the user must follow directions. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.


LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

NET CONTENTS:

Produced for



Bayer CropScience LP P.O. Box 12014, 2 T.W. Alexander Drive Research Triangle Park, North Carolina 27709 1-866-99BAYER (1-866-992-2937)

