

264-1119

5/11/2012

4/11



U S ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
1200 Pennsylvania Ave N W
Washington D C 20460

EPA Reg Number

264-1119

Date of Issuance

MAY 11 2012

NOTICE OF PESTICIDE

Registration
 Reregistration
(under FIFRA as amended)

Term of Issuance

Unconditional

Name of Pesticide Product

EverGol Prime

Name and Address of Registrant (include ZIP Code)

Bayer CropScience LP
2 T W Alexander Drive
P O Box 12014
Research Triangle Park, NC 27709

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant the above named pesticide is hereby registered under the Federal Insecticide Fungicide and Rodenticide Act

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment the Administrator on his motion may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you

1. Submit and/or cite all data required for registration/registration review of your product when the Agency requires all registrant of similar products to submit such data
2. Make the following label changes before you release the product for shipment
 - a. Revise the EPA Reg No so it reads "EPA Reg No 264-1119"
 - b. In the "On Farm Seed Piece Application" table under "Directions" revise the first sentence to read as follows "Apply specified dosage as a diluted spray slurry using equipment that ensures uniform coverage of each seed piece"
 - c. For clarity and consistency in the "Commercial Seed Treatment" table in the "Application Rates" column revise each "Maximum number of application for each crop listed is 3 per year" to read "Maximum number of applications per year is 3"
 - d. For clarity and consistency in the "Commercial Seed Treatment" table in the "Application Rates" column insert the information to deliver "xx mg ai/seed" for alfalfa, cereal grains, corn and sorghum, sunflower and safflower, and rice. For oilseeds add the information to deliver "xx g ai/100 lbs"
 - e. In the "Commercial Seed Treatment" table in the "Directions" column revise the first two sentences of the dye statement to read as follows "All seed must be adequately dyed with a suitable colorant to prevent its accidental use as food for man or feed for animals"

- f For clarity and consistency on page 6 of the label in the "Directions" column, remove the row divider between rice and the rest of the crop use sites – the Directions apply to all of the crops
 - g On page 6 of the label, for clarity and consistency based on the BEAD memo from 3/24/11 revise footnote #2 to read "Based on 4,500 cottonseed per pound "
 - h On page 6 of the label, for clarity and consistency in footnote #3 adjust seed per pound to reflect each corn variety (e g , popcorn, sweet, and field) based on the BEAD memo of 3/24/11
 - 1 Add the batch code information Please be reminded that the 40 CFR part 156 140(a)(4) requires a batch code, lot number, or other code identifying the batch of the pesticide product distributed and sold be placed on non-refillable containers The code may appear either on the label or durably marked on the container itself
- 3 For future submissions, make the following changes to your Confidential Statement of Formula (EPA Form 8570-4)
- a Box 4 Revise the EPA Reg No so it reads "EPA Reg No 264-1119 "
- 4 Submit 2 copies of your final printed label before release for shipment

<p>Signature of Approving Official</p>  <p>Mark Corbin, Acting Branch Chief Insecticide-Rodenticide Branch, Registration Division (7505P)</p>	<p>Date</p> <p>MAY 11 2012</p>
---	--------------------------------

EPA Form 8570 6

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e) Your release for shipment of the product constitutes acceptance of these conditions A stamped copy of the label is enclosed for your records If you have any questions, please contact Marianne Lewis at 703-308-8043 or lewis marianne@epa gov

Mark Corbin
Acting Branch Chief,
Insecticide-Rodenticide Branch
Registration Division (7505P)

Enclosure

3/11

EverGol Prime

For use as a fungicide for protection against listed soilborne, seedborne, and post-emergence diseases for listed crop plants

ACTIVE INGREDIENT
Penflufen
OTHER INGREDIENTS

22.7%
77.3%
TOTAL 100.0%

Contains 2 lb penflufen/gallon (241 g penflufen/liter)

EPA Reg No ~~264-XXX~~ 1119

EPA Est

KEEP OUT OF REACH OF CHILDREN CAUTION

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1 800 334 7577
For PRODUCT USE Information Call 1-866 99BAYER (1 866-992-2937)

FIRST AID

IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to by a poison control center or doctor • Do not give anything by mouth to an unconscious person
IF ON SKIN	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15 20 minutes • Call a poison control center or doctor for treatment advice
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing call 911 or ambulance then give artificial respiration preferably mouth to mouth if possible • Call a poison control center or doctor for further treatment advice
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 20 minutes • Remove contact lenses if present after the first 5 minutes then continue rinsing • Call a poison control center or doctor for treatment advice

In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No 1 800 334 7577
Have a product container or label with you when calling a poison control center or doctor or going for treatment

ACCEPTED
5/11/12
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
Registered under
EPA Reg No 264-1119

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed Harmful if absorbed through skin Harmful if inhaled Causes moderate eye irritation Avoid contact with skin eyes or clothing Avoid breathing spray mist

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are barrier laminate butyl rubber nitrile rubber neoprene rubber polyvinyl chloride or viton If you want more options follow the instructions for category C on an EPA chemical resistance category selection chart

All mixers, loaders, applicators and other handlers must wear

long sleeved shirt and long pants

socks plus shoes

chemical resistant gloves except when bagging or sewing bags of treated seeds

Follow manufacturer's instructions for cleaning/maintaining PPE If no such instructions for washables exist use detergent and hot water Keep and wash PPE separately from other laundry

USER SAFETY RECOMMENDATIONS

Wash hands before eating drinking chewing gum using tobacco or using the toilet

Remove clothing/PPE immediately if pesticide gets inside Then wash thoroughly and put on clean clothing

Remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark Runoff may be hazardous to aquatic organisms in water adjacent to treated areas Do not contaminate water when disposing of equipment washwater or rinsate Treated seed exposed on soil surface may be hazardous to wildlife Cover or collect seed spilled during loading

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

For use only in commercial seed treatment equipment Not for use in hopper box planter box slurry box or other on farm seed treatment applications except for potato seed pieces

The Agricultural Use Requirements statements below are specific to the potato seed piece and in furrow use

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170 (This Standard contains requirements for the protection of agricultural workers on farms forests nurseries and greenhouses and handlers of agricultural pesticides It contains requirements for training decontamination notification and emergency assistance It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours Exception if the product is soil injected or soil incorporated the Worker Protection Standard under certain circumstances allows workers to enter the treated area if there will be no contact with anything that has been treated PPE required for early reentry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants soil or water is coveralls chemical resistant gloves made of any waterproof material and shoes plus socks

ON FARM SEED PIECE APPLICATION

CROP	DISEASE	APPLICATION RATE	DIRECTIONS
<p>Potato</p>	<p>Black scurf stem & stolon canker caused by seedborne and soilborne (suppression) <i>Rhizoctonia solani</i></p> <p>Silver scurf caused by seedborne <i>Helminthosporium solani</i> (suppression)</p>	<p>Apply 0.13 fl oz per 100 lbs of seed pieces (2 g ai per 100 kg seed)</p> <p>Maximum number of application per year is 3</p>	<p>For disease control good coverage of the seed piece is required</p> <p>Apply specified dosage as a diluted spray using equipment that ensures uniform coverage of each seed piece. Do not apply more than 2.5 fl oz of slurry / 100 lbs of seed pieces. Apply only in areas with adequate ventilation or in areas that are equipped to remove spray mist or dust</p> <p>As part of the seed cutting and treating process application of an inert absorbent ingredient is recommended to improve suberization. For added <i>Fusarium</i> protection apply a Mancozeb containing product specifically designed for application to potatoes in place of the inert absorbent</p> <p>Do not use treated seed pieces for food feed or fodder</p> <p>For added protection against soilborne black scurf apply penflufen as an in furrow spray in addition to the seed piece application</p> <p>Do not exceed more than 0.143 lbs of active ingredient/A (65 g ai/A) per crop season for seed piece treatment application. This seed piece has been treated with xx mg ai/seed (2 g ai/100 Kg seed)</p>

IN-FURROW APPLICATION TOGETHER WITH SEED PIECE TREATMENT

CROP	DISEASE	APPLICATION RATE	DIRECTIONS
<p>Potato and other tuberous and corm vegetables (Crop Subgroup 1C)</p> <p>Arracacha Arrowroot Artichoke (Chinese and Jerusalem) Canna (edible) Cassava (bitter and sweet) Chayote (root) Chufa Dasheen (taro) Ginger Leren Potato Sweet Potato Tanier (cocoyam) Turmeric Yam Bean Yam (true)</p>	<p>Black scurf including stem & stolon canker caused by soilborne <i>Rhizoctonia solani</i></p>	<p>In furrow treatment in combination with seed piece treatment</p> <p>Apply a minimum of 0.4 fl oz up to a maximum of 0.57 fl oz per 1,000 row feet (4.55 ml per 100 row meters)</p> <p>Lower rates provide reduced levels of protection</p> <p>Maximum number of application per year is 3</p>	<p>Apply as an in furrow spray or drench at planting</p> <p>Mount spray nozzles to direct the spray as a 6 to 8 inch band onto the seed pieces in the furrow just before the seed is covered or apply as a soil drench in transplant water</p> <p>Do not exceed more than 0.143 lbs of active ingredient/A (65 g ai/A) per crop season through combined seed piece treatment and in furrow application. This seed piece has been treated with xx-mg ai/seed (2 g ai/100 Kg seed)</p>

IN-FURROW APPLICATION ONLY

CROP	DISEASE	APPLICATION RATE	DIRECTIONS
<p>Potato and other tuberous and corm vegetables (Crop Subgroup 1C)</p> <p>Arracacha Arrowroot Artichoke (Chinese and Jerusalem) Canna (edible) Cassava (bitter and sweet) Chayote (root) Chufa Dasheen (taro) Ginger Leren Potato Sweet Potato Tanier (cocoyam) Turmeric Yam Bean Yam (true)</p>	<p>Black scurf including stem & stolon canker caused by soilborne <i>Rhizoctonia solani</i></p>	<p>In furrow treatment without seed piece treatment</p> <p>Apply a minimum of 0.5 fl oz per 1 000 row feet (5 ml per 100 row meters) up to a maximum of 0.57 fl oz per 1 000 row feet (4.5 ml per 100 row meters) Applying without a seed piece treatment in potatoes will result in reduced protection against seed borne diseases</p> <p>Maximum number of application per year is 3</p>	<p>Apply as an in furrow spray or drench at planting</p> <p>Mount spray nozzles to direct the spray as a 6 to 8 inch band onto the seed pieces in the furrow just before the seed is covered or apply as a soil drench in transplant water</p> <p>Do not exceed more than 0.143 lbs of active ingredient/A (65 g ai/A) per crop season for in furrow application</p>

COMMERCIAL SEED TREATMENT

CROP	DISEASE	APPLICATION RATE PER 100 LBS SEED	DIRECTIONS
<p>Alfalfa</p>	<p>Seed rot and damping off caused by <i>Rhizoctonia solani</i></p>	<p>Apply 0.96 fl oz (15 g ai per 100 kg seed)</p> <p>Maximum number of application for each crop listed is 3 per year</p>	<p>Apply using commercial slurry or mist type seed treatment equipment Uniform application of seed is necessary to ensure seed safety and best disease protection Dilute product with sufficient water to ensure complete seed coverage Consult a seed treatment specialist regarding specified slurry rates for the crop to be treated with EverGol Prime Use high quality properly cleaned seed</p>
<p>Beans and Peas including Soybean and Legume Foliage (Crop Group 6 and 7)</p> <p>Soybean Adzuki Bean Asparagus Bean Broad Bean (Succulent or Dry) Catjang Bean Chinese Longbean Field Bean Guar Bean Jackbean Kidney Bean Lablab Bean Lima Bean (Succulent or Dry) Moth Bean (Succulent or Dry) Mung Bean Navy Bean Pinto Bean Rice Bean Runner Bean Snap Bean Sword Bean Tepary Bean Urd Bean Wax Bean Yardlong Bean Blackeyed Pea (Succulent or Dry) Chickpea Cowpea (Succulent or Dry) Crowder Pea Dwarf Pea Edible Pod Pea English Pea Field Pea Garden Pea Green Pea Pigeon Pea (Succulent or Dry) Snow Pea Southern Pea (Succulent or Dry) Sugar Snap Pea Grain Lupin White Lupin Sweet Lupin White Sweet Lupin Lentil</p>	<p>Seed rot and damping off caused by <i>Rhizoctonia solani</i></p>	<p>Apply 0.32 fl oz (5 g ai per 100 kg seed)</p> <p>To deliver 0.0081 mg ai per seed apply 0.16 fl oz of the product per 140 000 seeds¹</p> <p>Maximum number of application for each crop listed is 3 per year</p>	<p>All seed must be adequately dyed with a colorant The purchaser of this product is responsible for ensuring that all-seed-treated with this product are adequately dyed with a suitable colorant to prevent its accidental use as food for man or feed for animals Refer to 21CFR part 2.25 Any colorant added to treated seed must be cleared for use under 40CFR Part 153.155</p>
<p>Oil Seeds (Crop Group 20)</p> <p>Borage Calendula Canola</p>	<p>Seed rot and damping off caused by <i>Rhizoctonia solani</i></p>	<p>Apply 0.96 fl oz per 100 lbs seed (15 g ai per 100 kg</p>	

7/11

CROP	DISEASE	APPLICATION RATE PER 100 LBS SEED	DIRECTIONS
Castor Oil Plant Chinese Tallowtree Crambe Cuphea Echim Euphorbia Evening Primrose Flax Seed Gold of Pleasure Hare s Ear Mustard Jojoba Lesquerella Lunaria Meadowfoam Milkweed Mustard Seed Niger Seed Oil Radish Poppy Seed Rapeseed Rose Hip Sesame Stokes Aster Sweet Rocket Tallowwood Tea Oil Plant Veronia and cultivars varieties and or hybrids of these		seed) Maximum number of application for each crop listed is 3 per year	
Cotton	Seed rot and damping off caused by <i>Rhizoctonia solani</i>	Apply 0.64 fl oz per 100 lbs seed (10 g ai per 100 kg seed) To deliver 0.009 mg ai per seed apply 0.128 fl oz of the product per 100,000 seeds ² Maximum number of application for each crop listed is 3 per year	
Cereal grains Crop Group 15 and 16 (except corn sorghum and rice) Wheat barley oats buckwheat millet (pearl and proso) rye teosinte triticale wild rice	Common bunt Covered smut False loose smut Flag smut Leaf strip Loose smut Stinking smut True loose smut Seed rot pre emergence damping off and seedling blight caused by soilborne <i>Rhizoctonia solani</i>	Apply 0.32 fl oz per 100 lbs seed (5 g ai per 100 kg seed) Maximum number of application for each crop listed is 3 per year	
Corn (field, sweet popcorn) Sorghum	Seed rot and damping off caused by <i>Rhizoctonia solani</i>	Apply 0.64 fl oz per 100 lbs seed (10 g ai per 100 kg seed) To deliver 0.026 mg ai per seed apply 0.29 fl oz of the product per 80,000 seeds ³ Maximum number of application for each crop listed is 3 per year	
Sunflower and Safflower	Seed rot and damping off caused by <i>Rhizoctonia</i>	Apply 0.96 fl oz per 100 lbs seed	

CROP	DISEASE	APPLICATION RATE PER 100 LBS SEED	DIRECTIONS
	<i>solani</i>	(15 g ai per 100 kg seed) Maximum number of application for each crop listed is 3 per year	
Rice	Seed rot and damping off caused by <i>Rhizoctonia solani</i> Seedborne false smut (suppression) Seedborne kernel smut (suppression)	Apply 0.32 fl oz per 100 lbs seed (5 g ai per 100 kg seed) Maximum number of application for each crop listed is 3 per year	

¹ Based on 2 800 soybean seeds per pound

² Based on 4 800 cottonseed per pound

³ Based on 1 750 corn seed per pound

SEED BAG TAG

The Federal Seed Act requires that the container of seed treated with EverGol Prime must be labeled with the following statements

- This seed has been treated with EverGol Prime which contains penflufen
- Do not use treated seed for food feed or oil production Excess treated seed may be used for ethanol production only if (1) by products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by products that are used in agronomic practice
- User is responsible for ensuring that the seed bag meets all the requirements under the Federal Seed Act

In addition the US Environmental Protection Agency requires the following statements on the container of seed treated with EverGol Prime

- Store away from feeds and foodstuffs
- Wear long sleeved shirt long pants shoes socks and chemical resistant gloves when handling treated seed
- After the seeds have been planted do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours Exception Once the seeds are planted in soil or other planting media the Worker Protection Standard allows workers to enter the treated area without restriction if there will be no worker contact with the treated seeds in the soil or planting media
- Dispose of all excess treated seed Left over treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements Do not contaminate water bodies when disposing of planting equipment washwaters
- Dispose of seed packaging in accordance with local requirements
- Use planting equipment that will plant treated seed into the soil to a minimum depth of 0.5 inch
- Crops for which there are registered uses of penflufen which includes alfalfa potatoes (Crop Subgroup 1C) legume vegetables (Crop Group 6/7) cereal grains (Crop Group 15/16) and oil seeds (Crop Group 20) may be replanted at any time All other crops may be planted no less than 30 days after planting treated seeds
- This pesticide is toxic to fish Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark
- Treated seed exposed on soil surface may be hazardous to wildlife Cover or collect seeds spilled during loading
- Runoff may be hazardous to aquatic organisms in water adjacent to treated areas
- Do not plant treated rice seed directly into flooded field Do not soak treated rice seed
- For alfalfa Do not exceed 0.00225 lb active ingredient/A (1.02 g ai/A) per crop season The maximum number of applications per year is 3 This seed has been treated with xx mg ai/seed (15 g ai/100 Kg seed)
- For beans and peas (excluding soybeans) Do not exceed 0.0384 lb active ingredient/A (17.5 g ai/A) per crop season The maximum number of applications per year is 3 This seed has been treated with xx mg ai/seed (5 g ai/100 Kg seed)
- For soybeans Do not exceed 0.0384 lb active ingredient/A (17.5 g ai/A) per crop season The maximum number of applications per year is 3 This seed has been treated with xx mg ai/seed (5 g ai/100 Kg seed)
- For oil seed crops (based on canola seeding rate) Do not exceed 0.00123 lb active ingredient/A (0.6 g ai/A) per crop season The maximum number of applications per year is 3 This seed has been treated with xx mg ai/seed (15 g ai/100 Kg seed)
- For cotton Do not exceed 0.0019 lb active ingredient/A (0.86 g ai/A) per crop season The maximum number of applications per

9/11

- year is 3 This seed has been treated with xx mg ai/seed (10 g ai/100 Kg seed)
- For cereal grains except corn sorghum and rice (based on wheat seeding rate) Do not exceed 0 0078 lb active ingredient/A (3 5 g ai/A) per crop season The maximum number of applications per year is 3 This seed has been treated with xx mg ai/seed (5 g ai/100 Kg seed)
- For corn Do not exceed 0 0033 lb active ingredient/A (1 5 g ai/A) per crop season The maximum number of applications per year is 3 This seed has been treated with xx mg ai/seed (10 g ai/100 Kg seed)
- For sunflower and safflower Do not exceed 0 00123 lb active ingredient/A (0 6 g ai/A) per crop season The maximum number of applications per year is 3 This seed has been treated with xx mg ai/seed (15 g ai/100 Kg seed)
- For rice (dry seeded) Do not exceed 0 006 lb active ingredient/A (2 7 g ai/A) per crop season The maximum number of applications per year is 3 This seed has been treated with xx mg ai/seed (5 g ai/100 Kg seed)

INFORMATION

For use as a fungicide for protection against listed soilborne seedborne and post emergence diseases of listed crop plants

CROP ROTATION

Listed seed treatment crops and potatoes may be replanted at any time All other crops may be planted after 30 days

STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage disposal or cleaning of equipment

PESTICIDE STORAGE

Store in original container away from feed and food Store in cool dry area Do not store in direct sunlight Do not allow prolonged storage in temperatures that exceed 105°F (40°C) or in temperatures that fall below 14°F (10°C)

PESTICIDE DISPOSAL

To avoid wastes use all material in this container by application according to label directions If wastes cannot be avoided offer remaining product to a waste facility or pesticide disposal program (often such programs are run by state or local governments or by industry)

CONTAINER DISPOSAL

Rigid, Non refillable containers (equal to or less than 5 gallons)

Non refillable container Do not reuse or refill this container Offer for recycling if available Triple rinse or pressure rinse container (or equivalent) promptly after emptying Triple rinse as follows Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip Fill the container 1/4 full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times

Pressure rinse as follows Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds Drain for 10 seconds after the flow begins to drip

Once container is rinsed offer for recycling if available or puncture and dispose of in a sanitary landfill

Rigid Non refillable containers (greater than 5 gallons or 50 lbs)

Non refillable Containers

Non refillable containers Do not reuse or refill this container Refer to Bottom Discharge IBC or Top Discharge IBC Drums Kegs information as follows

Bottom Discharge IBC (e g – Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container To pressure rinse the container before final disposal empty the remaining contents from the IBC into application equipment or mix tank Raise the bottom of the IBC by 1 5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal Completely remove the top lid of the IBC Use water pressurized to at least 40 PSI to rinse all interior portions Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing Continue pressure rinsing for 2 minutes or until rinsate becomes clear Replace the lid and close bottom valve

Top Discharge IBC Drums Kegs (e g – Snyder 120 Next Gen Bonar B120 Drums Kegs)

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container To triple rinse the container before final disposal empty the remaining contents from this container into application equipment or mix tank Fill the container at least 10 percent full with water Agitate vigorously or recirculate water with the pump for 2 minutes Rinse all interior surfaces Pour or pump rinsate into application equipment or rinsate collection system Repeat this procedure two more times

Once container is rinsed offer for recycling if available or puncture and dispose of in a sanitary landfill

Refillable Containers

Refillable container – Refer to Bottom Discharge IBC or Top Discharge IBC Drums Kegs information as follows Refill this container with pesticide only Do not reuse this container for any other purpose Contact your Ag retailer or Bayer CropScience for container return disposal and recycling information

Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the refiller To pressure rinse the container before final disposal empty the remaining contents from the IBC into application equipment or mix tank Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal Completely remove the top lid of the IBC Use water pressurized to at least 40 PSI to rinse all interior portions Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing Continue pressure rinsing for 2 minutes or until rinsate becomes clear Replace the lid and close bottom valve

Top Discharge IBC Drums Kegs (e.g. – Snyder 120 Next Gen Bonar B120 Drums Kegs)

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the refiller To triple rinse the containers before final disposal empty the remaining contents from this container into application equipment or mix tank Fill the container at least 10 percent full with water Agitate vigorously or recirculate water with the pump for 2 minutes Rinse all interior surfaces Pour or pump rinsate into application equipment or rinsate collection system Repeat this procedure two more times

Once container is rinsed offer for recycling if available or puncture and dispose of in a sanitary landfill

End users are authorized to remove tamper evident cables as required to remove the product from the container unless the container is equipped with one way valves and refilling or returning is planned If this is the case end users are not authorized to remove tamper evident cables one way valves or clean container

IMPORTANT READ BEFORE USE

Read the entire Directions for Use Conditions Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, disclaimer of Warranties and Limitations of Liability:

Treatment of highly mechanically damaged seed or seed of known low vigor and poor quality may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Bayer CropScience, no claims are made to guarantee germination of carry over seed.

CONDITIONS The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

NET CONTENTS

PRODUCED FOR



Bayer CropScience

Bayer CropScience LP
P O Box 12014, 2 T W Alexander Drive
Research Triangle Park, North Carolina 27709
1 866-99BAYER (1-866 992-2937)