

264-1119

3/19/2014

1/11

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460



OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

MAR 19 2014

Jamin Huang
Bayer CropScience LP
2 T.W. Alexander Drive
P.O. Box 12014
Research Triangle Park, NC 27709

Subject: Amended label to add rate ranges while maintaining approved highest rate
EverGol Prime
EPA Reg. No. 264-1119
Submissions dated October 16, 2013

Dear Mr Huang:

The proposed labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable provided that you submit two copies of your final printed label before you release the product for shipment. Products shipped after 18 months from the date of this amendment or the next printing of the label whichever occurs first, must bear the new revised label. Your release for shipment of the product bearing the amended label constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e).

A stamped copy is enclosed for your records. If you have any questions, please contact Marianne Lewis at (703) 308-8043 or lewis.marianne@epa.gov.

Regards,

A handwritten signature in cursive script, appearing to read "Venus Eagle".

Venus Eagle, PM 01
Insecticide-Rodenticide Branch
Registration Division (7505P)

Enclosure

2/11

ACCEPTED

3/19/14

Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for the
pesticide registered under:

EverGol Prime

EPA Reg. No: 264-1119

For use as a fungicide for protection against listed soilborne, seedborne, and post-emergence diseases for listed crop plants.

ACTIVE INGREDIENT:

Penflufen 22.7%

OTHER INGREDIENTS: 77.3%

TOTAL: 100.0%

Contains 2 lb penflufen/gallon (241 g penflufen/liter)

EPA Reg. No. 264-1119

EPA Est.

KEEP OUT OF REACH OF CHILDREN

CAUTION

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
<p>In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride, or viton. If you want more options, follow the instructions for category C on an EPA chemical resistance category selection chart.

All mixers, loaders, applicators and other handlers must wear:

- long sleeved shirt and long pants,
- socks plus shoes,
- chemical resistant gloves, except when bagging or sewing bags of treated seeds,

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate. Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect seed spilled during loading.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use only in commercial seed treatment equipment. Not for use in hopper box, planter box, slurry box, or other on-farm seed treatment applications except for potato seed pieces.

The Agricultural Use Requirements statements below are specific to the potato seed piece and in-furrow use.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: if the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early reentry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is coveralls, chemical resistant gloves made of any waterproof material, and shoes plus socks.

ON FARM SEED PIECE APPLICATION

CROP	DISEASE	APPLICATION RATE	DIRECTIONS
Potato	<p>Black scurf, stem & stolon canker, caused by seedborne and soilborne (suppression) <i>Rhizoctonia solani</i></p> <p>Silver scurf caused by seedborne <i>Helminthosporium solani</i> (suppression)</p>	<p>Apply 0.13 fl oz per 100 lbs of seed pieces (2 g ai per 100 kg seed)</p> <p>Maximum number of application per year is 3.</p>	<p>For disease control, good coverage of the seed piece is required.</p> <p>Apply specified dosage as a diluted spray slurry using equipment that ensures uniform coverage of each seed piece. Do not apply more than 2.5 fl oz of slurry / 100 lbs of seed pieces. Apply only in areas with adequate ventilation or in areas that are equipped to remove spray mist or dust.</p> <p>As part of the seed cutting and treating process, application of an inert absorbent ingredient is recommended to improve suberization. For added <i>Fusarium</i> protection apply a Mancozeb containing product specifically designed for application to potatoes in place of the inert absorbent.</p> <p>Do not use treated seed pieces for food, feed, or fodder.</p> <p>For added protection against soilborne black scurf, apply penflufen as an in-furrow spray in addition to the seed piece application.</p> <p>Do not exceed more than 0.143 lbs of active ingredient/A (65 g ai/A) per crop season for seed-piece treatment application.</p>

IN-FURROW APPLICATION TOGETHER WITH SEED PIECE TREATMENT

CROP	DISEASE	APPLICATION RATE	DIRECTIONS
<p>Potato and other tuberous and corn vegetables (Crop Subgroup 1C)</p> <p>Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Canna (edible), Cassava (bitter and sweet), Chayote (root), Chufa, Dasheen (taro), Ginger, Leren, Potato, Sweet Potato, Tanier (cocoyam), Turmeric, Yam Bean, Yam (true)</p>	<p>Black scurf, including stem & stolon canker caused by soilborne <i>Rhizoctonia solani</i></p>	<p>In-furrow treatment in combination with seed piece treatment:</p> <p>Apply a minimum of 0.4 fl oz up to a maximum of 0.57 fl oz per 1,000 row feet (4-5.5 ml per 100 row meters). Lower rates provide reduced levels of protection.</p> <p>Maximum number of application per year is 3.</p>	<p>Apply as an in-furrow spray or drench at planting.</p> <p>Mount spray nozzles to direct the spray as a 6 to 8 inch band onto the seed pieces in the furrow just before the seed is covered or apply as a soil drench in transplant water.</p> <p>Do not exceed more than 0.143 lbs of active ingredient/A (65 g ai/A) per crop season through combined seed-piece treatment and in-furrow application.</p>

8/11

IN-FURROW APPLICATION ONLY

CROP	DISEASE	APPLICATION RATE	DIRECTIONS
Potato and other tuberous and corm vegetables (Crop Subgroup 1C) Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Canna (edible), Cassava (bitter and sweet), Chayote (root), Chufa, Dasheen (taro), Ginger, Leren, Potato, Sweet Potato, Tanier (cocoyam), Turmeric, Yam Bean, Yam (true)	Black scurf, including stem & stolon canker caused by soilborne <i>Rhizoctonia solani</i>	In-furrow treatment without seed piece treatment: Apply a minimum of 0.5 fl oz per 1,000 row feet (5 ml per 100 row meters) up to a maximum of 0.57 fl oz per 1,000 row feet (4-5.5 ml per 100 row meters). Applying without a seed piece treatment in potatoes will result in reduced protection against seed borne diseases. Maximum number of application per year is 3.	Apply as an in-furrow spray or drench at planting. Mount spray nozzles to direct the spray as a 6 to 8 inch band onto the seed pieces in the furrow just before the seed is covered or apply as a soil drench in transplant water. Do not exceed more than 0.143 lbs of active ingredient/A (65 g ai/A) per crop season for in-furrow application.

COMMERCIAL SEED TREATMENT

CROP	DISEASE	APPLICATION RATE PER 100 LBS SEED	DIRECTIONS
Alfalfa	Seed rot and damping-off caused by <i>Rhizoctonia solani</i>	Apply 0.64 - 0.96 fl oz (10 - 15 g ai per 100 kg seed) To deliver 0.0002 - 0.0003 mg ai per seed, apply 0.0027 - 0.004 fl oz of the product per 100,000 seeds. (Based on an average of 213,000 seeds per lb) Maximum number of application per year is 3.	Apply using commercial slurry or mist-type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Dilute product with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding specified slurry rates for the crop to be treated with EverGol Prime. Use high quality, properly cleaned seed.
Beans and Peas including Soybean and Legume Foliage (Crop Group 6 and 7) Soybean, Adzuki Bean, Asparagus Bean, Broad Bean (Succulent or Dry), Catjang Bean, Chinese Longbean, Field Bean, Guar Bean, Jackbean, Kidney Bean, Lablab Bean, Lima Bean (Succulent or Dry), Moth Bean, (Succulent or Dry), Mung Bean, Navy Bean, Pinto Bean, Rice Bean, Runner Bean, Snap Bean, Sword Bean, Tepary Bean, Urd Bean, Wax Bean, Yardlong Bean, Blackeyed Pea (Succulent or Dry), Chickpea, Cowpea (Succulent or Dry), Crowder Pea, Dwarf Pea, Edible-Pod Pea, English Pea, Field Pea, Garden Pea, Green Pea, Pigeon Pea (Succulent or Dry), Snow Pea, Southern Pea (Succulent or Dry), Sugar Snap	Seed rot and damping-off caused by <i>Rhizoctonia solani</i>	Apply 0.16 - 0.32 fl oz (2.5 - 5 g ai per 100 kg seed) Soybeans: To deliver 0.00405 - 0.0081 mg ai per seed, apply 0.08 - 0.16 fl oz of the product per 140,000 seeds. (Based on an average of 2,800 seeds per lb) Beans: To deliver 0.0071 - 0.0142 mg ai per seed, apply 0.1 - 0.2 fl oz of the product per 100,000 seeds. (Based on an average of 1,600 seeds per lb)	All seed must be adequately dyed with a suitable colorant to prevent its accidental use as food for man or feed for animals. Refer to 21CFR, part 2.25. Any colorant added to treated seed must be cleared for use under 40CFR, part 153.155.

6/11

CROP	DISEASE	APPLICATION RATE PER 100 LBS SEED	DIRECTIONS
Pea, Grain Lupin, White Lupin, Sweet Lupin, White Sweet Lupin, Lentil		<p>Peas: To deliver 0.00355 - 0.0071 mg ai per seed, apply 0.05 - 0.1 fl oz of the product per 100,000 seeds. (Based on an average of 3,181 seeds per lb)</p> <p>Maximum number of application per year is 3.</p>	
Oil Seeds (Crop Group 20) Borage, Calendula, Canola, Castor Oil Plant, Chinese Tallowtree, Crambe, Cuphea, Echium, Euphorbia, Evening Primrose, Flax Seed, Gold of Pleasure, Hare's Ear Mustard, Jojoba, Lesquerella, Lunaria, Meadowfoam, Milkweed, Mustard Seed, Niger Seed, Oil Radish, Poppy Seed, Rapeseed, Rose Hip, Sesame, Stokes Aster, Sweet Rocket, Tallowwood, Tea Oil Plant, Veronia and cultivars, varieties, and or hybrids of these.	Seed rot and damping-off caused by <i>Rhizoctonia solani</i>	<p>Apply 0.96 fl oz per 100 lbs seed</p> <p>(15 g ai per 100 kg seed or 6.8 g per 100 lbs)</p> <p>Maximum number of application per year is 3.</p>	
Cotton	Seed rot and damping-off caused by <i>Rhizoctonia solani</i>	<p>Apply 0.32 - 0.64 fl oz per 100 lbs seed</p> <p>(5 - 10 g ai per 100 kg seed)</p> <p>To deliver 0.005 - 0.01 mg ai per seed, apply 0.071 - 0.142 fl oz of the product per 100,000 seeds. (Based on an average of 4,500 cottonseeds per lb)</p> <p>Maximum number of application per year is 3.</p>	
Cereal grains Crop Group 15 and 16 (except corn, sorghum and rice) Wheat, barley, oats, buckwheat, millet (pearl and proso), rye, teosinte, triticale, wild rice,	Common bunt Covered smut False loose smut Flag smut Leaf strip Loose smut Stinking smut True loose smut Seed rot, pre-emergence damping-off and seedling blight caused by soilborne	<p>Apply 0.16 - 0.32 fl oz per 100 lbs seed</p> <p>(2.5 - 5 g ai per 100 kg seed)</p> <p>To deliver 0.00085 - 0.0017 mg ai per seed, apply 0.0125 - 0.025 fl oz of the product per 100,000 seeds. (Based on an average of 13,000 cereal grains per lb)</p>	

7/11

CROP	DISEASE	APPLICATION RATE PER 100 LBS SEED	DIRECTIONS
	<i>Rhizoctonia solani</i>	Maximum number of application per year is 3.	
Corn (field, sweet, popcorn), Sorghum	Seed rot and damping-off caused by <i>Rhizoctonia solani</i>	<p>Apply 0.16 - 0.64 fl oz per 100 lbs seed (2.5 - 10 g ai per 100 kg seed)</p> <p>Field Corn: To deliver 0.00675 - 0.027 mg ai per seed, apply 0.0757 - 0.303 fl oz of the product per 80,000 seeds. (Based on an average of 1,681 seeds per lb)</p> <p>Sweet Corn: To deliver 0.0035 - 0.014 mg ai per seed, apply 0.0505 - 0.202 fl oz of the product per 100,000 seeds. (Based on an average of 3,150 seeds per lb)</p> <p>Popcorn: To deliver 0.0037 - 0.015 mg ai per seed, apply 0.052 - 0.208 fl oz of the product per 100,000 seeds. (Based on an average of 3,061 seeds per lb)</p> <p>Sorghum: To deliver 0.0006 - 0.0025 mg ai per seed, apply 0.0087 - 0.035 fl oz of the product per 100,000 seeds. (Based on an average of 18,000 seeds per lb)</p> <p>Maximum number of application per year is 3.</p>	
Sunflower and Safflower	Seed rot and damping-off caused by <i>Rhizoctonia solani</i>	<p>Apply 0.32 - 0.96 fl oz per 100 lbs seed (5 - 15 g ai per 100 kg seed)</p> <p>Sunflower: To deliver 0.0041 - 0.0124 mg ai per seed, apply 0.0058 - 0.0174 fl oz of the product per 100,000 seeds. (Based on an average of 5,500 seeds per lb)</p> <p>Safflower: To deliver 0.0017 - 0.005 mg ai</p>	

8/11

CROP	DISEASE	APPLICATION RATE PER 100 LBS SEED	DIRECTIONS
		per seed, apply 0.023 - 0.07 fl oz of the product per 100,000 seeds. (Based on an average of 13,608 seeds per lb) Maximum number of application per year is 3.	
Rice	Seed rot and damping-off caused by <i>Rhizoctonia solani</i> Seedborne false smut (suppression) Seedborne kernel smut (suppression)	Apply 0.32 fl oz per 100 lbs seed (5 g ai per 100 kg seed) To deliver 0.001 mg ai per seed, apply 0.015 fl oz of the product per 100,000 seeds. (Based on an average of 21,850 seeds per lb) Maximum number of application per year is 3.	

SEED BAG TAG

The Federal Seed Act requires that the container of seed treated with EverGol Prime must be labeled with the following statements:

- This seed has been treated with EverGol Prime, which contains penflufen.
- Do not use treated seed for food, feed, or oil production. Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- User is responsible for ensuring that the seed bag meets all the requirements under the Federal Seed Act.

In addition, the US Environmental Protection Agency requires the following statements on the container of seed treated with EverGol Prime:

- Store away from feeds and foodstuffs.
- Do not allow children, pets or livestock to have access to treated seeds.
- Wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves when handling treated seed.
- After the seeds have been planted, do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: Once the seeds are planted in soil or other planting media, the Worker Protection Standard allows workers to enter the treated area without restriction if there will be no worker contact with the treated seeds in the soil or planting media
- Dispose of all excess treated seed. Left over treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- Use planting equipment that will plant treated seed into the soil to a minimum depth of 0.5 inch.
- Crops for which there are registered uses of penflufen, which includes alfalfa, potatoes (Crop Subgroup 1C), legume vegetables (Crop Group 6/7), cereal grains (Crop Group 15/16), and oil seeds (Crop Group 20) may be replanted at any time. All other crops may be planted no less than 30 days after planting treated seeds.
- This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.
- Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect seeds spilled during loading.
- Runoff may be hazardous to aquatic organisms in water adjacent to treated areas.
- Do not plant treated rice seed directly into flooded field. Do not soak treated rice seed.
- The maximum number of application per year is 3.
- Alfalfa: Do not exceed 0.00225 lb ai/A (1.02 g ai/A) per crop season.
- Beans and Peas (excluding soybeans): Do not exceed 0.0384 lb ai/A (17.5 g ai/A) per crop season.
- Soybeans: Do not exceed 0.0384 lb ai/A (17.5 g ai/A) per crop season.
- Oil Seed Crops: Do not exceed 0.00123 lb ai/A (0.6 g ai/A) per crop season.

- 9/11
- Cotton: Do not exceed 0.0019 lb ai/A (0.86 g ai/A) per crop season.
 - Cereal Grains (except corn, sorghum, and rice) do not exceed 0.0078 lb ai/A (3.5 g ai/A) per crop season.
 - Corn: Do not exceed 0.0033 lb ai/A (1.5 g ai/A) per crop season.
 - Sunflower and Safflower: Do not exceed 0.00123 lb ai/A (0.6 g ai/A) per crop season.
 - Rice (Dry seeded): Do not exceed 0.006 lb ai/A (2.7 g ai/A) per crop season.

INFORMATION

For use as a fungicide for protection against listed soilborne, seedborne, and post-emergence diseases of listed crop plants.

CROP ROTATION:

Listed seed treatment crops and potatoes may be replanted at any time. All other crops may be planted after 30 days.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

PESTICIDE STORAGE

Store in original container away from feed and food. Store in cool, dry area. Do not store in direct sunlight. Do not allow prolonged storage in temperatures that exceed 105°F (40°C) or in temperatures that fall below 14°F (-10°C).

PESTICIDE DISPOSAL

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER HANDLING

Rigid, Non-refillable containers (equal to or less than 5 gallons)

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill.

Rigid Non-refillable containers (greater than 5 gallons or 50 lbs)

Non-refillable Containers

Non-refillable containers - Do not reuse or refill this container. Refer to Bottom Discharge IBC or Top Discharge IBC, Drums, Kegs information as follows.

Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Top Discharge IBC, Drums, Kegs (e.g.– Snyder 120 Next Gen, Bonar B120, Drums, Kegs).

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. To triple rinse the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill.

Refillable Containers

Refillable container – Refer to Bottom Discharge IBC or Top Discharge IBC, Drums, Kegs information as follows. Refill this container with pesticide only. Do not reuse this container for any other purpose. Contact your Ag retailer or Bayer CropScience for container return, disposal and recycling information.

Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Top Discharge IBC, Drums, Kegs (e.g.– Snyder 120 Next Gen, Bonar B120, Drums, Kegs).

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To triple rinse the containers before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill.

End users are authorized to remove tamper evident cables as required to remove the product from the container unless the container is equipped with one way valves and refilling or returning is planned. If this is the case, end users are not authorized to remove tamper evident cables, one way valves or clean container.

11/11

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Bayer CropScience, no claims are made to guarantee germination of carry-over seed.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

NET CONTENTS:

PRODUCED FOR



Bayer CropScience

Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

EverGol Prime (PENDING) 10/09/13