


264-1121

5/11/2012

1/8

 <p style="text-align: center;">U S ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505C) 1200 Pennsylvania Ave N W Washington D C 20460</p> <p style="text-align: center;">NOTICE OF PESTICIDE <u> X </u> Registration <u> </u> Reregistration (under FIFRA as amended)</p>	EPA Reg Number 264-1121	Date of Issuance MAY 11 2012
	Term of Issuance Unconditional	
	Name of Pesticide Product Prosper EverGol	
<p>Name and Address of Registrant (include ZIP Code)</p> <p>Bayer CropScience LP 2 T W Alexander Drive P O Box 12014 Research Triangle Park, NC 27709</p>		
<p>Note Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce In any correspondence on this product always refer to the above EPA registration number</p>		
<p>On the basis of information furnished by the registrant the above named pesticide is hereby registered under the Federal Insecticide Fungicide and Rodenticide Act</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency In order to protect health and the environment the Administrator on his motion may at any time suspend or cancel the registration of a pesticide in accordance with the Act The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others</p> <p>This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you</p> <ol style="list-style-type: none"> 1 Submit and/or cite all data required for registration/registration review of your product when the Agency requires all registrant of similar products to submit such data 2 Make the following label changes before you release the product for shipment <ol style="list-style-type: none"> a Revise the EPA Reg No so it reads "EPA Reg No 264-1121 " b For clarity and consistency in the "Commercial Seed Treatment" table in the "Application Rates" column insert the information to deliver "xx g ai/100 lb seed" for canola, rapeseed and mustard c For clarity and consistency in the "Commercial Seed Treatment" table in the "Directions" column revise the tank mix instructions to read as follows "Pre-test all tank mixes to determine the physical compatibility between formulations Follow all cautions and most restrictive limitations on labeling of all products used in mixtures" d Add the batch code information Please be reminded that the 40 CFR part 156 140(a)(4) requires a batch code, lot number, or other code identifying the batch of the pesticide product distributed and sold be placed on <u>non-refillable</u> containers The code may appear either on the label or durably marked on the container itself 		

3 For future submissions, make the following changes to your Confidential Statement of Formula (EPA Form 8570-4)

a Box 4 Revise the EPA Reg No so it reads "EPA Reg No 264-1121 "

4 Submit 2 copies of your final printed label before release for shipment

Signature of Approving Official
Mark Corbin
Mark Corbin, Acting Branch Chief
Insecticide-Rodenticide Branch, Registration Division (7505P)

Date
MAY 11 2012

EPA Form 8570 6

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e) Your release for shipment of the product constitutes acceptance of these conditions A stamped copy of the label is enclosed for your records If you have any questions, please contact Marianne Lewis at 703-308-8043 or lewis marianne@epa gov

Mark Corbin
Acting Branch Chief,
Insecticide-Rodenticide Branch
Registration Division (7505P)

Enclosure

3/8

Prosper EverGol

For use as a fungicide and insecticide seed dressing for protection against listed soilborne, seedborne, and post-emergence diseases and insect pests of listed crop plants

ACTIVE INGREDIENTS

Clothianidin	22.3%
Penflufen	0.82%
Trifloxystrobin	0.55%
Metalaxyl	0.55%

OTHER INGREDIENTS

75.78%

TOTAL 100.00%

Contains 2.42 lb clothianidin 0.09 lb penflufen 0.06 lb trifloxystrobin 0.06 lb metalaxyl per gallon
Contains 290 g clothianidin 10.7 g penflufen 7.15 g trifloxystrobin 7.15 g metalaxyl per liter

EPA Reg No 264-XXXX

EPA Est


KEEP OUT OF REACH OF CHILDREN

CAUTION

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1 800-334-7577
For PRODUCT USE Information Call 1 866-99BAYER (1-866 992-2937)

FIRST AID

IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to by a poison control center or doctor • Do not give anything by mouth to an unconscious person
IF ON SKIN	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15-20 minutes • Call a poison control center or doctor for treatment advice
<p>In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No 1 800 334 7577 Have a product container or label with you when calling a poison control center or doctor or going for treatment</p>	



PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed Harmful if absorbed through skin Avoid contact with skin eyes or clothing

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are barrier laminate butyl rubber nitrile rubber neoprene rubber polyvinyl chloride or viton If you want more options follow the instructions for category C on an EPA chemical resistance category selection chart

All mixers, loaders, applicators and other handlers must wear

long sleeved shirt and long pants

socks plus shoes

chemical resistant gloves except when bagging or sewing bags of treated seeds

Follow manufacturer s instructions for cleaning/maintaining PPE If no such instructions for washables exist use detergent and hot water Keep and wash PPE separately from other laundry

USER SAFETY RECOMMENDATIONS

Wash hands before eating drinking chewing gum using tobacco or using the toilet

Remove clothing/PPE immediately if pesticide gets inside Then wash thoroughly and put on clean clothing

Remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing

ENVIRONMENTAL HAZARDS

This pesticide is highly toxic to bees exposed directly (contact) and is toxic to birds mammals fish and aquatic invertebrates Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark Runoff may be hazardous to aquatic organisms in water adjacent to treated areas Do not contaminate water when disposing of equipment washwater or rinsate Treated seed exposed on soil surface may be hazardous to wildlife Cover or collect seed spilled during loading

This product has properties and characteristics associated with chemicals detected in ground water This chemical may leach into ground water if used in areas where soils are permeable particularly where the water table is shallow Therefore contain any product spills or equipment leaks and dispose of wastes according to the disposal instructions on this label

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

For use only in commercial seed treatment equipment Not for use in hopper box planter box slurry box or other on farm seed treatment applications

COMMERCIAL SEED TREATMENT

CROP	DISEASE/PEST	APPLICATION RATE	DIRECTIONS
<p>Canola (including <i>Brassica napus</i> and <i>Brassica rapa</i>) Rapeseed and Mustard (oilseed and condiment)</p>	<p>For control of listed insect pests and for control of seed rot damping off seedling blight and early season root rot caused by <i>Pythium Rhizoctonia Fusarium</i> and <i>Alternaria</i>. Also controls seedborne Blackleg. The insecticide component clothianidin controls flea beetles on seeds and seedlings of canola (including <i>Brassica napus</i> and <i>Brassica rapa</i>) mustard (oilseed and condiment) and rapeseed. For protection of seeds and emerging seedlings only.</p>	<p>Apply 21.5 fl oz of Prosper EverGol to 100 lbs of seed (14 liters of product per 100 kgs of seed). Maximum number of application per year is 3.</p> <p><i>Pre-test</i></p>	<p>Apply using commercial slurry or mist type seed treatment equipment. Mix thoroughly before use. Use high quality properly cleaned seed. All tank mixes should be pre-tested to determine physical compatibility between formulations. Follow all cautions and limitations on labeling of all products used in mixtures. Prior to and during application Prosper EverGol must be thoroughly agitated to ensure uniform mixing.</p> <p><i>Pre-test</i></p> <p><i>and m restrict</i></p>

SEED BAG TAG

The Federal Seed Act requires that the container of seed treated with Prosper EverGol must be labeled with the following statements

- This seed has been treated with Prosper EverGol which contains metalaxyl, clothianidin, trifloxystrobin, and penflufen.
- Do not use treated seed for food, feed, or oil production. Excess treated seed may be used for ethanol production only if (1) by products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by products that are used in agronomic practice.
- User is responsible for ensuring that the seed bag meets all the requirements under the Federal Seed Act.

In addition, the US Environmental Protection Agency requires the following statements on the container of seed treated with Prosper EverGol:

- Store away from feeds and foodstuffs.
- Wear long sleeved shirt, long pants, shoes, socks, and chemical resistant gloves when handling treated seed.
- Areas planted with treated seed may be replanted immediately with canola, cotton, corn (field, pop, and sweet), cereal grains, except rice, mustard, rapeseed, sorghum, soybean, and potatoes (tuberous and corm subgroup 1C). These areas may also be replanted no less than 30 days after planting treated seeds with grasses, non grass livestock feeds, dried beans, fruiting vegetables (including cucurbits), leafy vegetables (including brassica vegetables), and sugar beets. Do not plant any other crop in the treated area for at least twelve months after treated seeds are planted.
- After the seeds have been planted, do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. Exception: Once the seeds are planted in soil or other planting media, the Worker Protection Standard allows worker to enter the treated area without restriction if there will be no worker contact with the treated seeds in the soil or planting media.
- Dispose of all excess treated seed. Left over treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- This contains metalaxyl that is known to leach through soil into ground water under certain conditions as result of agricultural use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.
- Dispose of seed packaging in accordance with local requirements.
- Use planting equipment that will plant treated seed into the soil to a minimum depth of 0.5 inch.
- This compound is toxic to birds and mammals. Treated seed exposed on soil surface may be hazardous to birds and mammals. Cover or collect seeds spilled during loading.
- This compound may be highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.
- Regardless of type of application, do not apply more than 0.2 lbs of the active ingredient clothianidin per acre per season.

- For Canola (including *Brassica napus* and *Brassica rapa*) rapeseed and Mustard (based on canola rapeseed or mustard seeding rate) Do not exceed 0.00123 lb penflufen/A (0.6 g ai/A) per crop season. The maximum number of applications per year is 3. This seed has been treated with xx mg active ingredients (ai) of Prosper EverGol /seed which includes penflufen (15 g ai/100 Kg seed), clothianidin (406.5 g ai/100 Kg seed), trifloxystrobin (10 g ai/100 Kg seed) and metalaxyl (10 g ai/100 Kg seed).
- Runoff may be hazardous to aquatic organisms in water adjacent to treated areas.
- This pesticide is toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark.

INFORMATION

Prosper EverGol is a insecticide and fungicide seed treatment for use on canola (including *Brassica napus* and *Brassica rapa*) mustard (oilseed and condiment) and rapeseed for control of listed insect pests and for control of seed rot, damping off, seedling blight and early season root rot caused by *Pythium*, *Rhizoctonia*, *Fusarium* and *Alternaria*. Also controls seedborne Blackleg. The insecticide component, clothianidin, controls flea beetles on seeds and seedlings of canola (including *Brassica napus* and *Brassica rapa*) mustard (oilseed and condiment) and rapeseed. For protection of seeds and emerging seedlings only.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

PESTICIDE STORAGE

Store in original container away from feed and food. Store in cool, dry area. Do not store in direct sunlight. Do not allow prolonged storage in temperatures that exceed 105°F (40°C) or in temperatures that fall below 14°F (-10°C).

PESTICIDE DISPOSAL

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER DISPOSAL

Rigid, Non refillable containers (equal to or less than 5 gallons)

Non refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill.

Rigid Non refillable containers (greater than 5 gallons or 50 lbs)

Non refillable Containers

Non refillable containers. Do not reuse or refill this container. Refer to Bottom Discharge IBC or Top Discharge IBC Drums, Kegs information as follows:

Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Top Discharge IBC, Drums, Kegs (e.g. – Snyder 120 Next Gen, Bonar B120 Drums, Kegs)

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. To triple rinse the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed offer for recycling if available or puncture and dispose of in a sanitary landfill

Refillable Containers

Refillable container – Refer to Bottom Discharge IBC or Top Discharge IBC Drums Kegs information as follows Refill this container with pesticide only Do not reuse this container for any other purpose Contact your Ag retailer or Bayer CropScience for container return disposal and recycling information

Bottom Discharge IBC (e g – Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the refiller To pressure rinse the container before final disposal empty the remaining contents from the IBC into application equipment or mix tank Raise the bottom of the IBC by 1 5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal Completely remove the top lid of the IBC Use water pressurized to at least 40 PSI to rinse all interior portions Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing Continue pressure rinsing for 2 minutes or until rinsate becomes clear Replace the lid and close bottom valve

Top Discharge IBC Drums Kegs (e g – Snyder 120 Next Gen Bonar B120 Drums Kegs)

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the refiller To triple rinse the containers before final disposal empty the remaining contents from this container into application equipment or mix tank Fill the container at least 10 percent full with water Agitate vigorously or recirculate water with the pump for 2 minutes Rinse all interior surfaces Pour or pump rinsate into application equipment or rinsate collection system Repeat this procedure two more times

Once container is rinsed offer for recycling if available or puncture and dispose of in a sanitary landfill

End users are authorized to remove tamper evident cables as required to remove the product from the container unless the container is equipped with one way valves and refilling or returning is planned If this is the case end users are not authorized to remove tamper evident cables one way valves or clean container

8/8

IMPORTANT READ BEFORE USE

Read the entire Directions for Use Conditions Disclaimer of Warranties and Limitations of Liability before using this product If terms are not acceptable return the unopened product container at once

By using this product user or buyer accepts the following Conditions disclaimer of Warranties and Limitations of Liability

Treatment of highly mechanically damaged seed or seed of known low vigor and poor quality may result in reduced germination and/or reduction of seed and seedling vigor Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment Due to seed quality conditions beyond the control of Bayer CropScience no claims are made to guarantee germination of carry over seed

CONDITIONS The directions for use of this product are believed to be adequate and must be followed carefully However it is impossible to eliminate all risks associated with the use of this product Crop injury ineffectiveness or other unintended consequences may result because of such factors as weather conditions presence of other materials or the manner of use or application all of which are beyond the control of Bayer CropScience All such risks shall be assumed by the user or buyer

DISCLAIMER OF WARRANTIES TO THE EXTENT CONSISTENT WITH APPLICABLE LAW BAYER CROPSCIENCE MAKES NO WARRANTIES EXPRESS OR IMPLIED OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein TO THE EXTENT CONSISTENT WITH APPLICABLE LAW BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT

LIMITATIONS OF LIABILITY TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT WARRANTY TORT NEGLIGENCE STRICT LIABILITY OR OTHERWISE SHALL NOT EXCEED THE PURCHASE PRICE PAID OR AT BAYER CROPSCIENCE S ELECTION THE REPLACEMENT OF PRODUCT

NET CONTENTS

PRODUCED FOR



Bayer CropScience

**Bayer CropScience LP
P O Box 12014, 2 T W Alexander Drive
Research Triangle Park, North Carolina 27709
1 866-99BAYER (1 866 992-2937)**

PROSPER EVERGOL (PENDING) 03/31/10 Resubmitted 12/06/11 03/20/12 04 17 12 05 11 12V3