



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

June 30, 2021

Dave Wilson
Senior Regulatory Affairs Manager
Bayer U.S.- Crop Science
700 Chesterfield Pkwy W
Chesterfield, MO 63017

Subject: Registration Review Label Mitigation for Metalaxyl
Product Name: EVERGOL ENERGY
EPA Registration Number: 264-1122
Application Date: 06/28/2018
Decision Number: 566383

Dear Mr. Wilson:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with Metalaxyl Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Page 2 of 2
EPA Reg. No. 264-1122
Decision No. 566383

If you have any questions about this letter, please contact Srijana Shrestha by phone at 703-305-6471, or via email at shrestha.srijana@epa.gov.

Sincerely,

A handwritten signature in blue ink, appearing to read "Linda Arrington".

Linda Arrington, Branch Chief
Risk Management and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

Enclosure: Stamped Label

PROTHIOCONAZOLE	GROUP	3	FUNGICIDE
PENFLUFEN	GROUP	7	FUNGICIDE
METALAXYL	GROUP	4	FUNGICIDE

EverGol Energy

ABN: Acceleron D-961 Fungicide Seed Treatment

For: Use as a fungicide seed dressing for protection against listed soilborne, seedborne, and early season post-emergence diseases of listed crop plants.

ACTIVE INGREDIENTS:

Prothioconazole:	07.18%
Penflufen:	03.59%
Metalaxyl:	05.74%
OTHER INGREDIENTS:	83.49%
TOTAL:	100.00%

Contains 0.64 pounds Prothioconazole per U.S. gallon (76.8 grams AI/liter)

Contains 0.32 pounds Penflufen per U.S. gallon (38.4 grams AI/liter)

Contains 0.51 pounds Metalaxyl per U.S. gallon (61.4 grams AI/liter)

EPA Reg. No. 264-1122

EPA Est.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours a Day 1-800-334-7577
For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)


Please refer to [back panel] [booklet] for additional precautionary statements and directions for use. [Note to reviewer: Location of additional precautionary statements and directions for use will vary between those listed, depending on container type/size.]

Net Contents:

PRODUCED FOR



Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167
1-866-99BAYER (1-866-992-2937)



FIRST AID

If Swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on Skin or Clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

- Harmful if swallowed.
- Harmful if absorbed through skin.
- Avoid contact with skin, eyes, or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride, or viton.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Socks plus shoes
- Chemical resistant gloves, except when bagging or sewing bags of treated seeds

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, and freshwater/estuarine/marine aquatic plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate. Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect seed spilled during loading.

Ground Water Advisory

This product has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow. Therefore contain any product spills or equipment leaks and dispose of wastes according to the disposal instructions on this label.

CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Bayer CropScience, no claims are made to guarantee germination of carry-over seed.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

DIRECTIONS FOR USE

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Read the entire label before using this product.**

For use only in commercial seed treatment equipment. Do not use in hopper box, planter box, slurry box, or other on-farm seed treatment applications.

PRODUCT INFORMATION

EverGol Energy:

- is a fungicide seed dressing for protection against listed soil-borne, seed-borne and early season post-emergence diseases of listed crop plants.

APPLICATION INSTRUCTIONS

- Apply using commercial slurry or mist-type seed treatment equipment.
- Uniform application of seed is necessary to ensure seed safety and best disease protection.
- Dilute product with sufficient water to ensure complete seed coverage
- Apply to high quality, properly cleaned seed.

USE RESTRICTIONS

- For use only in commercial seed treatment facilities. Do not use in hopper box, planter box, slurry box, or other on-farm seed treatment applications.
- Maximum usage when applying both metalaxyl- and mefenoxam- containing products to the same crop within the same season: Do not apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop. The maximum application rate for metalaxyl- and mefenoxam-containing products (including seed treatments, foliar applications, soil application) is 12.3 lb per acre per calendar year. DO NOT apply more than 12.3 lb of active ingredient metalaxyl per acre per calendar year.
- All seed must be adequately dyed with a suitable colorant to prevent its accidental use as food for man or feed for animals. Refer to 21CFR, part 2.25. Any colorant added to treated seed must be cleared for use under 40CFR, part 153.155.

Refer to the specific use directions and restrictions in each Crop, Crop Group or Crop Subgroup table.

Compatibility

It is essential that before using EverGol Energy seed treatment in any tank mixture the compatibility of the mixture be established. Add EverGol Energy at the labeled rate to a clean quart jar containing approximately one-half the amount of water intended for a final slurry application rate. Next, follow with all other tank mix components that will be used in the total slurry application. Add last the remaining balance of water. The total amount of volume is determined by the seed size and how much is necessary to ensure complete and uniform coverage and distribution on the seed, as well as the type of commercial seed treating application equipment that will be used.

DO NOT USE MIXTURES THAT CURDLE, PRECIPITATE, OR GEL. USE TANK MIXTURES IMMEDIATELY AFTER MIXING WITH ADEQUATE AGITATION.

RESISTANCE MANAGEMENT RECOMMENDATION

For resistance management, please note that EverGol Energy contains a Group 3 (demethylation inhibitors), Group 7 (succinate dehydrogenase inhibitor) and Group 4 (phenylamides) fungicide. Any fungal population may contain individuals naturally resistant to EverGol Energy and other Group 3, Group 7, or Group 4 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of EverGol Energy or other Group 3, 7 and 4 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Bayer CropScience LP at 1-866-99BAYER (1-866-992-2937). You can also contact your pesticide distributor or university extension specialist to report resistance.

SPECIFIC CROP DIRECTIONS

CROP USE DIRECTIONS

COMMERCIAL SEED TREATMENT

ALFALFA	
Disease(s) Controlled	Product Rate (Per 100 lbs Seed)
Seed rot and damping-off caused by <i>Rhizoctonia</i>	3.0 fl oz
Crop Specific Directions and Restrictions: <ul style="list-style-type: none"> • To deliver 0.0006 – 0.0008 mg ai per seed, apply 0.013 – 0.015 fl oz of the product per 100,000 seeds. (Based on an average of 213,000 seeds per lb). 	

BEANS AND PEAS (DRIED)	
Crops of Crop Group 6-C and Soybean Including: Adzuki Bean, Black-eyed Pea, Broad Bean, Catjang, Chickpea, Cowpea, Crowder Pea, Field Bean, Field Pea, Guar, Kidney Bean, Lablab Bean, Lentil, Lima Bean, Moth Bean, Mung Bean, Navy Bean, Pigeon Pea, Pinto Bean, Rice Bean, Southern Pea, Soybean, Tepary Bean, and Urd Bean	
Disease(s) Controlled	Product Rate (Per 100 lbs Seed)
Seed rot and damping-off caused by <i>Rhizoctonia</i> , <i>Fusarium</i> and <i>Pythium</i> ¹	1.0 fl oz
Seed decay caused by <i>Phomopsis</i>	
Seed decay	
Disease(s) Suppressed	
<i>Ascochyta</i> blight	
Seed rot, damping-off, and seedling blight caused by seedborne <i>Botrytis cinerea</i>	
Crop Specific Directions and Restrictions: <ul style="list-style-type: none"> • To deliver 0.017 - 0.021 mg ai per seed, apply 0.45 – 0.55 fl oz of the product per 140,000 soybean seeds. (Based on an average of 2,800 soybeans per lb). • To deliver 0.02943 – 0.03597 mg ai per seed, apply 0.5625 – 0.6875 fl oz of the product per 100,000 bean seeds. (Based on an average of 1,600 beans per lb). • To deliver 0.01485 – 0.01815 mg ai per seed, apply 0.2826 – 0.3454 fl oz of the product per 100,000 pea seeds. (Based on an average of 3,181 peas per lb). 	
¹ Early season <i>Phytophthora</i> . For longer season control add sufficient metalaxyl product (such as, Allegiance) to supply a total metalaxyl amount of 15 to 30 gal/100kg of seed.	

CEREAL GRAINS

Crops of Crop Group 15 (except corn, sorghum, and rice) including: Barley, Triticale, Wheat, Oats, Rye, Buckwheat, and Millet (Pearl and Proso)

Disease(s)	Product Rate (Per 100 lbs Seed)
<p align="center">Disease(s) Controlled</p> Common bunt Covered smut False loose smut Flag smut Leaf stripe Loose smut Stinking smut Stem smut True loose smut Seed rot, pre-emergence damping-off and seedling blight caused by soilborne and seed-borne <i>Rhizoctonia</i> , <i>Fusarium</i> , <i>Cochliobolus</i> and <i>Pythium</i> Seed decay	1.0 fl oz
<p align="center">Disease(s) Suppressed</p>	
Common root rot, foot rot, and crown rot Rust, <i>Septoria</i> and powdery mildew	

Crop Specific Directions and Restrictions:

- To deliver 0.0036 – 0.0044 mg ai per seed, apply 0.0693 – 0.0847 fl oz of the product per 100,000 wheat and barley seeds. (Based on an average of 13,000 wheat or barley seeds per lb).
- To deliver 0.00252 – 0.00308 mg ai per seed, apply 0.0477 – 0.0583 fl oz of the product per 100,000 Triticale seeds. (Based on an average of 18,840 Triticale seeds per lb).
- To deliver 0.0026 - 0.0032 mg ai per seed, apply 0.05 – 0.062 fl oz of the product per 100,000 oat and rye seeds. (Based on an average of 18,000 oat or rye seeds per lb).
- To deliver 0.0027 - 0.0033 mg ai per seed, apply 0.051 – 0.062 fl oz of the product per 100,000 buckwheat seeds. (Based on an average of 17,650 buckwheat seeds per lb)
- To deliver 0.0005 – 0.0007 mg ai per seed, apply 0.011 – 0.013 fl oz of the product per 100,000 millet seeds. (Based on an average of 84,000 millet seeds per lb)

CORN (FIELD CORN, POPCORN, AND SWEET CORN)	
Disease(s) Controlled	Product Rate (Per 100 lbs Seed)
Seed rot and damping-off caused by <i>Rhizoctonia</i> , <i>Fusarium</i> and <i>Pythium</i>	0.5 – 2.0 fl oz
Seed decay	
Crop Specific Directions and Restrictions:	
<ul style="list-style-type: none"> To deliver 0.0156 – 0.0623 mg ai per seed, apply 0.24 – 0.95 fl oz of the product per 80,000 field corn seeds. (Based on an average of 1,681 field corn per lb). To deliver 0.0086 -0.0342 mg ai per seed, apply 0.16 – 0.65 fl oz of the product per 100,000 popcorn seeds. (Based on an average of 3,061 popcorn per lb). To deliver 0.0083 – 0.0332 mg ai per seed, apply 0.16 – 0.64 fl oz of the product per 100,000 sweet corn seeds. (Based on an average of 3,150 sweet corn per lb). 	

RICE	
Disease(s)	Product Rate (Per 100 lbs Seed)
Disease(s) Controlled	1.0 – 2.0 fl oz
Seed rot and damping-off caused by <i>Rhizoctonia</i> , <i>Fusarium</i> and <i>Pythium</i>	
Seed decay	
Disease(s) Suppressed	
Seed-borne false smut	
Seed-borne kernel smut	
Crop Specific Directions and Restrictions:	
<ul style="list-style-type: none"> To deliver 0.0024 – 0.0048 mg ai per seed, apply 0.046 – 0.092 fl oz of the product per 100,000 Rice seeds. (Based on an average of 21,850 Rice seeds per lb). Must use high rate for moderate to high seedling disease pressure or for suppression of seed-born false smut and kernel smut. 	

SORGHUM	
Disease(s) Controlled	Product Rate (Per 100 lbs Seed)
Seed rot and damping-off caused by <i>Rhizoctonia</i> , <i>Fusarium</i> and <i>Pythium</i>	0.5 – 2.0 fl oz
Seed decay	
Crop Specific Directions and Restrictions:	
<ul style="list-style-type: none"> To deliver 0.0015 – 0.006 mg ai per seed, apply 0.028 – 0.111 fl oz of the product per 100,000 seeds. (Based on an average of 18,000 seeds per lb) 	

SUGAR BEET		
Disease(s)	Product Rate (Per 100 lbs Raw Seed)	Product Rate (Per raw seed unit*)
Seed rot and damping-off caused by <i>Rhizoctonia solani</i> , <i>Fusarium</i> and <i>Pythium</i>	0.9 - 1.0 fl oz (10.35 - 11.51 g ai per 100 kg seed)	0.019 - 0.022 fl oz (0.001 - 0.0011 mg ai/seed)
Crop Specific Directions and Restrictions:		
<ul style="list-style-type: none"> Apply EverGol Energy as a commercial seed treatment to pelleted sugar beet seeds at the rate given in the application table above using commercial slurry or mist-type seed treatment equipment. Consult a seed treatment specialist regarding specified slurry rates for the crop to be treated with EverGol Energy. Use high quality, properly cleaned seed. To deliver 0.0011 mg ai per seed, apply 0.022 fl oz of the product per 100,000 seeds. (*Based on a unit of raw sugarbeet seed unit of 100,000 seeds weighing 2.2 lbs. or 45,454 seeds per lb.) 		

SEED TAG LABELING

The Federal Seed Act requires that the container of seed treated with EverGol Energy must be labeled with the following statements:

- This seed has been treated with:
 - **EverGol Energy**, which contains **7.18% Prothioconazole**, **3.59% Penflufen**, and **5.74% Metalaxyl**
- Do not use treated seed for food, feed, or oil production. User is responsible for ensuring that the seed bag meets all the requirements under the Federal Seed Act.

The following statements are also required on the container of seed treated with **EverGol Energy**:

- Crops on this label and other crops for which tolerances exist for metalaxyl, prothioconazole and penflufen may be replanted at any time. All other crops may be planted after 30 days.
- **Store treated seeds away from food and feedstuffs.**
- **Do not allow children, pets, or livestock to have access to treated seed.**
- **Wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves when handling treated seed.**
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- This compound is toxic to birds and mammals. Treated seeds exposed on soil surface may be hazardous to birds and mammals. Cover or collect treated seeds spilled during loading.
- Dispose of all excess treated seed. Leftover treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements.
- Do not contaminate water bodies when disposing of planting equipment wash waters.
- Treated seed must be adequately covered with soil at planting.
- Use planting equipment that will plant treated seed into the soil to a minimum depth of 0.5 inch or following local Agronomical practice.
- Dispose of seed packaging or containers in accordance with local requirements. Do not use empty seed bags for any other purpose.
- To reduce seed dust which can drift onto blooming crops or weeds, ensure that planting equipment is functioning properly in accordance with manufacturer's recommendations. Surplus seed or empty seed containers should be stored or disposed according to state or local regulations.
- After the seeds have been planted, do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours. Exception: Once the seeds are planted in soil or other planting media, the Worker Protection Standard allows worker to enter the treated area without restriction if there will be no worker contact with the treated seeds in the soil or planting media.
- This product contains chemicals which are known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.
- Do not plant treated rice seed directly into flooded field. Do not soak treated rice seed.
- For alfalfa: Do not exceed 0.00225 lb penflufen /A (1.02 g ai/A) per crop season.
- For beans (excluding soybeans): Do not exceed 0.0384 lb penflufen /A (17.5 g ai/A) per crop season.
- For peas: Do not exceed 0.0384 lb penflufen /A (17.5 g ai/A) per crop season.
- For soybeans: Do not exceed 0.0384 lb penflufen /A (17.5 g ai/A) per crop season.
- For Barley and Wheat: Do not exceed 0.0078 lb penflufen /A (3.5 g ai/A) per crop season.
- For Triticale: Do not exceed 0.0078 lb penflufen /A (3.5 g ai/A) per crop season.
- For rice (dry seeded): Do not exceed 0.006 lb penflufen /A (2.7 g ai/A) per crop season.
- For Oats and Rye: Do not exceed 0.0078 lb penflufen /A (3.5 g ai/A) per crop season.
- For Buckwheat: Do not exceed 0.0078 lb penflufen /A (3.5 g ai/A) per crop season.
- For Millet: Do not exceed 0.0078 lb penflufen /A (3.5 g ai/A) per crop season.
- For Field Corn: Do not exceed 0.0078 lb penflufen /A (3.5 g ai/A) per crop season.
- For Popcorn: Do not exceed 0.0078 lb penflufen /A (3.5 g ai/A) per crop season.
- For Sweet Corn: Do not exceed 0.0078 lb penflufen /A (3.5 g ai/A) per crop season.
- For Sorghum: Do not exceed 0.0078 lb penflufen /A (3.5 g ai/A) per crop season.
- This product has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow. Therefore contain any product spills or equipment leaks and dispose of wastes according to the disposal instructions on this label.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)
For MEDICAL, TRANSPORTATION or OTHER Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

Pesticide Storage

Store in original container away from feed and food. Store in cool, dry area. Do not store in direct sunlight. Do not allow prolonged storage in temperatures that exceed 105°F (40°C) or in temperatures that fall below 14°F (-10°C).

Pesticide Disposal

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling

[Dilutable Seed Treatment Products in Non-Refillable Plastic Containers]

Rigid, Non-refillable containers (equal to or less than 5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

Rigid Non-refillable containers (greater than 5 gallons or 50 lbs)

Non-refillable Containers

Non-refillable containers - Do not reuse or refill this container. Refer to Bottom Discharge IBC or Top Discharge IBC, Drums, Kegs information as follows.

Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Top Discharge IBC, Drums, Kegs (e.g.– Snyder 120 Next Gen, Bonar B120, Drums, Kegs).

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. To triple rinse the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

Refillable Containers

Refillable container – Refer to Bottom Discharge IBC or Top Discharge IBC, Drums, Kegs information as follows. Refill this container with pesticide only. Do not reuse this container for any other purpose. Contact your Ag retailer or Bayer CropScience for container return, disposal and recycling information.

Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Top Discharge IBC, Drums, Kegs (e.g.– Snyder 120 Next Gen, Bonar B120, Drums, Kegs).

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To triple rinse the containers before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

End users are authorized to remove tamper evident cables as required to remove the product from the container unless the container is equipped with one way valves and refilling or returning is planned. If this is the case, end users are not authorized to remove tamper evident cables, one way valves or clean container.

EverGol Energy (PENDING) 06/13/2018, 06/14/2018, 09/15/2020, 06/21/2021