



## OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

January 17, 2024

Megan Mader  
Regulatory Affairs Manager, Fruits and Vegetables  
Bayer CropScience LP  
800 N. Lindbergh Blvd.  
St. Louis, MO 63167

Subject: Label Amendment - Registration Review Mitigation for Trifloxystrobin  
Product Name: EVERGOL XTEND  
EPA Registration Number: 264-1124  
Application Date: 09/11/2019 and 06/01/2022  
Decision Number: 555167

Dear Megan Mader:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Trifloxystrobin Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling

before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact DeMariah Koger by phone at (202)-566-2288, or via email at [Koger.demariah@epa.gov](mailto:Koger.demariah@epa.gov).

Sincerely,

A handwritten signature in blue ink, appearing to read "Linda Arrington".

Linda Arrington, Branch Chief  
Risk Management and Implementation Branch 4  
Pesticide Re-Evaluation Division  
Office of Pesticide Programs

ENCLOSURE: Stamped label

**ACCEPTED**

Jan 17, 2024

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 264-1124

PENFLUFEN	GROUP	<b>7</b>	FUNGICIDE
TRIFLOXYSTROBIN	GROUP	<b>11</b>	FUNGICIDE

# EverGol® Xtend

For use as a fungicide seed dressing for protection against listed soilborne, seedborne, and early season post-emergence diseases of listed crop plants.

**ACTIVE INGREDIENTS:**

Penflufen .....	13.3%
Trifloxystrobin .....	13.3%

**OTHER INGREDIENTS:** .....

	73.4%
<b>TOTAL :</b>	100.0%

Contains 1.28 lb. penflufen; 1.28 lb. trifloxystrobin per gallon (154 g penflufen; 154 g trifloxystrobin per liter)

**EPA Reg. No. 264-1124** **EPA Est.**

## KEEP OUT OF REACH OF CHILDREN

### CAUTION

### PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577  
 For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

See [Back][Side] Panel for First Aid Instructions and [Leaflet][Booklet] for Complete Precautionary Statements and Directions for Use. (Note to reviewer: Location of additional precautionary statements, directions for use will vary between those listed, depending on container type/size.)

### FIRST AID

<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF ON SKIN:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>

**In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.**

# PRECAUTIONARY STATEMENTS

## CAUTION

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed, if absorbed through the skin, or inhaled. Causes moderate eye irritation. Avoid breathing vapor or dust and contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

**All mixers, loaders, applicators and other handlers must wear:**

- long sleeved shirt and long pants,
- socks plus shoes,
- waterproof gloves, except when bagging or sewing bags of treated seeds,

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Applying this product when rain is not predicted for the next 24 hours will help reduce potential risk to aquatic invertebrates by reducing pesticide runoff from the treatment area into water bodies. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift or run-off from treated areas is hazardous to fish and aquatic organisms in water adjacent to treated areas. Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect seed spilled during loading.

This product has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow. Therefore contain any product spills or equipment leaks and dispose of wastes according to the disposal instructions on this label.

# DIRECTIONS FOR USE

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

For use only in commercial seed treatment equipment. Not for use in hopper box, planter box, slurry box, or other on-farm seed treatment applications.

## COMMERCIAL SEED TREATMENT

CROPS	DISEASES	APPLICATION RATES PER 100 LBS SEED	DIRECTIONS
<p><b>Alfalfa</b></p>	<p>Seed rot and damping-off caused by <i>Rhizoctonia solani</i></p>	<p>Apply 1 – 1.5 fl oz</p> <p><b>NOTE:</b> To deliver 0.0004 – 0.0006 mg ai per seed, apply 0.005 – 0.007 fl oz of the product per 100,000 seeds. (Based on an average of 213,000 seeds per lb)</p> <p>Maximum number of application per year is 3.</p>	<p>Apply using commercial slurry or mist-type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Dilute product with sufficient water to ensure complete seed coverage.</p> <p>Use high quality, properly cleaned seed.</p>
<p><b>Beans and Peas including Soybean (Crop Group 6)</b></p> <p>Soybean, Adzuki Bean, Asparagus Bean, Broad Bean (Succulent or Dry), Catjang Bean, Chinese Longbean, Field Bean, Guar Bean, Jackbean, Kidney Bean, Lablab Bean, Lima Bean (Succulent or Dry), Moth Bean (Succulent or Dry), Mung Bean, Navy Bean, Pinto Bean, Rice Bean, Runner Bean, Snap Bean, Sword Bean, Tepary Bean, Urd Bean, Wax Bean, Yardlong Bean, Blackeyed Pea (Succulent or Dry), Chickpea, Cowpea (Succulent or Dry), Crowder Pea, Dwarf Pea, Edible-Pod Pea, English Pea, Field Pea, Garden Pea, Green Pea, Pigeon Pea (Succulent or Dry), Snow Pea, Southern Pea (Succulent or Dry), Sugar Snap Pea, Grain Lupin, White Lupin, Sweet Lupin, White Sweet Lupin, Lentil</p>	<p>Seed rot and damping-off caused by <i>Rhizoctonia</i>, and <i>Fusarium</i></p> <p>Seed rot, damping-off, and seedling blight caused by seedborne <i>Botrytis cinerea</i></p> <p>Seed decay caused by <i>Phomopsis</i></p> <p><i>Ascochyta</i> blight (suppression)</p> <p>Seed decay</p>	<p>Apply 0.25 – 0.5 fl oz (5 – 10 g ai per 100 kg seed)</p> <p><b>NOTE:</b> To deliver 0.0081-0.016 mg ai per seed, apply 0.125 -0.25 fl oz of the product per 140,000 seeds. <sup>1</sup></p> <p>Maximum number of application per year is 3.</p>	<p>All seed must be adequately dyed with a suitable colorant to prevent its accidental use as food for man or feed for animals. Refer to 21CFR, part 2.25. Any colorant added to treated seed must be cleared for use under 40CFR, part 153.155.</p>
<p><b>Wheat</b></p>	<p>Loose smut</p> <p>Seed rot, pre-emergence damping-off and seedling blight caused by soilborne <i>Rhizoctonia solani</i>,</p> <p>Seed decay</p>	<p>Apply 0.25 – 0.5 fl oz</p> <p><b>NOTE:</b> To deliver 0.0018– 0.0035 mg ai per seed, apply 0.019 – 0.038 fl oz of the product per 100,000 wheat seeds. (Based on an average of 13,000 wheat seeds per lb)</p> <p>Maximum number of application per year is 3.</p>	

<p><b>Corn (field, sweet, popcorn), Sorghum</b></p>	<p>Seed rot and damping-off caused by <i>Rhizoctonia</i> and <i>Fusarium</i></p> <p>Seed decay</p>	<p>Apply 0.25 – 0.5 fl oz</p> <p><b>NOTE:</b> To deliver 0.013-0.026 mg ai per seed, apply 0.114 - 0.228 fl oz of the product per 80,000 seeds.<sup>2</sup></p> <p>Maximum number of application per year is 3.</p>	
<p><b>Cotton</b></p>	<p>Seed rot and damping-off caused by <i>Rhizoctonia solani</i></p>	<p>Apply 0.25 – 1.0 fl oz</p> <p><b>NOTE:</b> To deliver 0.0045 – 0.018 mg ai per seed, apply 0.05-0.2 fl oz of the product per 100,000 seeds.<sup>3</sup></p> <p>Maximum number of application per year is 3.</p>	
<p><b>Rice</b></p>	<p>Seed rot and damping-off caused by <i>Rhizoctonia</i> and <i>Fusarium</i></p> <p>Seed decay</p> <p>Seed-borne false smut (suppression)</p> <p>Seed-borne kernel smut (suppression)</p>	<p>Apply 0.25 – 0.5 fl oz</p> <p><b>NOTE:</b> To deliver 0.0010 – 0.0021 mg ai per seed, apply 0.011 – 0.023 fl oz of the product per 100,000 Rice seeds. (Based on an average of 21,850 Rice seeds per lb)</p> <p>Maximum number of application per year is 3.</p>	

<sup>1</sup> Based on 2,800 soybean seeds per pound.

<sup>2</sup> Based on 1,750 corn seed per pound.

<sup>3</sup> Based on 4,800 cottonseed per pound.

## SEED BAG TAG

The Federal Seed Act requires that the container of seed treated with EverGol Xtend must be labeled with the following statements:

- This seed has been treated with EverGol® Xtend which contains Trifloxystrobin and Penflufen.
- Do not use treated seed for food, feed, or oil production. Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- User is responsible for ensuring that the seed bag meets all the requirements under the Federal Seed Act.
- Rice paddy water must be held for a minimum of 7 days after application.

In addition, the US Environmental Protection Agency requires the following statements on the container of seed treated with EverGol Xtend:

- Store away from feeds and foodstuffs.
- Wear long sleeved shirt, long pants, shoes, socks, and chemical resistant gloves when handling treated seed.
- Crops on this label and other crops for which tolerances exist for trifloxystrobin and penflufen may be replanted at any time. All other crops may be planted after 30 days.
- After the seeds have been planted, do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: Once the seeds are planted in soil or other planting media, the Worker Protection Standard allows worker to enter the treated area without restriction if there will be no worker contact with the treated seeds in the soil or planting media.
- Dispose of all excess treated seed. Left over treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- Use planting equipment that will plant treated seed into the soil to a minimum depth of 0.5 inch.
- This compound is toxic to birds and mammals. Treated seed exposed on soil surface may be hazardous to birds and mammals. Cover or collect seeds spilled during loading.
- Do not plant treated rice seed directly into flooded field. Do not soak treated rice seed.
- For alfalfa: Do not exceed 0.00225 lb. penflufen /A (1.02 g ai/A) per crop season. The maximum number of applications per year is 3. This seed has been treated with xx mg active ingredients (ai) of EverGol Xtend /seed which includes penflufen (5-15 g ai/100 Kg seed) and trifloxystrobin (5-15 g ai/100 Kg seed).
- For beans and peas (excluding soybeans): Do not exceed 0.0384 lb. penflufen /A (17.5 g ai/A) per crop season. The maximum number of applications per year is 3. This seed has been treated with xx mg active ingredients (ai) of EverGol Xtend /seed which includes penflufen (2.5-5 g ai/100 Kg seed) and trifloxystrobin (2.5-5 g ai/100 Kg seed).
- For soybeans: Do not exceed 0.0384 lb. penflufen /A (17.5 g ai/A) per crop season. The maximum number of applications per year is 3. This seed has been treated with xx mg active ingredients (ai) of EverGol Xtend /seed which includes penflufen (2.5-5 g ai/100 Kg seed) and trifloxystrobin (2.5-5 g ai/100 Kg seed).
- For Barley, Triticale, and Wheat (based on wheat seeding rate): Do not exceed 0.0078 lb. penflufen /A (3.5 g ai/A) per crop season. The maximum number of applications per year is 3. This seed has been treated with xx mg active ingredients (ai) of EverGol Xtend /seed which includes penflufen (2.5-5 g ai/100 Kg seed) and trifloxystrobin (2.5-5 g ai/100 Kg seed).
- For Cotton (based on cotton seeding rate): Do not exceed 0.0019 lb. penflufen /A (0.86 g ai/A) per crop season. The maximum number of applications per year is 3. This seed has been treated with xx mg active ingredients (ai) of EverGol Xtend /seed which includes penflufen (2.5-5 g ai/100 Kg seed) and trifloxystrobin (2.5-5 g ai/100 Kg seed).
- For Corn (field, sweet, popcorn) and sorghum (based on corn or sorghum seeding rate): Do not exceed 0.0033 lb. penflufen /A (1.5 g ai/A) per crop season. The maximum number of applications per year is 3. This seed has been treated with xx mg active ingredients (ai) of EverGol Xtend /seed which includes penflufen (2.5-5 g ai/100 Kg seed) and trifloxystrobin (2.5-5 g ai/100 Kg seed).
- For rice (dry seeded): Do not exceed 0.006 lb. Penflufen /A (2.7 g ai/A) per crop season. The maximum number of applications per year is 3. This seed has been treated with xx mg active ingredients (ai) of EverGol Xtend /seed which includes penflufen (2.5-5 g ai/100 Kg seed) and trifloxystrobin (2.5-5 g ai/100 Kg seed).
- This product has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow. Therefore contain any product spills or equipment leaks and dispose of wastes according to the disposal instructions on this label.

## FUNGICIDE RESISTANCE MANAGEMENT

For resistance management, please note that EverGol Xtend contains both a Group 7 and Group 11 fungicide. Any fungal population may contain individuals naturally resistant to EverGol Xtend and other Group 7 or Group 11 fungicides. A gradual or total loss of pest control may occur over time if these (fungicides) are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed. [Note to reviewer: We disagree with the statement "Rotate the use of (name of product) or other Group (mode of action group number) fungicides within a growing season sequence with different groups that control the same pathogens. Evergol Xtend is used for seed treatment, and seed treatment scenarios are different compared to multiple foliar applications where the risk of resistance development is much higher. We believe the forementioned statement would lead to confusion for the grower since seed treatment is a once off application and there are no opportunities to rotate. We are already addressing the management strategy of using different groups because EverGol Xtend is a combination product of two active ingredients with two different MoA's. The foliar fungicides that would be used after emergence have no effect on seed and seedling diseases on the EverGol Xtend label. In the case of Aschochyta and loose smut we recommend adding the statement "Because of the minimal amount of EverGol Xtend applied to the seed that relocates to the upper parts of the plant and the dual mode of action, the potential for development of resistance is low. Seed treatments of EverGol Xtend can, therefore, be followed by foliar use and resistance management guidelines of foliar use products containing the same mode of action as specified on the respective labels."]

To delay fungicide resistance, take one or more of the following steps:

- Because of minimal amount of EverGol Xtend applied to seed that relocates to the upper parts of the plant and the dual mode of action, the potential for development of resistance is low. Seed treatments of EverGol Xtend can, therefore, be followed by foliar use and resistance management guidelines of foliar use products containing the same mode of action as specified on the respective labels.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Bayer CropScience at 1-866-99BAYER (1-866-992- 2937). You can also contact your pesticide distributor or university extension specialist to report resistance.

## INFORMATION

EverGol Xtend is a fungicide seed dressing for protection against listed soilborne, seedborne, and early season post-emergence diseases of listed crop plants.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

### PESTICIDE STORAGE

Store in original container away from feed and food. Store in cool, dry area. Do not store in direct sunlight. Do not allow prolonged storage in temperatures that exceed 105°F (40°C) or in temperatures that fall below 14°F (-10°C).

### PESTICIDE DISPOSAL

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

### CONTAINER DISPOSAL

#### **Rigid, Non-refillable containers (equal to or less than 5 gallons)**

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.



Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill.

### **Rigid Non-refillable containers (greater than 5 gallons or 50 lbs)**

#### **Non-refillable Containers**

Non-refillable containers - Do not reuse or refill this container. Refer to Bottom Discharge IBC or Top Discharge IBC, Drums, Kegs information as follows.

Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal.

Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Top Discharge IBC, Drums, Kegs (e.g.– Snyder 120 Next Gen, Bonar B120, Drums, Kegs).

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. To triple rinse the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill.

#### **Refillable Containers**

Refillable container – Refer to Bottom Discharge IBC or Top Discharge IBC, Drums, Kegs information as follows. Refill this container with pesticide only. Do not reuse this container for any other purpose. Contact your Ag retailer or Bayer CropScience for container return, disposal and recycling information.

Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Top Discharge IBC, Drums, Kegs (e.g.– Snyder 120 Next Gen, Bonar B120, Drums, Kegs).

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To triple rinse the containers before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill.

End users are authorized to remove tamper evident cables as required to remove the product from the container unless the container is equipped with one way valves and refilling or returning is planned. If this is the case, end users are not authorized to remove tamper evident cables, one way valves or clean container. See Container Disposal instructions under Storage and Disposal.

## IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, disclaimer of Warranties and Limitations of Liability.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Bayer CropScience, no claims are made to guarantee germination of carry-over seed.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

### NET CONTENTS:

PRODUCED FOR



Bayer CropScience LP  
800 N. Lindbergh Blvd.  
St. Louis, MO 63167  
1-866-99BAYER (1-866-992-2937)

EverGol Xtend (PENDING) 09/09/2019, 09/10/2019, 05/25/2022, 06/05/2023, 06/07/2023, 06/30/2023, 10/27/2023