



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505T)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

264-1235

Date of Issuance:

3/26/24

NOTICE OF PESTICIDE:

☒ Registration
☐ Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

RGOF0064 Seed Treatment
Insecticide and Fungicide

Name and Address of Registrant (include ZIP Code):

Christie Villegas
Bayer U.S. – Crop Science
700 Chesterfield Pkwy W
Chesterfield, MO 63017

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Continues page 2

Signature of Approving Official:

Venus Eagle, Product Manager 01
Invertebrate-Vertebrate Branch III, Registration Division (7505T)

Date:

3/26/24

2. If, after formal consultation with the appropriate Service(s), additional modifications are identified for any chemical in this product in the Service's Biological Opinion(s) for that chemical, EPA will notify Bayer in writing consistent with the terms in that Biological Opinion of any necessary required changes. Bayer, must submit an application for amendment incorporating any required changes, including amended labels, consistent with the timeline specified in EPA's notification. If Bayer, LLC fails to comply with this term, Bayer has agreed in prior written acceptance on August 24th, 2022, of these terms that EPA may cancel the registration under an expedited process under FIFRA 6(e).
3. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) were provided and are satisfied for HDPE plastic.
4. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 264-1235."
5. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.


If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF(s):

- Basic CSF dated 01/23/2024
- Alternate CSF 1 dated 01/23/2024

If you have any questions, please contact Scott Campbell at campbell.scott@epa.gov

Enclosure



IMIDACLOPRID	GROUP	4A	INSECTICIDE
METALAXYL	GROUP	4	FUNGICIDE
TRIFLOXYSTROBIN	GROUP	11	FUNGICIDE

RGOF0064 Seed Treatment Insecticide and Fungicide

[ABN: GAUCHO OPTI SOYBEAN SEED TREATMENT]

A SEED TREATMENT FOR SOYBEANS TO PROTECT THE SEED AGAINST LISTED PESTS, CONTROL OF SEED ROT, AND DAMPING-OFF DISEASE ON SOYBEAN- FOR COMMERCIAL SEED TREATMENT ONLY

ACTIVE INGREDIENTS:	
Imidacloprid: 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine	26.00%
Metalaxyl: N-(2,6-dimethylphenyl)-N-(methoxyacetyl)alanine methyl ester	6.25%
Trifloxystrobin	2.08%
OTHER INGREDIENTS:	65.67%
TOTAL:	100.00%
Contains 0.626 pounds Metalaxyl per gallon, 0.208 pounds Trifloxystrobin per gallon, 2.604 pounds Imidacloprid per gallon.	

EPA Reg. No. 264-XXXX	EPA Est.
------------------------------	-----------------

KEEP OUT OF REACH OF CHILDREN
CAUTION
Precaución

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

See Back [Side] Panel for First Aid Instructions and Leaflet [Booklet] for Complete Precautionary Statements and Directions for Use.

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• DO NOT induce vomiting unless told to do so by a poison control center or doctor.• DO NOT give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.• Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for treatment advice.

In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577.
Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Workers involved with treating the seed (i.e., operating the treating equipment, loading the chemical, or calibrating the treating equipment); packaging the treated seed (i.e., bagging, closing, tagging, stacking, or forklifting); cleanout of the treatment chamber; or loading and planting treated seeds must wear:

- Long-sleeved shirt and long pants,
- Socks plus shoes, and
- Chemical-resistant gloves made of barrier laminate, butyl rubber (≥14 mils), nitrile rubber (≥14 mils), neoprene rubber (≥14 mils), natural rubber (≥14 mils), polyethylene, polyvinyl chloride (≥14 mils), or Viton (≥14 mils).

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is highly toxic to bees, birds, fish and aquatic invertebrates. For terrestrial uses, **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters. Treated seed exposed on the soil surface may be hazardous to wildlife. Cover or collect seeds spilled during loading.

Ensure that planting equipment is functioning properly in accordance with manufacturing specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.

Groundwater Advisory: This product has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow. Therefore contain any product spills or equipment leaks and dispose of wastes according to the disposal instructions on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all label directions carefully before use.

RGOF0064 Seed Treatment Insecticide and Fungicide is a liquid seed treatment for soybeans to protect the seed against listed pests, control of seed rot, and damping-off disease and is for commercial seed treatment only. Not for use on agricultural establishments in hopper-box, planter-box, slurry-box or other seed treatment applications at or immediately before planting.

Seed treated with RGOF0064 Seed Treatment Insecticide and Fungicide must be visually identifiable from untreated seed by the use of an approved colorant or dye to prevent accidental use of treated seed as food for humans or feed for animals. Refer to 21 CFR, § 2.25. Any colorant or dye added to treated seed must be cleared for use in accordance with 40 CFR, § 153.155(c).

RESISTANCE MANAGEMENT RECOMMENDATION

For resistance management, please note that RGOF0064 Seed Treatment Insecticide and Fungicide contains both a Group 4 and Group 11 fungicide and a Group 4a insecticide. Any fungal or insect population may contain individuals naturally resistant to RGOF0064 Seed Treatment Insecticide and Fungicide and other Group 4 or Group 11 fungicides or Group 4a insecticides. A gradual or total loss of pest control may occur over time if these fungicides/insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide and insecticide resistance, take one or more of the following steps:

- Rotate the use of RGOF0064 Seed Treatment Insecticide and Fungicide or other Group 4 and Group 11 fungicides and Group 4a insecticides within a growing season sequence with different groups that control the same pests.
- Use tank mixtures with fungicides or insecticides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer. **DO NOT** rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the

individual components of the mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):

- Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
- Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
- When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
- Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
- The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated disease management program for fungicide and insecticide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant or pest resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pests.
- For further information or to report suspected resistance contact Bayer CropScience as 1-866-99BAYER (1-866-992-2937).

COMPATIBILITY TESTING AND TANK MIX PARTNERS

This product may be used in combination with other registered Bayer CropScience seed treatment fungicides and insecticides. Pre-test for compatibility with other seed treatment products.

Follow more restrictive labeling of any tank mix partner. **DO NOT** tank mix with any product, which contains a prohibition on tank mixing. RGOF0064 Seed Treatment Insecticide and Fungicide may be applied as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol byproducts that are used in agronomic practice.

Compatibility

It is essential that before using RGOF0064 Seed Treatment Insecticide and Fungicide in any tank mixture the compatibility of the mixture be established. Add RGOF0064 Seed Treatment Insecticide and Fungicide at the labeled rate to a clean quart jar containing approximately one-half the amount of water intended for a final slurry application rate. Next, follow with all other tank mix components that will be used in the total slurry application.

Add last the remaining balance of water. The total amount of volume is determined by the seed size and how much is necessary to ensure complete and uniform coverage and distribution on the seed, as well as the type of commercial seed treating application equipment that will be used.

DO NOT USE MIXTURES THAT CURDLE, PRECIPITATE, OR GEL. USE TANK MIXTURES IMMEDIATELY AFTER MIXING WITH ADEQUATE AGITATION.

Mixing Instructions

1. When applied with seed treatment application equipment designed to treat seed utilizing a treatment slurry mixture, add approximately 1/2 to 2/3 of the required water to the treatment slurry tank followed by the accompanying seed treatment products and RGOF0064 Seed Treatment Insecticide and Fungicide.
2. The order of product addition is dependent on the accompanying seed treatment products and application equipment used.
3. Allow each slurry component to disperse completely prior to the next addition until a uniform suspension is obtained.
4. After all slurry components have been added and dispersed completely add the remaining 1/2 to 1/3 of required water for proper slurry volume and mix until uniform.
5. **DO NOT** store mixed slurries for greater than 72 hours.
6. Maintain adequate agitation until treatment slurry is applied.

For additional instructions for use, slurry component mixing order, and other application suggestions contact your Bayer CropScience representative.

SEED TREATMENT USE DIRECTIONS

	Diseases Controlled	Product Rate	gram ai/100 kg seed	mg ai/seed
Soybean	Aid in control and suppression of seedborne disease and early season seedling disease- Root & Stem Rot- <i>Rhizoctonia solani</i> , <i>Fusarium spp.</i> , <i>Pythium spp.</i> , & <i>Phytophthora</i> Seed Decay- <i>Alternaria spp.</i> , <i>Aspergillus spp.</i> , <i>Cladosporium spp.</i> , <i>Penicillium spp.</i> ,	3.0 fl oz/cwt* (equivalent to 1.5 fl oz/140,000 seed unit)	Imidacloprid – 61.11	Imidacloprid – 0.1012
			Metalaxyl – 14.82	Metalaxyl – 0.0243
			Trifloxystrobin – 4.94	Trifloxystrobin – 0.0081
	Pest Protection			
	Protection from damage caused by: seed corn maggot, soybean aphids, & over-wintering bean leaf beetle		Total – 80.87	Total – 0.134

*Based on 2,800 soybean seeds/lb

Restrictions:

- Seed treated with this product must be visually identifiable from untreated seed by the use of an approved colorant or dye to prevent accidental use of treated seed as food for humans or feed for animals. Refer to 21 CFR, Part 2.25. Any colorant or dye added to treated seed must be cleared for use in accordance with 40 CFR, Part 153.155(c).
- Maximum usage when applying both metalaxyl- and mefenoxam-containing products to the same crop within the same season: **DO NOT** apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop. The maximum application rate for metalaxyl and mefenoxam-containing products (including seed treatments, foliar applications, soil application) is 12.3 lb per acre per calendar year. **DO NOT** apply more than 12.3 lb of active ingredient metalaxyl per acre per calendar year.
- Do not graze or feed livestock on soybean forage or hay.

*Deliver 0.134 mg a.i. per seed based on 2800 soybean seeds per pound. **DO NOT** apply by seed treatment more than 0.0027 lbs of trifloxystrobin, 0.0084 lbs of metalaxyl or 0.0349 lbs of imidacloprid per acre per season.

ROTATIONAL PLANT-BACK INTERVALS*:

Treated areas may be replanted with any crop specified on this label as soon as practical after the last application. For crops for which no tolerances for the active ingredients have been established, a 12-month plant-back interval must be observed.

IMMEDIATE PLANT-BACK: All crops on this label.

30-DAY PLANT-BACK: Barley, bulb vegetables (crops of crop group 03-07), corn (field, pop, and sweet), rapeseed, sorghum, wheat, cereals (including buckwheat, millet, oats, rice, rye, and triticale) and safflower.

12-MONTH PLANT-BACK: All other crops

* Cover crops for soil building or erosion control may be planted any time, but do NOT graze or harvest for food or feed.

NOTIFICATION of the crop rotational restriction must be conveyed to the grower by appropriate seed tag labeling or bag printing on all seed units.

SEED BAG LABELING REQUIREMENT

Federal Seed Act requires that containers containing treated seeds shall be labeled with the following information.

- This seed has been treated with:
 - RGOF0064 Seed Treatment Insecticide and Fungicide which contains 6.25% Metalaxyl, 2.08% Trifloxystrobin and 26.00% Imidacloprid.
- DO NOT** use for feed, food or oil purposes.

The U.S. Environmental Protection Agency requires the following statements on containers containing seed treated with RGOF0064 Seed Treatment Insecticide and Fungicide (264-REGL):

- Store away from feeds and foodstuffs.**
- DO NOT allow children, pets, or livestock to have access to treated seeds.**
- Wear long pants, long-sleeved shirt and chemical resistant gloves when handling treated seed.**
- Treated seeds exposed on the soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during

loading and planting (such as in row ends).

- Dispose of all excess treated seed by burying seed away from bodies of water.
- **DO NOT** contaminate bodies of water when disposing of planting equipment wash water.
- **DO NOT** plant treated seeds in areas subject to seasonal or extended periods of standing water.
- Dispose of seed packaging or containers in accordance with local requirements.
- **DO NOT** graze or feed livestock on soybean forage or hay.
- Excess treated seed may be used for ethanol production if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- Treated seed must be adequately covered with soil at planting (1-1.5 inch depth).
- Ensure that planting equipment is functioning properly in accordance with manufacturing specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.
- **ROTATIONAL PLANT-BACK INTERVALS:** Treated areas may be replanted with soybean as soon as practical after the last application. For crops for which no tolerances for the active ingredients have been established, a 12-month plant-back interval must be observed. **IMMEDIATE PLANT-BACK:** Soybean. **30-DAY PLANT-BACK:** Barley, bulb vegetables (crops of crop group 03-07), corn (field, pop, and sweet), rapeseed, sorghum, wheat, cereals (including buckwheat, millet, oats, rice, rye, and triticale) and safflower. **12-MONTH PLANT-BACK:** All other crops. Cover crops for soil building or erosion control may be planted any time, but do NOT graze or harvest for food or feed
- Pollinator Precautions: Imidacloprid is highly toxic to bees. Ensure that planting equipment is functioning properly in accordance with manufacturer specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.
- Groundwater Advisory: Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.
- Store treated seed in cool, dry conditions.
- When using formulations that do not contain dye, to comply with 40 CFR153.155, all seed treated with an economic poison must be colored with an EPA approved dye such as one of the dyes listed in 40 CFR Sections 180.910 and 180.920 to distinguish and prevent subsequent inadvertent use as a food for man or feed animals.
- After the seeds have been planted, **DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours. Exception: If the treated seed is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, chemical-resistant gloves made of any waterproof material, and chemical-resistant footwear.
- This seed has been treated with 0.099 mg imidacloprid, 0.024 mg metalaxyl, and 0.008 mg trifloxystrobin per seed.
- Maximum usage when applying both metalaxyl- and mefenoxam-containing products to the same crop within the same season: **DO NOT** apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop. The maximum application rate for metalaxyl and mefenoxam-containing products (including seed treatments, foliar applications, soil application) is 12.3 lb per acre per calendar year. **DO NOT** apply more than 12.3 lb of active ingredient metalaxyl per acre per calendar year.
- **DO NOT** apply by seed treatment more than 0.0027 lbs of trifloxystrobin, 0.0084 lbs of metalaxyl or 0.0349 lbs of imidacloprid per acre per season.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE

Store in a cool dry location. **DO NOT** store in direct hot sunlight. **DO NOT** store above 90°F for prolonged periods of time. After prolonged storage in intense cold, allow product to return to approximately 75°F before using.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING

Rigid, Non-refillable containers small enough to shake (i.e., with capacities equal to or less than 5 gallons).

Non-refillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure-rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available, or puncture and dispose of in a sanitary landfill or incineration, or if allowed by state and local authorities, by burning. If burned stay out of smoke.

Rigid Non-refillable containers that are too large to shake (i.e., with capacities greater than 5 gallons or 50 lbs)

Non-refillable Containers

Non-refillable containers - **DO NOT** reuse or refill this container. After emptying product from container, either return container to Bayer CropScience per instructions from Bayer CropScience Customer Service Center (1-800-527-4781), or rinse and either recycle or dispose of the container as follows:

Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Top Discharge IBC, Drums, Kegs (e.g.– Snyder 120 Next Gen, Bonar B120, Drums, and Kegs)

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. To triple rinse the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

Refillable Containers

Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. **DO NOT** remove tamper evident cables or one-way valves or clean the container if the container is equipped with one-way valves and refilling or returning is planned. After emptying product from container, either return container to Bayer CropScience per instructions from Bayer CropScience Customer Service Center (1-800-527-4781) or rinse and either recycle or dispose of the container as follows:

Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Top Discharge IBC, Drums, Kegs (e.g. – Snyder 120 Next Gen, Bonar B120, Drums, and Kegs)

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To triple rinse the containers before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

Refillable Containers

End users are authorized to remove tamper evident cables as required to remove the product from the container unless the container is equipped with one-way valves and refilling or returning is planned. If this is the case, end users are not authorized to remove tamper evident cables, one-way valves or clean container. See Container Disposal instructions under Storage and Disposal.

IMPORTANT: READ BEFORE USE

CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Bayer CropScience, no claims are made to guarantee germination of carry-over seed.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

NET CONTENTS:

PRODUCED FOR



Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167
1-866-99BAYER (1-866-992-2937)

RGOF0064 Seed Treatment Insecticide and Fungicide (MASTER) 3/26/24