



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505T)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

264-1239

Date of Issuance:

10/10/24

NOTICE OF PESTICIDE:

☒ Registration
☐ Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Acceleron 26FI Fungicide and
Insecticide Seed Treatment

Name and Address of Registrant (include ZIP Code):

Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO, 63167

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Continues page 2

Signature of Approving Official:

Venus Eagle, Product Manager 01
Invertebrate & Vertebrate Branch 3,
Registration Division (7505T)

Date:

10/10/24

2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 264-1239."
3. If, after formal consultation with the appropriate Service(s), additional modifications are identified in the Service's Biological Opinion(s) for imidacloprid, EPA will notify Bayer CropScience LP in writing consistent with the terms in the Biological Opinion of any necessary required changes. Bayer CropScience LP must submit an application for amendment incorporating any required changes, including amended labels, consistent with the timeline specified in EPA's notification. If Bayer CropScience LP fails to comply with this term, Bayer CropScience LP has agreed in prior written acceptance on June 12, 2024, that EPA may cancel the registration under an expedited process under FIFRA 6(e).
4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF(s):

- Basic CSF dated 08/24/2023

If you have any questions, please contact Ralph Narain at 202-566-2853 or at Narain.Ralph@epa.gov.

Enclosure: Stamped label

IMIDACLOPRID	GROUP	4A	INSECTICIDE
METALAXYL	GROUP	4	FUNGICIDE
PROTHIOCONAZOLE	GROUP	3	FUNGICIDE
FLUOXASTROBIN	GROUP	11	FUNGICIDE

Acceleron 26FI Fungicide and Insecticide

Seed Treatment

[ABN: Acceleron Prime Fungicide Insecticide]

A SEED TREATMENT PRODUCT FOR CONTROL OF SEED ROT AND DAMPING-OFF DISEASES ON SOYBEAN – FOR COMMERCIAL SEED TREATMENT ONLY


ACTIVE INGREDIENTS:

Imidacloprid: 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine	22.41%
Metalaxyl: N-(2,6-dimethylphenyl)-N-(methoxyacetyl)alanine methyl ester.....	4.31%
Prothioconazole: 2-[2-(1-Chlorocyclopropyl)-3-(2-chlorophenyl)-2-hydroxypropyl]- 1,2- dihydro-3H-1,2,4-triazole-3-thione.....	2.16%
Fluoxastrobin: ((2E)-(2-[6-(2-chlorophenoxy)-5-fluoro-4pyrimidinyl] oxy phenyl)-(5, 6-dihydro-1, 4, 2-dioxazin-3yl) methanone O-methyloxime).....	2.16%

OTHER INGREDIENTS:

TOTAL:

Contains 2.17 pounds Imidacloprid per gallon (260 grams ai/liter)
 Contains 0.42 pounds Metalaxyl per gallon (50 grams ai/liter)
 Contains 0.21 pounds Prothioconazole per gallon (25 grams ai/liter)
 Contains 0.21 pounds Fluoxastrobin per gallon (25 grams ai/liter)



68.96%
100.00%

EPA Reg. No. 264-XXXX

EPA Est.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
 For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

Please refer to [back panel] [booklet] for additional precautionary statements and directions for use. [Note to reviewer: Location of additional precautionary statements and directions for use will vary between those listed, depending on container type/size.]

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.

**In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577.
Have a product container or label with you when calling a poison control center or doctor, or going for treatment.**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Causes Moderate eye irritation. Avoid contact with skin, eyes, or clothing. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Workers involved with treating the seed (i.e., operating the treating equipment, loading the chemical, or calibrating the treating equipment); packaging the treated seed (i.e., bagging, closing, tagging, stacking, or forklifting); cleanout of the treatment chamber; or loading and planting treated seeds must wear:

- Long-sleeved shirt and long pants
- Socks plus shoes
- Chemical-resistant gloves made of barrier laminate, butyl rubber (≥ 14 mils), nitrile rubber (≥ 14 mils), neoprene rubber (≥ 14 mils), polyethylene, polyvinyl chloride (≥ 14 mils), or Viton (≥ 14 mils)

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Fluoxastrobin is toxic to fish and aquatic invertebrates. Prothioconazole-desthio and prothioconazole-S-methyl degradates of prothioconazole, are toxic to aquatic invertebrates. Imidacloprid is toxic to bees, birds, fish, and aquatic invertebrates. Runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Imidacloprid is highly toxic to bees exposed to direct treatment and effects may be possible as a result of exposure to translocated residues on blooming crops or weeds. Ensure that planting equipment is functioning properly in accordance with manufacturing specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds. The active ingredients in this product can be persistent for several months or longer. For terrestrial uses, do not apply to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect seed spilled during loading.

Surfacewater Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this product from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Groundwater Advisory

Imidacloprid, metalaxyl, prothioconazole-desthio (a degradate of prothioconazole), and fluoxastrobin are known to leach through soil into groundwater under certain conditions as a result of agricultural use. This product may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Read the entire label before using this product.

PRODUCT INFORMATION

Acceleron 26FI Fungicide and Insecticide Seed Treatment is a systemic fungicide and insecticide seed treatment specifically for control and suppression of seedborne disease and early season seedling disease of root and stem rot caused by *Rhizoctonia solani*, *Fusarium spp.*, *Pythium spp.*, and *Phytophthora spp.*, and Seed Decay caused by, *Fusarium spp.*, *Aspergillus spp.*, *Penicillium spp.*, and *Phomopsis spp.*, and protection from damage caused by seed corn maggot, early-season soybean aphids, and over-wintering bean leaf beetle.

USE RESTRICTIONS

- For use only in commercial seed treatment facilities. Not for use in agricultural establishments in hopper-box, planter-box, slurry-box, or other seed treatment applications at or immediately before planting.
- Seed treated with this product must be visually identifiable from untreated seed by the use of an approved colorant or dye to prevent accidental use of treated seed as food for humans or feed for animals. Refer to 21 CFR, Part 2.25. Any colorant or dye added to treated seed must be cleared for use in accordance with 40 CFR, Part 153.155(c).
- Maximum usage when applying metalaxyl- and mefenoxam-containing products, imidacloprid-containing products, fluoxastrobin-containing products and prothioconazole-containing products to the same crop within the same season: **DO NOT** apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop. The maximum application rates of Acceleron 26FI are located in the CROP USE DIRECTIONS section of this label.
- **DO NOT** graze or feed livestock on soybean forage or hay
- **DO NOT** allow children, pets, or livestock to have access to treated seed.
- **DO NOT** use treated seed for food, feed, or oil.
- Not for Sale, Sale Into, Distribution and /or Use in Nassau and Suffolk Counties of New York State.
- Refer to the specific use directions and restrictions in the crop table.
- Store treated seed away from feeds and foodstuffs.
- Plant treated seed into the soil at a minimum depth of 1/2 inch or greater to minimize exposure.

ROTATIONAL CROPS*

In the event of a crop failure or harvest of a crop grown from seed treated with Acceleron 26FI Fungicide and Insecticide, the field may be replanted immediately with any crops, except those with the 30-DAY MINIMUM PLANTBACK: Barley, bulb vegetables (crops of crop group 03-07), corn (field, pop, and sweet), rapeseed, sorghum, wheat, cereals (including buckwheat, millet, oats, rice, rye, and triticale) and safflower. After treating with other products, **DO NOT** rotate any crops except soybeans until after the longest plant-back interval for the active ingredient, as specified on the product label containing that active ingredient.

* Cover crops for soil building or erosion control may be planted any time, but do NOT graze or harvest for food or feed, unless a crop tolerance exists.

NOTIFICATION of the crop rotational restriction must be conveyed to the grower by appropriate seed tag labeling or bag printing on all seed units.

RESISTANCE MANAGEMENT RECOMMENDATION

For resistance management, please note that Acceleron 26FI Fungicide and Insecticide Seed Treatment contains Group 3, 4, and Group 11 fungicides and a Group 4A insecticide. Any fungal or insect population may contain individuals naturally resistant to Acceleron 26FI Fungicide and Insecticide Seed Treatment and other Group 3, 4 or Group 11 fungicides or Group 4A insecticides. A gradual or total loss of pest control may occur over time if these fungicides/insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide and insecticide resistance, take one or more of the following steps:

- Rotate the use of Acceleron 26FI Fungicide and Insecticide Seed Treatment or other Group 3, 4, and Group 11 fungicides and Group 4A insecticides within a growing season sequence with different groups that control the same pests.
- Use tank mixtures with fungicides or insecticides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer. **DO NOT** rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of the mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).

- Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
- The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated disease management program for fungicide and insecticide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant or pest resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pests.
- For further information or to report suspected resistance contact Bayer CropScience as 1-866-99BAYER (1-866-992-2937).

APPLICATION INSTRUCTIONS

- Uniform application of seed is necessary to ensure seed safety and best disease protection.
- Dilute with a sufficient amount of water to achieve uniform application to seed which is necessary to ensure seed safety and best disease control.
- Use high quality, properly cleaned seed.

COMPATIBILITY TESTING AND TANK MIX PARTNERS

May be used in combination with other registered Bayer CropScience fungicide seed treatments and insecticides. Pre-test for compatibility with other seed treatment products.

Compatibility

It is essential that before using Acceleron 26FI Fungicide and Insecticide Seed Treatment in any tank mixture the compatibility of the mixture be established. Add Acceleron 26FI Fungicide and Insecticide Seed Treatment at the labeled rate to a clean quart jar containing approximately one-half the amount of water intended for a final slurry application rate. Next, follow with all other tank mix components that will be used in the total slurry application.

Add last the remaining balance of water. The total amount of volume is determined by the seed size and how much is necessary to ensure complete and uniform coverage and distribution on the seed, as well as the type of commercial seed treating application equipment that will be used.

DO NOT USE MIXTURES THAT CURDLE, PRECIPITATE, OR GEL. USE TANK MIXTURES IMMEDIATELY AFTER MIXING WITH ADEQUATE AGITATION.

Mixing Instructions

1. When applied with seed treatment application equipment designed to treat seed utilizing a treatment slurry mixture, add approximately 1/2 to 2/3 of the required water to the treatment slurry tank followed by the accompanying seed treatment products and Acceleron 26FI Fungicide and Insecticide Seed Treatment.
2. The order of product addition is dependent on the accompanying seed treatment products and application equipment used.
3. Allow each slurry component to disperse completely prior to the next addition until a uniform suspension is obtained.
4. After all slurry components have been added and dispersed completely add the remaining 1/2 to 1/3 of required water for proper slurry volume and mix until uniform. For containers over 15 gallons, mix the product with the built-in mixer for 5-10 minutes before each use.
5. **DO NOT** store mixed slurries for greater than 72 hours. Always recirculate or mix thoroughly before using.
6. Maintain adequate agitation until treatment slurry is applied.

For additional instructions for mixing, use, slurry component mixing order, and other application suggestions contact your Bayer CropScience representative.

Follow the most restrictive labeling of any tank mix partner. **DO NOT** tank mix with any product, which contains a prohibition on tank mixing. Acceleron 26FI Fungicide and Insecticide Seed Treatment may be applied as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

CROP USE DIRECTIONS

	Diseases Controlled	Product Rate	gram ai/100 kg seed	mg ai/seed
Soybean	Aid in control and suppression of seedborne disease and early season seedling disease- Root & Stem Rot- <i>Rhizoctonia solani</i> , <i>Fusarium spp.</i> , <i>Pythium spp.</i> , and <i>Phytophthora spp.</i> Seed Decay- <i>Fusarium spp.</i> , <i>Aspergillus spp.</i> , <i>Penicillium spp.</i> , and <i>Phomopsis spp.</i>	4.6 fl oz/100 lbs seed (2.3 fl oz/140,000 seed unit*)	Imidacloprid - 78	Imidacloprid – 0.126
			Metalaxyl - 15	Metalaxyl - 0.025
			Fluoxastrobin - 7.5	Fluoxastrobin – 0.012
	Pest Controlled		Prothioconazole - 7.5	Prothioconazole – 0.012
	Protection from damage caused by: seed corn maggot, early-season soybean aphids, & over-wintering bean leaf beetle		Total - 108	Total - 0.175

* Based on 2,800 soybean seeds/lb

For soybeans with average size of 2,800 soybean seed/lb and average seeding rate of 140,000 seed/a the recommended rate of 4.6 fl oz/100 lb seed will result in: 0.039 lb/acre of imidacloprid, 0.0075 lb/acre of metalaxyl, 0.0037 lb/a of fluoxastrobin, and 0.0037 lb/a of prothioconazole

SEED BAG LABEL REQUIREMENTS

Federal Seed Act requires that containers containing treated seeds shall be labeled with the following information.

- This seed has been treated with:
 - Acceleron 26FI Fungicide and Insecticide Seed Treatment (EPA Reg. No. 264-REGO) which contains 22.41% Imidacloprid, 4.31% Metalaxyl, 2.16% Prothioconazole, and 2.16% Fluoxastrobin.
- DO NOT** use for feed, food or oil purposes.

The U.S. Environmental Protection Agency requires the following statements on containers containing seed treated with Acceleron 26FI Fungicide and Insecticide Seed Treatment (EPA Reg. No. 264-REGO):

- In the event of a crop failure or harvest of a crop grown from seed treated with Acceleron 26FI Fungicide and insecticide, the field may be replanted immediately with any crops. After treating with other products, do not rotate any crops except soybeans until after the longest plant-back interval for the active ingredient, as specified on the product label containing that active ingredient. Cover crops for soil building or erosion control may be planted any time, but do NOT graze or harvest for food or feed, unless a crop tolerance exists.
- Store treated seed away from feeds and foodstuffs.**
- DO NOT allow children, pets, or livestock to have access to treated seeds.**
- When opening this bag or loading/pouring the treated seed, wear long sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves.**
- Treated seeds exposed on the soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting (such as in row ends).
- Dispose of all excess treated seed by burying seed away from bodies of water.
- DO NOT** contaminate bodies of water when disposing of planting equipment wash water.
- DO NOT** plant treated seeds in areas subject to seasonal or extended periods of standing water.
- Dispose of seed packaging or containers in accordance with local requirements.
- DO NOT** use empty seed bags for any other purpose
- Treated seed must be adequately covered with soil at planting.
- Imidacloprid is highly toxic to bees exposed to direct treatment and effects may be possible as a result of exposure to translocated residues on blooming crops or weeds. Ensure that planting equipment is functioning properly in accordance with

manufacturing specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.

- Plant treated seed into the soil at a minimum depth of 1/2 inch or greater to minimize exposure.
- **DO NOT** plant treated seed by broadcasting to the soil surface. Ensure that all planted seeds are thoroughly incorporated by the planter during planting, additional incorporation may be required to thoroughly cover exposed seeds.
- Seed treated with this product must be visually identifiable from untreated seed by the use of an approved colorant or dye to prevent accidental use of treated seed as food for humans or feed for animals. Refer to 21 CFR, Part 2.25. Any colorant or dye added to treated seed must be cleared for use in accordance with 40 CFR, Part 153.155(c).
- Maximum usage when applying metalaxyl- and mefenoxam-containing products, imidacloprid-containing products, fluoxastrobin-containing products and prothioconazole-containing products to the same crop within the same season: **DO NOT** apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop.
- To reduce seed dust which can drift onto blooming crops or weeds, ensure that planting equipment is functioning properly in accordance with manufacturer's recommendations. Surplus seed or empty seed containers should be stored or disposed according to state or local regulations.
- This seed has been treated with 0.078 lbs. imidacloprid, 0.015 lbs. metalaxyl, 0.0075 lbs. fluoxastrobin, and 0.0075 lbs. prothioconazole per 100 lbs. of seed (based on 2800 soybean seed/lb, 0.126 mg. imidacloprid, 0.025 mg. metalaxyl, 0.012 mg. fluoxastrobin, and 0.012 mg. prothioconazole per seed)

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE

Store in a cool dry location. **DO NOT** store in direct hot sunlight. **DO NOT** store above 90°F for prolonged periods of time. After prolonged storage in intense cold, allow product to return to approximately 75°F before using.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING

Rigid, Non-refillable containers small enough to shake (i.e., with capacities equal to or less than 5 gallons).

Non-refillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure-rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available, or puncture and dispose of in a sanitary landfill or incineration, or if allowed by state and local authorities, by burning. If burned stay out of smoke.

Rigid Non-refillable containers that are too large to shake (i.e., with capacities greater than 5 gallons)

Non-refillable Containers

Non-refillable containers - **DO NOT** reuse or refill this container. After emptying product from container, either return container to Bayer CropScience per instructions from Bayer CropScience Customer Service Center (1-800-527-4781), or rinse and either recycle or dispose of the container as follows:

Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Top Discharge IBC, Drums, Kegs (e.g.– Snyder 120 Next Gen, Bonar B120, Drums, and Kegs)

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. To triple rinse the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the

container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

Refillable Containers

Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. **DO NOT** remove tamper evident cables or one-way valves or clean the container if the container is equipped with one-way valves and refilling or returning is planned. After emptying product from container, either return container to Bayer CropScience per instructions from Bayer CropScience Customer Service Center (1-800-527-4781) or rinse and either recycle or dispose of the container as follows:

Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Top Discharge IBC, Drums, Kegs (e.g. – Snyder 120 Next Gen, Bonar B120, Drums, and Kegs)

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To triple rinse the containers before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

Refillable Containers

End users are authorized to remove tamper evident cables as required to remove the product from the container unless the container is equipped with one-way valves and refilling or returning is planned. If this is the case, end users are not authorized to remove tamper evident cables, one-way valves or clean container. See Container Disposal instructions under Storage and Disposal.

IMPORTANT: READ BEFORE USE

CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Bayer CropScience, no claims are made to guarantee germination of carry-over seed.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

NET CONTENTS:

PRODUCED FOR



Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167
1-866-99BAYER (1-866-992-2937)

Acceleron 26FI Fungicide and Insecticide Seed Treatment (PENDING) 9/15/2023, 09/11/2024, 10/01/2024