

## DIRECTIONS

Apply as a water-base spray at dilutions recommended below, preferably using power sprayer with mechanical agitation. When using low-pressure or hand sprayer, stir spray solution thoroughly, and agitate manually throughout entire spraying operation. Spray animals until thoroughly wet. Average size animal requires approximately 2 quarts of spray. Always stir well before using.

**FOR CATTLE** . . . To control TICKS (Lone Star, Gulf Coast, Winter and Fever Ticks) . . . LICE (Short Nose, Long Nose and Tail Lice) . . . also HORN flies, fleas, gnats and mosquitoes on Cattle — apply at dilution of 1 to 150. To control Gulf Coast Tick, treatment of ears and around ears is essential, repeating treatment in 10 to 14 days. **FOR MANGE MITES** (Sarcoptic, Psoroptic or Chorioptic, apply at a dilution of 1 to 90, repeating at 7 to 10 days. **FOR SHEEP AND GOATS** . . . To control SHEEP TICKS (Keds) and Lice on Sheep and Goats, apply at a dilution of 1 to 600 and wet thoroughly, repeating treatment in 10 to 14 days. **FOR SHEEP SCAB**, apply at a dilution of 1 to 150 repeating treatment at 7 to 10 days. **FOR FLEECE WORMS** — dilute 1 to 150 and wet thoroughly repeating treatment in 10 to 14 days. When used in a dipping vat, immerse sheep not less than 12 minute, ducking head at least twice. Dip solution should be checked frequently by vat-side test to assure proper dipping strength.

**CAUTION:** Do not use as spray on any animal within 30 days of slaughter nor as a dip within 60 days of slaughter.

For Use in **CATTLE OILERS** Back Rubbers

To control Horn Flies and aid in the control of lice on range and beef cattle, dilute one part with 19 parts of fuel oil or mineral seal oil and mix thoroughly. One quart makes 3 gallons finished oiler solution.

**CAUTION:** Calves less than six months old should not be exposed to back rubber treatment. Do not permit

animals access to backrubber within 30 days of slaughter.

**CAUTION:** Follow directions explicitly. Do not exceed recommended dilutions. Do not use on dairy cattle, dairy goats, or suckling calves, inside dairy barns, around feed or forage for animals. Do not use in milk rooms or poultry houses. Do not use in combination with arsenical sprays. Do not apply to animals under 6 months of age. Do not treat animals in cold stormy weather. Do not treat overheated or sick animals. Do not use on humans, household pets, dogs, cats or poultry. Make sure you have a perfect mixture in spray solution. If oily layer or particles form, do NOT apply spray to animals. Do not reuse empty container. Destroy by perforation or crushing. Bury or discard in safe place.

**CAUTION:** DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

**WARNING:** This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

**WARNING:** Keep out of reach of children.

**WARNING:** May Be Fatal If Swallowed! May be absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after use. Do not store near feed and food products.

**ANTIDOTE:** Human - Internal - If swallowed, give one tablespoonful of salt in a glass of warm water. Repeat until vomit is clear. Call doctor immediately. External - In case of contact, wash immediately with plenty of soap and water.



EPA Reg. No. 270-18

FARNAM

# TOXANOX

**PLUS Emulsifiable  
Livestock Spray**

An emulsifiable livestock spray, specially formulated for use on cattle, sheep and goats. Combines the quick kill characteristics of Lindane with the positive killing power of Toxaphene. Mixes with water at most economical dilutions.

**Controls Horn Flies, House Flies, Stable Flies, Ticks, Lice and Mange Mites (Sarcoptic, Psoroptic and Chorioptic) on cattle, sheep and goats. Also Sheep Ticks (Keds), Sheep Scab and Fleece Worms!**

Active Ingredients: \*Toxaphene 40.5%, Gamma Isomer of Benzene Hexachloride from Lindane 1.80%, Deodorized Kerosene 47.2%  
Inert Ingredients: 10.5%

\*Technical Chlorinated Camphene (Chlorine content 67-69%)

**WARNING!** Keep out of reach of children.

(See Other Precautions On Back Panel)

**32 FL. OZ. (1 QUART)**

8701 No. 29th St.

PRINTED IN U.S.A.

Manufactured Exclusively For . . .

**THE FARNAM CO.**



**OMAHA, NEBRASKA**

## DIRECTIONS

Apply as a water base spray at dilutions recommended below, preferably using power sprayer with mechanical agitation. When using low pressure or hand sprayer, stir spray solution thoroughly, and agitate manually throughout entire spraying operation. Spray animals until thoroughly wet. Average size animal requires approximately 2 quarts of spray. Always stir well before using.

**FOR CATTLE** . . . To control TICKS: Lone Star, Gulf Coast, Winter and Fever Ticks . . . LICE (Short Nose, Long Nose and Tail Lice) . . . also HORN flies, fleas, gnats and mosquitoes on Cattle — apply at dilution of 1 to 150. To control Gulf Coast Tick, treatment of ears and around ears is essential, repeating treatment in 10 to 14 days. **FOR MANGE MITES** Sarcoptic, Psoroptic or Chorioptic, apply at a dilution of 1 to 90, repeating at 7 to 10 days. **FOR SHEEP AND GOATS** . . . To control SHEEP TICKS, Keds and Lice on Sheep and Goats, apply at a dilution of 1 to 600 and wet thoroughly, repeating treatment in 10 to 14 days. **FOR SHEEP SCAB**, apply at a dilution of 1 to 150 repeating treatment at 7 to 10 days. **FOR FLEECE WORMS** — dilute 1 to 150 and wet thoroughly repeating treatment in 10 to 14 days. When used in a dipping vat, immerse sheep not less than 12 minute, ducking head at least twice. Dip solution should be checked frequently by vatside test to assure proper dipping strength.

**CAUTION** Do not use as spray on any animal within 30 days of slaughter nor as a dip within 60 days of slaughter.

For Use in **CATTLE OILERS** Back Rubbers

To control Horn Flies and aid in the control of lice on range and beef cattle dilute one part with 19 parts of fuel oil or mineral seal oil and mix thoroughly. One quart makes 5 gallons finished oiler solution.

**CAUTION** Calves less than six months old should not be exposed to back rubber treatment. Do not permit

animals access to backrubber within 30 days of slaughter.

**CAUTION** Follow directions explicitly. Do not exceed recommended dilutions. Do not use on dairy cattle, dairy goats, or suckling calves, inside dairy barns, around feed or forage for animals. Do not use in milk rooms or poultry houses. Do not use in combination with arsenical sprays. Do not apply to animals under 6 months of age. Do not treat animals in cold stormy weather. Do not treat overheated or sick animals. Do not use on humans, household pets, dogs, cats or poultry. Make sure you have a perfect mixture in spray solution. If oily layer or particles form, do NOT apply spray to animals. Do not reuse empty container. Destroy by perforation or crushing. Bury or discard in safe place.

**CAUTION: DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME**

**WARNING:** This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

**WARNING:** Keep out of reach of children.

**WARNING:** May Be Fatal If Swallowed! May be absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after use. Do not store near feed and food products.

**ANTIDOTE:** Human: Internal — If swallowed, give one tablespoonful of salt in a glass of warm water. Repeat until vomit is clear. Call doctor immediately. External — In case of contact, wash immediately with plenty of soap and water.



EPA Reg. No. 270-18

**FARNAM**

# TOXANOX

**PLUS Emulsifiable  
Livestock Spray**

An emulsifiable livestock spray, specially formulated for use on cattle, sheep and goats. Combines the quick kill characteristic of Lindane with the positive killing power of Toxaphene. Mixes with water at most economical dilutions.

**Controls Horn Flies, House Flies, Stable Flies, Ticks, Lice and Mange Mites (Sarcoptic, Psoroptic and Chorioptic) on cattle, sheep and goats. Also Sheep Ticks (Keds), Sheep Scab and Fleece Worms!**

Active Ingredients: \*Toxaphene 40.5%, Gamma Isomer of Benzene Hexachloride from Lindane 1.80%, Deodorized Kerosene 47.2%  
Inert Ingredients: 10.5%

\*Technical Chlorinated Camphene (Chlorine content 67-69%)

**WARNING!** Keep out of reach of children.

(See Other Precautions On Back Panel)

**32 FL. OZ. (1 QUART)**

8701 No. 29th St.

PRINTED IN U.S.A.

Manufactured Exclusively For

**THE FARNAM CO. OMAHA, NEBRASKA**

