

100-1000

(FRONT PANEL)

FARNAM  
3%  
CLODRIN INSECTICIDE\*  
DUST

Livestock Insecticide Powder  
For Automatic Dusting Stations

FOR DAIRY CATTLE--To Control Hornflies and Lice  
FOR BEEF CATTLE--To Control Hornflies

3% CLODRIN INSECTICIDE DUST is a specially prepared livestock dusting powder ideal for use in automatic dusting stations. Also for use in conventional hand and power dusters. No feed-off period is required when used on animals being fed for slaughter.

Active Ingredients

Dimethyl Phosphate of alpha-Methylbenzyl	
2-hydroxy-2-isobutanoic (Clover).....	3.0%
Inert ingredients.....	97.0%
	TOTAL 100.0%

CAUTION: Keep out of reach of children.  
(See label for additional caution)

\*Registered Trademark of Farnam Chemical Company

Net Wt. \_\_\_\_\_

THE FARNAM COMPANY

OMAHA, NEBRASKA

ACCEPTED

59

FARNAM


# RONENOX

EMULSIFIABLE LIVESTOCK SPRAY

For Hornflies, Lice and Screw Worms on Beef Cattle and Non-Lactating Dairy Cattle. Dilute one part of RonenoX with 100 parts of water. Spray liberally on the head, neck, and body of the animal. Repeat every 7 to 10 days. For Lope Star and Winter Ticks on Beef Cattle and Non-Lactating Dairy Cattle. Dilute one part of RonenoX with 100 parts of water. Spray liberally on the head, neck, and body of the animal. Repeat every 7 to 10 days. For Wool Maggots, Screw Worms, Hornflies on Sheep and Non-Lactating Goats. Dilute one part of RonenoX with 100 parts of water. Spray liberally on the head, neck, and body of the animal. Repeat every 7 to 10 days. For Dog and Sheep Ticks on Sheep and Non-Lactating Goats. Dilute one part of RonenoX with 100 parts of water. Spray liberally on the head, neck, and body of the animal. Repeat every 7 to 10 days. For Fleas on Swine. Dilute one part of RonenoX with 100 parts of water. Spray liberally on the head, neck, and body of the animal. Repeat every 7 to 10 days. For Fleas on Cattle, Horses, and Dogs. Dilute one part of RonenoX with 100 parts of water. Spray liberally on the head, neck, and body of the animal. Repeat every 7 to 10 days.

NEBRASKA

NEBRASKA



ONE FLUID GALLON

CAUTION: Keep Out of Reach of Children

THE FARNAM CO., 8701 North 29th Street, OMAHA, NEBRASKA

## DIRECTIONS

For Hornflies, Lice and Screw Worms on Beef Cattle and Non-Lactating Dairy Cattle. Dilute one part of RonenoX with 100 parts of water. Spray liberally on the head, neck, and body of the animal. Repeat every 7 to 10 days.

For Lope Star and Winter Ticks on Beef Cattle and Non-Lactating Dairy Cattle. Dilute one part of RonenoX with 100 parts of water. Spray liberally on the head, neck, and body of the animal. Repeat every 7 to 10 days.

For Wool Maggots, Screw Worms, Hornflies on Sheep and Non-Lactating Goats. Dilute one part of RonenoX with 100 parts of water. Spray liberally on the head, neck, and body of the animal. Repeat every 7 to 10 days.

For Dog and Sheep Ticks on Sheep and Non-Lactating Goats. Dilute one part of RonenoX with 100 parts of water. Spray liberally on the head, neck, and body of the animal. Repeat every 7 to 10 days.

For Fleas on Swine. Dilute one part of RonenoX with 100 parts of water. Spray liberally on the head, neck, and body of the animal. Repeat every 7 to 10 days.

For Fleas on Cattle, Horses, and Dogs. Dilute one part of RonenoX with 100 parts of water. Spray liberally on the head, neck, and body of the animal. Repeat every 7 to 10 days.