

[.270.60]



*Contains
Trichlorfon*

WARNING:
Keep out of Reach of
Children. See Back Panel
for Warning Statements.

NET CONTENTS: 20 GRAMS

Manufactured Exclusively For

FARNAM COMPANIES · OMAHA, NEBRASKA

**For the Removal of Bots (*Gasterophilus spp.*), ascarids
(*Parascaris equorum*) and pinworms (*Oxyuris equi*) from horses.**

DOSAGE

Give one packet (10 grams) for the average horse and mature horse. For smaller breeds and younger horses, administer one or more packets (5 grams) for each 250 pounds of body weight. For most effective control, treatment should be repeated at 4 to 6 month intervals. BOT-X Horse Wormer can be used at any time, but do not repeat more frequently than every 30 days.

DIRECTIONS

Use required dosage dry or dissolved in water and mix with the amount of feed to be consumed at one feeding.

Withdrawal of feed 12 to 18 hours prior to treatment will assure complete ingestion. Allow horse to consume all of the medicated feed before additional feed is given. Remove and destroy any medicated feed not consumed within 12 hours.

ACTIVE INGREDIENT:

Dimethyl 2,2,2 trichloro 1-hydroxyethyl phosphonate 90%

INERT INGREDIENTS:

(Colloidal Silica and Dextrose) 10%

U.S. Patent No. 2,791,295 Canadian Patent No. 529,519 100%
†Trichlorfon

RESTRICTIONS

Use only as directed. Do not treat horses to be used for food. Do not treat sick or debilitated horses or foals under 4 months of age, mares in the last month of pregnancy or animals other than horses. Do not exceed recommended dose. Warning: BOT-X is a cholinesterase inhibitor. Do not use this product on animals immediately or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals. Do not administer intravenously or anesthetics, especially muscle relaxants, for a period of 2 weeks after using this product.

NOTICE TO VETERINARIANS

Symptoms of overdosage in horses are ataxia and depression. Atropine is antidote. Give 100 milligrams or 2 milligrams per 100 pounds of body weight.

WARNING

Not for human use. Keep out of reach of children. Do not get in eyes or on skin. Wash hands with soap and water after use. If swallowed by humans, induce vomiting by giving 1 table spoon of 3% hydrogen peroxide solution of water. Call physician immediately.

TO PHYSICIAN

This material is a cholinesterase inhibitor. Atropine and 2-PAM are antidotes. Give 100 mg for treatment. Do not wash stomach or induce vomiting.

USDA REG. NO. 270-60

FARNAM COMPANIES

801 North 19th Street, Omaha, Nebraska 68102

FARNAM COMPANIES · OMAHA, NEBRASKA

NET CONTENTS, 20 GRAMS