

CAUTION

Baits must be placed in areas inaccessible to children, pets, wildlife and domestic animals, or in tamper proof bait boxes. Do not contaminate water by cleaning of equipment or disposal of wastes. Keep out of lakes, streams, or ponds. Apply this product only as specified on this label.

**DO NOT REUSE CONTAINER
DESTROY WHEN EMPTY**

If swallowed by humans, domestic animals, wildlife or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated, as in the case of hemorrhage caused by overdose of Bishydroxycoumarin. In case of ingestion, call a Physician.


***TOX-HID Wisconsin Alumni Research Foundation Patent Pending
TOX-HID Trademark of Wisconsin Alumni Research Foundation**

E.P.A. REG. NO. 270-109

E.P.A. EST. NO. 6658-MO-1

Manufactured Exclusively For
FARNAM COMPANIES
Omaha, Nebraska U.S.A. 68112

PRINTED IN U.S.A.



FARNAM

Kills Rats and Mice

Pelleted Rat Bait


A TOX-HID* COATED WARFARIN PELLETTED BAIT

Each particle of the warfarin in this product has been specially coated with a material that reduces detection of the warfarin by the rodents at the time the bait is consumed. The coating may help assure greater acceptance of the bait in cases where rodent awareness of uncoated warfarin is responsible for low bait acceptability.

ACTIVE INGREDIENTS:

Warfarin (3-Alpha-Acetylbenzyl)-4-Hydroxycoumarin) 0.025%

INERT INGREDIENTS: 99.975%



DIRECTIONS

NORWAY AND ROOF RATS: Open package and place four to six ounces of Farnam Pelleted Bait in dry container next to burrows, runways, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than fifteen days and continue baiting until all signs of feeding have stopped. Replace stale, damp, or moldy baits with fresh bait.

HOUSE MICE: Open package and place table-spoonful amounts (1/4 to 1/2 ounce) of Farnam Pelleted Bait in shallow containers where mice or signs of mice have been seen. Bait stations at intervals of eight to twelve feet are recommended. Assure an uninterrupted supply of bait for a period of not less than fifteen days.

This rodenticide contains warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and, upon repeated feedings causes fatal hemorrhage in rats and mice.

CAUTION: Keep Out of Reach of Children

See back panel for additional cautions

NET WT. 16 OZ. (1 LB.)

FARNAM COMPANIES • OMAHA, NEBRASKA, U.S.A.

