# ATTACHMENT TO CONFIDENTIAL STATEMENT OF FORMULA

Date:

10/24/89

Product Name: FARNAM GO-FLY

EPA Registration Number:

270-145

Registrant:

Farnam Companies, Inc. 301 West Osborn Road Phoenix, Arizona 85013

COMPONENT:

Pyrethrin 20%

SUPPLIERS:

Fairfield American 238 Wilson Avenue

Newark, NJ 07015

MGK

EPA REG. NO.: 1021-24

EPA REG. NO.: 4816-17

8810 Tenth Avenue North Minneapolis, MN 55427

Prentiss

EPA REG. NO.:

655-75

C.B. 2000

Floral Park, NY 11001

COMPONENT:

Piperonyl Butoxide Tech.

SUPPLIERS:

Fairfield American 238 Wilson Avenue Newark, NJ 07015

EPA REG. NO.: 4816-72

MGK

8810 Tenth Avenue North

Minneapolis, MN 55427

Prentiss

C.B. 2000

Floral Park, NY 11001

EPA REG. NO.: 1021-974

EPA REG. NO.: 655-113

## **FARNAM**

#### GO-FLY

# DDVP AND NATURAL PYRETHRINS

Ready to use fly spray for Horses, Ponies, and Other Livestock

100% active fly spray formulated with effective combination of DDVP

natural pyrethrins.

Direct application to horses provides protection from House Flies,

Horn Flies, Stable Flies, Mosquitoes and Face Flies.

Also controls House Flies when used as a space spray in stables.

# ACTIVE INCREDIENTS:

Pyrethrins	0.030%
2,2-Dichlorovinyl Dimethyl Phosphate	0.180%
Related Compounds	0.014%
Technical Piperonyl Butoxide*	0.250%
INERT INGREDIENTS	99.526%
TOTAL	100.000%

Equivalent to 0.20% (butylcarbityl) (6-propylpiperonyl) ether and 0.05% related compounds.

# KEEP OUT OF REACH OF CHILDREN

## CAUTION

See back panel for additional precautionary statements

NET CONTENTS: 32 FL. CZ. (1 Quart)

Page 1 of 4

## FARNAM GO-FLY

Versatile and easy to use Farnam Go-Fly needs no mixing or preparation. It may be used with conventional hand or automatic sprayers for direct application to animals or as a stable and barn space spray.

#### PRECAUTIONARY STATEMENTS

## HAZARDS TO HUMAN AND DOMESTIC ANIMALS

#### CAUTION

For animal use only. Do not use on humans. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid inhalation of vapor or spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly with scap and water after handling and before eating or smoking. Do not contaminate water, feed or foodstuffs.

# STATEMENT OF PRACTICAL TREATMENT


IF SWALLCWED: Call a physician immediately. Castric lavage is indicated if material was taken internally. DO NOT INDUCE VOMITING unless other treatment is not available. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. Apply respiration if indicated.

IF CN SKIN: Wash contaminated skin with soap and water.

IF IN EYES: Flush eyes with plenty of water for at least 15 minutes. Get medical attention immediately.

Atropine is antidotal and may be administered in conjunction with 2-PAM.



BACK LABEL (Cont)

## ENVIRONMENTAL HAZARDS

Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

#### DIRECTIONS

HCRSES: Apply as a finely atomized spray uniformly distributed over each animal at a rate of one to two fluid ounces per animal. Mist thoroughly, covering all parts of the body, but do not wet the skin. Reduce application rate by one-half for young animals.

DAIRY CATTLE: May also be used on dairy cattle (for milking dairy animals apply spray 20 to 30 minutes before beginning to milk or after milking has been completed. Do not spray udders or lactating dairy animals), and beef cattle at the same rate of application as for horses.

OTHER LIVESTOCK: For use on hogs, sheep or goats, apply at a rate of one ounce per grown animal, or 1/2 ounce for young animals

DO NOT APPLY IN EXCESS OF RATES STATED PER ANIMAL

STAPLES AND BARNS: To control House Flies in stables, dairy barns, or livestock barns, apply as a fog or mist at the rate of 2 fluid ounces per 1,000 cubic feet uniformly distributed throughout the area. Before fogging close doors, windows and other openings to reduce air movement as much as possible. Leave closed for 10 minutes after application is complete, then ventilate the area. May be repeated twice daily, morning and afternoon, as

BACK LABEL (Cont)

needed. Do not apply as barn spray if animals which have received direct application within eight hours remain in the barn.

# STORAGE AND DISPOSAL

Do not reuse empty container. Destroy it by perforating or crushing, wrap and throw in trash.

EPA REG. NO.: 270-145

EPA EST. NO.: 34734-IA-1

Manufactured For:

FARNAM COMPANIES, INC.

Omaha, Nebraska 68112

(c) 1989 Farmam Companies, Inc.

Printed in U.S.A.

8DD9

