

LABEL - FRONT

FARNAM
EQUI-TROL™
Feed-Thru Fly Control

Highly palatable feed additive to prevent the development of stable flies and house
in the manure of treated horses.

ACTIVE INGREDIENT:

BY WEIGHT

2-Chloro-1-(2,4,5-trichlorophenyl) vinyl

dimethyl phosphate..... 2.468% *

INERT INGREDIENTS..... 97.532% **

TOTAL..... 100.000%

* Contains 11.2 gms. RABON® per pound

** Refers only to ingredients which are not larvicidal

† RABON® - Registered T.M. of Shell Chemical Company

ACCEPTED
APR 27 1984
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 270-164

FEED INGREDIENTS: Wheat Bran, Dehydrated Alfalfa Meal, Molasses, Artificial Flavor.

KEEP OUT OF REACH OF CHILDREN
CAUTION
See side panel for additional precautionary statements

NET WT. 3.75 LBS. (1.70 kg.)

(2)

BEST DOCUMENT AVAILABLE

SIDE PANEL

210-164

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

DO NOT USE ON HORSES INTENDED FOR SLAUGHTER.

Feed the recommended dosage to each horse separately to make certain he receives his full portion. This product should be fed top dressed on grain or mixed with the horses total ration to provide 70 mg. of Rabon per 100 lbs. of body weight. For an occasional finicky eater who does not accept new feeds readily, mix Equi-Trol with sweet feed or grain ration.

Start feeding Equi-Trol early in the Spring before flies begin to appear and continue feeding throughout the Summer and into the Fall until cold weather restricts fly activity.


Equi-Trol Feed-Thru Fly Control prevents the development of house flies and stable flies in the manure of treated horses but is not effective against existing adult flies.

In some cases, supplemental fly control measures may be needed in and around paddocks and buildings to control adult house flies and stable flies which can breed not only in manure but in other decaying vegetable matter or silage on premises.

In order to achieve optimum fly control Equi-Trol should be used in conjunction with other good management and sanitation practices.

DAILY DOSAGE

- 300 - 500 lb. Horse Feed 2/5 oz. (One Equi-Trol measuring cup filled to the blue line) per horse per day.
- 500 - 700 lb. Horse Feed 3/5 oz. (One Equi-Trol measuring cup filled to the red line) per horse per day.
- 700 - 900 lb. Horse Feed 4/5 oz. (One Equi-Trol measuring cup filled to the green line) per horse per day.
- 900 - 1100 lb. Horse Feed 1 oz. (One full Equi-Trol measuring cup) per horse per day.



For larger horses over 1100 lb. of body weight feed 1/4 oz. for each 250 lb. of body weight

SIDE PANEL

3

270-164

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by Storage or Disposal.

STORAGE: Store tightly closed in cool dry place away from heat and sunlight.

PRODUCT DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly before discarding in trash.

Manufactured for:

FARNAM COMPANIES, INC.

Omaha, Nebraska 68112 U.S.A.

E.P.A. REG. NO. 270-164

E.P.A. EST. NO. 45277-MO-2

©1983 Farnam Companies, Inc.

Printed in U.S.A.

