

270-165

09/25/2001

1/5

ACCEPTED

SEP 25 2001

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for this pesticide registered under Reg. No. 270-165

FARNAM
VITA PLUS®
WITH EQUITROL®

FEED-THRU FLY CONTROL

Now contains Equitrol® Feed thru fly control to prevent the development of Stable Flies and Houseflies in the manure of treated horses
The Nutritional Feed Supplement for All Classes of Horses
Vitamins · Minerals · Amino Acids · Selenium · Coat Conditioner
With EQUITROL® Feed-Thru Fly Control

ACTIVE INGREDIENT

Tetrachlorvinphos: (CAS #961-11-5)..... 1.234%*

OTHER INGREDIENTS 98.766%**

TOTAL 100.000%

*Contains 5.597 g RABON® per pound.

**Refers only to ingredients which are not larvicidal.

FEED INGREDIENTS

Wheat Middlings, Dehydrated Alfalfa Meal, Wheat Germ Meal, Molasses, Corn, Dicalcium Phosphate, Salt, Dried Whey, Vitamin A Supplement, Vitamin D₃ Supplement, Vitamin E Supplement, Vitamin B₁₂ Supplement, Riboflavin Supplement, Niacin, Calcium Pantothenate, Choline Chloride, Thiamine Mononitrate, Pyridoxide Hydrochloride (Vitamin B₆), Sodium Selenite, L Lysine Hydrochloride, dL-Methionine, Zinc Amino Acid Chelate, Manganese Amino Acid Chelate, Iron Amino Acid Chelate, Copper Amino Acid Chelate, Iodine Amino Acid Chelate, and Cobalt Amino Acid Chelate, Magnesium Sulfate, Potassium Chloride, propionic acid, and acetic acid (preservatives).

Keep out of Reach of Children

CAUTION

See side panel for additional precautionary statements

NET WT: XX

EPA REG. NO. 270-165

EPA EST. NO. 45277-MO-2

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
CAUTION**

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.


FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>In case of medical emergencies or health and safety inquiries, or in case of fire, leaking or damaged containers or for product use information, call toll free: (800) 234-2269</p>	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	

Personal Protective Equipment

- Applicators and Handlers must wear:
- Long-sleeved shirt and pants
 - Shoes and socks
 - Chemical resistant gloves

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.



User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water when disposing of equipment wash water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store tightly closed in cool, dry place inaccessible to children and pets.

PESTICIDE DISPOSAL: Pesticide that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

DISPOSAL: Completely empty container. Then, dispose of empty container in a sanitary landfill or by incineration or if allowed by State and local authorities, by burning. If burned, stay out of smoke. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

GENERAL INSTRUCTIONS AND INFORMATION

Feed the recommended dosage to each horse separately to make certain he receives his full portion. This product should be fed top dressed on grain or mixed with the horse's total ration to provide 70 mg of Rabon® per 100 lbs. of body weight. For an occasional finicky eater who does not accept new feeds readily, mix VITA PLUS® WITH EQUITROL® with sweet feed or grain ration.

Begin feeding VITA PLUS® WITH EQUITROL® early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall until cold weather restricts fly activity.

VITA PLUS® WITH EQUITROL® Feed-Thru Fly Control prevents the development of House Flies and Stable Flies in the manure of treated horses but is not effective against existing adult flies. In some cases, supplemental fly control measures may be needed in and around paddocks and buildings to control adult House Flies and Stable Flies which can breed not only in manure but in other decaying vegetable matter or silage on premises.

In order to achieve optimum fly control, VITA PLUS® WITH EQUITROL® should be used in conjunction with other good management and sanitation practices.

GENERAL PRECAUTIONS AND RESTRICTIONS

DO NOT USE ON HORSES INTENDED FOR SLAUGHTER

CAUTION: Follow label directions. The addition to feed of higher levels of this premix containing Selenium is not permitted.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift.

Only protected handlers may be in the area during application.

GUARANTEED ANALYSIS

Each pound contains not less than:

	Per Ounce	Per Pound
Lysine		0.05 %
Methionine		0.05 %
Calcium (min)	766 mg	2.75 %
Calcium (max)	836 mg	3.75 %
Phosphorous (min)	326 mg	1.15 %
Salt (min)	255 mg	0.9 %
Salt (max)	397 mg	1.4 %
Potassium (K)	241 mg	0.85 %
Magnesium (Mg)	12.5 mg	0.044 %
Copper (Cu)	4.0 mg	141 ppm
Selenium (Se)	120 mg	4 ppm
Zinc (Zn)	20.0 mg	705 ppm
Iodine (I)	1.0 mg	35 ppm
Iron (Fe)	100.0 mg	3,527 ppm
Cobalt (Co)	0.05 mg	18 ppm
Manganese (Mn)	10.0 mg	353 ppm
Vitamin A	25,000 IU	400,000 IU
Vitamin D ₃	2,500 IU	40,000 IU
Vitamin E	25 IU	400 IU
Vitamin B ₁₂	200 mg	3,200 mg/lb
Riboflavin (B ₂)	25 mg	400 mg/lb
d-Pantothenic Acid	62.5 mg	1,000 mg/lb
Thiamine (B ₁)	12.5 mg	200 mg/lb
Niacin	125.0 mg	2,000 mg/lb
Vitamin B ₆	5.0 mg	80 mg/lb
Choline	106.0 mg	1,696 mg/lb
Polyunsaturated Fatty Acids (as glycerides)		2.75 %

DIRECTIONS FOR USE (Application Instructions)

DAILY DOSAGE

- 300-500 lb. Horse Feed 2/5 oz. (One EQUI-TROL measuring cup filled to the blue line) per horse per day.
- 500-700 lb. Horse Feed 3/5 oz. (One EQUI-TROL measuring cup filled to the red line) per horse per day.
- 700-900 lb. Horse Feed 4/5 oz. (One EQUI-TROL measuring cup filled to the green line) per horse per day.
- 900-1100 lb. Horse Feed 1 oz. (One full EQUI-TROL measuring cup) per horse per day.

For larger horses over 1100 lbs. of body weight feed $\frac{1}{4}$ oz. for each 250 lbs. of body weight.

EPA REG. NO. 270-165

EPA EST. NO. 45277-MO-2

©RABON is a registered trademark of Boehringer Ingelheim.

FARNAM COMPANIES, INC.
 Horse Products Division
 P.O. Box 34820 • Phoenix, AZ 85067-4820
www.farnam.com
 (602) 285-1660, M-F, 7am-3:45pm, MST

Product Code: 013920

©2001 Farnam Companies, Inc.

Printed in U.S.A.

XXX