

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**DO NOT USE ON HORSES INTENDED FOR SLAUGHTER**

To protect horses from Horn Flies, House Flies, Mosquitoes, and Gnats, apply a light mist sufficient to wet the surface of the hair. To control Stable Flies, Horse Flies, and Deer Flies, apply at a rate of 2 ounces per adult horse, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control Blood-sucking Lice, apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 2 to 3 weeks if required.

**STORAGE:** Do not store under 32 ° F. Store in a place inaccessible to children and pets.

**DISPOSAL:** When empty, wrap container in several layers of newspaper and discard in trash.

**E.P.A. REG. NO. 270-172**

**E.P.A. EST. NO. 6658-MO-1**

Manufactured for:  
FARNAM COMPANIES, INC.  
Omaha, Nebraska 68112

296

Farnam's Water Base Fly Repel gives you the perfect show combination - a high-tech silicon shine and a pyrethrin fly repellent in one product. All the advantages of a silicon polish - detangles manes and tails quickly; a lustrous health looking coat that shines like a winner; silicone formula repels dust and dirt and prevents stains from grass, urine, manures and latigo - plus protection from annoying flies.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. Remove pets, bird and cover fish aquaria before space spraying or surface applications. Avoid contamination of feed and foodstuffs.

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person. Call a physician or Poison Control Center immediately. **IF ON SKIN:** Remove contaminated clothing and wash affected areas with soap and water. Seek medical attention if irritation persists. **IF IN EYES:** Flush eyes with plenty of water. Seek medical attention if irritation persists. **IF INHALED:** Remove victim to fresh air. Apply artificial respiration if indicated. Seek medical attention if indicated.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not apply directly to lakes, ponds, streams, tidal marshes or estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herein.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**DO NOT USE ON HORSES INTENDED FOR SLAUGHTER**

**INDOOR APPLICATION:**

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned with potable water before using.

Vacate treated areas and ventilate before reoccupying.

For maximum effectiveness, a combination of spot, surface treatment and fogging is recommended.

**SURFACE SPRAYING:** To control Cockroaches, Palmetto Bugs, Ants, Silverfish, Spiders, Crickets, Clovermites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Sawtoothed Grain Beetles, Spider Beetles, Cigarette Beetles, Drugstore Beetles, Meal Worms, Grain Mites, and Cadelles: Use a good sprayer adjusted to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers, behind and beneath sinks, stoves, refrigerators and cabinets, along baseboards, around door and window frames, into dark corners and in and around garbage compactors, garbage chutes and cans. For Silverfish, spray bookcases. For Ants, spray trails, nests, and points of entry. If surface application only is to be used, spray floors and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

In case of infested foods in cabinets, dispose of the infested material. Remove all contents and clean thoroughly. Spray interior surfaces of empty cabinets thoroughly. After spray has dried, cover shelves with paper or otherwise protect foods from direct contact with sprayed surfaces.

To control Carpet Beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

To control Bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

**Pet Areas:** To control Fleas and Brown Dog Ticks on premise: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals is recommended. Repeat treatment as needed.

To control Clothes Moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

**BEST AVAILABLE COPY**

**SPACE SPRAYING:** Close doors, windows and transoms and shut off ventilating system. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of House Flies, Fruit Flies, Gnats, Mosquitoes, Skipper Flies, Wasps, Hornets, Angoumois Grain Moths, Mediterranean Flour Moths, Indianmeal Moths, and Tobacco Moths, direct spray at an upward angle distributing it uniformly throughout the entire area at a rate of 3-6 ounces per 1000 cubic feet of space. Do not remain in treated areas. If insects have not dropped to floor in 3 minutes, repeat spraying as quantity sprayed was insufficient. Ventilate the area after treatment is completed. To aid in the prevention of breeding, spray on standing water and in garbage pails.

**FOGGING:** To kill exposed or accessible stages of the above named crawling insects, use a rate of 1 gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly whenever possible.

**ON LIVESTOCK:** To protect cattle, horses, and hogs from Horn Flies, House Flies, Mosquitoes and Gnats apply a light mist sufficient to wet the surface of the hair.

To control Stable Flies, Horse Flies and Deer Flies, apply at a rate of 2 ounces per adult animal sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control Blood-sucking Lice, apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 2 to 3 weeks if required.

To control Poultry Lice, spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

For control of Bedbugs and Mites in poultry houses, spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbug and mites hide.

**FLEAS AND TICKS ON DOGS AND CATS:** Start spraying at the animal's head, avoiding the eyes, nose and mouth as you spray. Make sure the animal's entire body is covered including the legs under body and tail. While spraying, fluff the hair so that the spray will penetrate the skin. Make sure spray wets thoroughly. Avoid contact with genitalia. Repeat as necessary.

**ADULT MOSQUITO CONTROL:** Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where Mosquitoes may rest (low, marshy places, stagnate pools or ponds and shore line). Apply at a rate of 2.5 pints per acre (0.1 ounce per 100 square feet). Using truck-mounted conventional misting equipment such as Rotomist® models 51C, 91 or 100HT traveling at 5 m.p.h. to treat a 300 foot swath calibrate equipment to deliver 0.9 gallon per minute.

**BEST AVAILABLE COPY**

**OUTDOORS:** (Camping, fishing, hunting) - Spray outer clothing and surroundings. Repels Biting Flies, Mosquitoes, Midges and Gnats. If sprayed every hour or so, will aid in keeping surroundings free from Mosquitoes. If in rare cases skin irritation results, avoid continued contact. Do not pour or squirt - use a good sprayer which will throw a heavy fine mist.

**USE AS A WIPE ON:** First, Brush animal to remove excess dirt and dust. For horses face, always apply as a wipe using a piece of clean, absorbent cloth or toweling (Turkish). Rub over the hair. It is best to apply by rubbing against the hair growth. Give special attention to the neck and facial areas where flies most often are seen. Only a light application is required. Avoid using an excessive amount on your horses. Do not wet skin. After application, brush thoroughly to bring out bright sheen on the coat. Repeat as required. Wash hands with soap and water after use.

**USE WITH ENCLOSED TOWEL APPLICATOR:** Open container cap. Remove foil seal; discard in appropriate place. Pull out top towel and close container cap securely. Recommend for use on dogs, cats, small domestic animals, horses, ponies and other stable and barn animals. To protect animals from Horn Flies, Deer Flies, Mosquitoes, House Flies, Horse Flies, Gnats, Lice, Fleas and Ticks; First brush animal to remove excess dirt and dust. Use towel to apply a light coating sufficient to wet the tips of the hair. It is best to apply by rubbing against the hair growth. Only a light application is required. Avoid using an excessive amount. Do not wet skin. After application, brush coat thoroughly. When finished discard used towel in appropriate place. Repeat as necessary. Wash hands with soap and water after use.

STORAGE AND DISPOSAL

Do not transport or store under 32°F. Store in a place inaccessible to children and pets.

Containers larger than one gallon:

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE AND SPILL PROCEDURES: Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste. Protect from freezing.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other State and local procedures.

Containers one gallon or less:

STORAGE: Store in original container in a cool dry place inaccessible to children and pets. Protect from freezing.

DISPOSAL: Wrap container in several layers of newspaper and discard in trash.

EPA REG. NO. 270-172

EPA EST. NO.

Distributed by:

FARNAM COMPANIES, INC.  
Omaha, NE 68112 U.S.A.

©1990 Farnam Companies, Inc.  
Product Code                      Printed in USA