


WATER BASE FLY REPEL



- Fly Spray for Horse
- High Tech Hair Polish
- Excellent for horses sensitive to oil base fly sprays
- Contains less than 1% oil
- Contains pyrethrins
- Leaves no oily residue to attract dust
- Ready-to-use
- To protect horse from Horn Flies, Stable Flies, Deer Flies, Mosquitoes, House Flies, Horse Flies, Gnats, and Lice
- Contains more active ingredients than the original Bronco Fly Spray
- A water based micro emulsion with pleasant citronella scent
- A concentrated pyrenone® micro emulsion
- Clear stable leaves no oily or objectionable residues
- Effective, fast acting insecticide
- Kills and repels biting flies
- For use in homes, institutions, restaurants, food processing plants, dairies, livestock and outdoors for mosquito control
- For control of fleas and brown dog ticks on premises and on pets
- Apply with conventional mechanical or compressed air equipment

ACTIVE INGREDIENTS

Pyrethrins.....	0.10%
*Piperonyl Butoxide, Technical.....	1.00%
OTHER INGREDIENTS .....	<u>98.90%</u>
TOTAL.....	100.00%

- Equivalent to min. 0.8% (butylcarbityl) (6-propylpiperonyl) ether and
- 0.2% related compounds
- Do not transport or store under 32°F

PYRENONE - Registered trademark of Fairfield American Corporation

<p>KEEP OUT OF REACH OF CHILDREN CAUTION See back panel for additional cautions</p>
---

**Net Contents:**

Farnam's Water Base Fly Repel gives you the perfect show combination - a high-tech silicon shine and a pyrethrin fly repellent in one product. All the advantages of a silicon polish - detangles manes and tails quickly; a lustrous healthy looking coat that shines like a winner; silicone formula repels dust and dirt and prevents stains from grass, urine, manures and latigo - plus protection from annoying flies.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS & COMESTIC ANIMALS  
CAUTION**

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. Remove pets, birds and cover fish aquaria before space spraying or surface applications. Avoid contamination of feed and foodstuffs.

<b>FIRST AID</b>	
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

IF INHALED:	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<p align="center"><b>In case of medical emergencies or health and safety inquiries, or in case of fire, leaking or damaged containers or for product use information, Call 602-285-1660</b></p>	
<p align="center">Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	

### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply where run-off is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.


#### DO NOT USE ON HORSES INTENDED FOR SLAUGHTER

##### INDOOR APPLICATION:

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during or thoroughly cleaned with potable water before using.

Vacate treated areas and ventilate before reoccupying.

For maximum effectiveness, a combination of spot, surface treatment and fogging is recommended.



**SURFACE SPRAYING:** To control Cockroaches, Palmetto Bugs, Ants, Silverfish, Spiders, Crickets, Clovermites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Sawtoothed Grain Beetles, Spider Beetles, Cigarette Beetles, Drugstore Beetles, Meal Worms, Grain Mites, and Cadelles: Use a good sprayer adjusted to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers, behind and beneath sinks, stoves, refrigerators and cabinets, along baseboards, around door and window frames, into dark corners and in and around garbage compactors, garbage chutes and cans. For Silverfish, spray bookcases. For Ants, spray trails, nests, and points of entry. If surface application only is to be used, spray floors and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

In case of infested foods in cabinets, dispose of the infested material. Remove all contents and clean thoroughly. Spray interior surfaces of empty cabinets thoroughly. After spray has dried, cover shelves with paper or otherwise protect foods from direct contact with sprayed surfaces.

To control Carpet Beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

To control Bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

**Pet Areas:** To control Fleas and Brown Dog Ticks on premise: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals is recommended. Repeat treatment as needed.

To control Clothes Moths: Remove any infested articles from storage; brush thoroughly and air for several hours in sunlight if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the

spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

**SPACE SPRAYING:** Close doors, windows and transoms and shut off ventilating system. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of House Flies, Fruit Flies, Gnats, mosquitoes, Skipper Flies, Wasps, Hornets, Angoumois Grain Moths, Mediterranean Flour Moths, Indianmeal Moths, and Tobacco Moths, direct spray at an upward angle distributing it uniformly throughout the entire area at a rate of 3-6 ounces per 1000 cubic feet of space. Do not remain in treated areas. If insects have not dropped to floor in 3 minutes, repeat spraying as quantity sprayed was insufficient. Ventilate the area after treatment is completed. To aid in the prevention of breeding, spray on standing water and in garbage pails.

**FOGGING:** To kill exposed or accessible stages of the above named crawling insects, use a rate of one gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly whenever possible.

**ON LIVESTOCK:** To protect cattle, horses, and hogs from Horn Flies, House Flies, Mosquitoes and Gnats apply a light mist sufficient to wet the surface of the hair.

To control Stable Flies, Horse Flies and Deer Flies, apply at a rate of 2 ounces per adult animal sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control Blood-sucking Lice, apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 2 to 3 weeks if required.

To control Poultry Lice, spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

For control of Bedbugs and mites in poultry houses, spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

6/7

**FLEAS AND TICKS ON DOGS AND CATS:** Start spraying at the animal's head, avoiding the eyes, nose and mouth as you spray. Make sure the animal's entire body is covered including the legs, underbody and tail. While spraying, fluff the hair so that the spray will penetrate the skin. Make sure spray wets thoroughly. Avoid contact with genitalia. Repeat as necessary.

**ADULT MOSQUITO CONTROL:** Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest (low, marshy places, stagnate pools, on ponds and shore line). Apply at a rate of 2.5 pints per acre (0.1 ounce per 100 square feet). Using truck-mounted conventional misting equipment such as Rotomist® models 51C, 91 or 100HT traveling at 5 mph to treat a 300 foot swatch, calibrate equipment to deliver 0.9 gallons per minute.

**OUTDOORS:** (Camping, fishing, hunting) - Spray outer clothing and surroundings. Repels biting flies, mosquitoes, midges and gnats. If sprayed every hour or so, will aid in keeping surroundings free from mosquitoes. If in rare cases skin irritation results, avoid continued contact. Do not pour or acquire - use a good sprayer which will throw a heavy fine mist.

**USE AS A WIPE ON:** First, brush animal to remove excess dirt and dust. For horse's face, always apply as a wipe using a piece of clean, absorbent cloth or toweling (Turkish). Rub over the hair. It is best to apply by rubbing against the hair growth. Give special attention to the neck and facial areas where flies most often are seen. Only a light application is required. Avoid using an excessive amount on your horses. Do not wet skin. After application, brush thoroughly to bring out bright sheen on the coat. Repeat as required. Wash hands with soap and water after use.

Delete this paragraph:

**USE WITH ENCLOSED TOWEL APPLICATOR:** Open container cap. Remove foil seal; discard in appropriate place. Pull out top towel and close container cap securely. Recommend for use on dogs, cats, small domestic animals, horses, ponies and other stable and barn animals. To protect animals from Horn Flies, Deer, Flies, mosquitoes, House Flies, Horse Flies, Gnats, Lice, Fleas and Ticks:

7/7

### STORAGE AND DISPOSAL

Do not transport or store under 32°F. Store in a place inaccessible to children and pets. Containers larger than one gallon: Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE AND SPILL PROCEDURES:** Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste. Protect from freezing.

Containers one gallon less: Store in original container in a cool dry place inaccessible to children and pets. Protect from freezing

**DISPOSAL:** If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or call 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

EPA Est. No. 270-172

EPA Est. No. \_\_\_\_\_

