

4- 9 270-238 111 155 - 12 8
IMMEDIATE CONTAINER - GLASS VIAL

DIAZINON LIQUID CONCENTRATE INSECT SPRAY

ACTIVE INGREDIENTS:

Diazinon [0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate].....25.0 %

INERT INGREDIENTS:75.0 %

KEEP OUT OF REACH OF CHILDREN

WARNING

See Back of Outer Package for Additional Precautions

EPA Reg. No. 270-238

EPA Est. No.: 731-MA-1

Net Contents: .08 oz

Not Reviewed. Registrant claims to be in accordance with
Registration of Pesticide Product Guidance for Phase 2
EPA Reg. No. 270-238, Active to Inert Change in Status.

OUTSIDE CONTAINER - BLISTER CARD
FRONT PANEL

UNI-SPRAY

Pre-measured treatments - use one bottle each time you spray

DIAZINON LIQUID CONCENTRATE INSECT SPRAY

Prevents and controls damage to yard and garden plants caused by
many common insects as listed*

ACTIVE INGREDIENTS:

Diazinon [0,0-diethyl 0-(2-isopropyl-6-methyl-4-
pyrimidinyl) phosphorothioate]..... 25.0%

INERT INGREDIENTS:..... 75.0%

Diazinon is a registered trademark of Ciba-Geigy Corporation

KEEP OUT OF REACH OF CHILDREN

WARNING

See Back For Additional Precautions

Each Uni-Spray bottle dilutes to make one quart of spray, enough for 1
average application*

No measuring - Accurate and Easy to Use

Mix exactly what you need

Convenient - No storage of partially used bottles

*Refer to back for complete list of pests controlled and Use Directions

Remove bottles as needed from back

NET CONTENTS: 8 bottles, 0.08 Fl. Oz. Each

OUTSIDE CONTAINER - BLISTER CARD
BACK PANEL

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Do not use in home. Keep out of reach of children and domestic pets. May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Do not use on household pet or humans. Do not permit children or pets to go onto the treated areas until the spray has completely dried.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control center immediately. If possible, vomiting should be induced under medical supervision. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES: Flush eyes with plenty of water. Call a physician immediately.

Note to Physician: Gastric lavage is indicated. If symptoms of cholin-

4 2 5

OUTSIDE CONTAINER - BLISTER CARD

BACK PANEL (cont.)

terase inhibition are present Atropine Sulfate is antidotal. 2-pam is also antidotal and may be used in conjunction with atropine.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife . Birds feeding on treated areas may be killed. Keep out of lakes, streams, and ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining in the treated areas. Do not apply when bees are actively visiting the crop, crop cover, or weeds blooming in treated areas. Applications should be timed to provide maximum possible interval between treatment and the next period of bee activity.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill, or store near heat or open flame.

OUTSIDE CONTAINER - BLISTER CARD

BACK PANEL (cont)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Use any clean sprayer. Spray with medium to fine mist for thorough coverage. Flush sprayer with clean water after each use. Spray only as directed. Make applications when insects first appear.

VEGETABLES: Foliar application for the listed crops and insects. Mix one Uni-Spray bottle of Diazinon in 1 qt. (32 oz) of water. To cover approximately 80 sq. ft. Generally, applications at 7 to 10 day intervals may be required. Do not feed treated vines to livestock.

CROP	INSECT
Beans, Peas	Aphids, Cucumber Beetles, Grasshoppers, Mexican Bean Beetles, Mites.
Broccoli, Cabbage, Cauliflower, Collards, Kale, Radishes, Turnips	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers.
Carrots, Celery, Endive, Lettuce, Parsnips, Spinach	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Mites.
Cantaloupes, Cucumbers, Muskmelons, Squash (summer, winter), Watermelon	Aphids, Thrips, Mites.
Peppers, Potatoes, Tomatoes	Aphids, Colorado Potato Beetles, Fall armyworms (tomato), Flea Beetles, Leafhoppers, Southern Armyworms (potato and tomato), Vinegar Flies [Drosophila ssp (tomato)].
Onions	Onion Thrips
Sweet Corn	Aphids, Flea Beetles, Grasshoppers, Mites

After spraying, do not pick vegetables within:

1 day - Peas, Tomatoes, Sweet Corn

3 days - Cantaloupes, Muskmelons, Winter Squash, Watermelons

OUTSIDE CONTAINER - BLISTER CARD
BACK PANEL (cont)

5 days - Broccoli, Cauliflower, Peppers

7 days - Beans, Cabbage, Cucumbers, Summer Squash

10 days - Carrots, Celery, Collards, Endive, Kale, Lettuce, Onions, Parsnips,
Radishes, Spinach, Turnips.

35 days - Potatoes

VEGETABLES: Preplant soil application. Mix one Uni-Spray bottle of Diazinon in water as specified below.

Cabbage, Carrots,
Corn, Lettuce, Peas,
Pole Beans
Radishes, Red Beets,
Snap Beans, Tomatoes,
Turnips

Cutworms (surface, subterranean) 1 bottle per 14 square feet in sufficient water to achieve thorough coverage. Apply broadcast just before planting. Immediately work into soil 2-3 inches for Surface Cutworms, 3-6 inches for Subterranean Cutworms.

Garden Symphylans (Garden Centipedes)
1 bottle per 6 square feet in sufficient water to achieve thorough coverage. Apply broadcast just before planting. Immediately work into soil 4-8 inches.

Mole Crickets: Carrots, Corn, Peas, Pole Beans, Red Beets, and Snap Beans only. 1 bottle per 55 square feet in sufficient water to achieve thorough coverage. Apply broadcast just before planting. Apply 1-2 days before sowing or transplanting. Immediately work into soil 1-2 inches.

Root Maggots: Cabbage only
1 bottle per 18 square feet in sufficient water to achieve thorough coverage. Apply broadcast just before planting. Immediately work into soil 3-4 inches before seeding, either in seed bed or in field or before transplanting.

OUTSIDE CONTAINER - BLISTER CARD

BACK PANEL (cont)

Wireworms: Cabbage, Carrots, Corn, Lettuce, Peas, Snap Beans, and Tomatoes only. 1 bottle per 14 square feet in sufficient water to achieve thorough coverage. Apply broadcast just before planting. Immediately work into soil 4-8 inches.

TREES & SHRUBS

Wet leaves and branches to the dripping point. Try to penetrate dense foliage.

Arborvitae, Azalea
Birch, Boxwood, Camellia
Douglas Fir, Elm, Hawthorn,
Holly Juniper, Lilac,
Locust Maple, Mimosa Oak,
Pine, Plum Poplar,
Rhododendron, Spruce Willow

Aphids, Bagworms, European Pine Shoot Moths, Holly Bud Moths, Dipterous Leafminers Douglas (Arborvitae, Azalea, Birch, Boxwood, Holly Oak), Leafhoppers, Mealybugs, Mites (Clover, Cyclamen, European Red, Privet, Twospotted Spider), Obscure Root Weevils, Omnivorous Leaf-tiers, Scale Crawlers, (cottoncushion, Euonymus, Lecanium, Pine Needle, Pit Scale, San Jose Soft)
1 bottle in 1 qt. (32 oz.)

Webworms (Juniper, Mimosa*), Whiteflies
1 bottle in 1 quart (32 oz) of water. Apply as a thorough foliar spray.

*Mimosa Webworms. Apply as a thorough foliar spray when adults first appear. Make two additional applications at 4-5 week intervals.

Apple and Thorn Skeletonizers, Hemlock Chermes. Oak Loopers, Obliquebanded Leaf-rollers, Pear Slugs, Tent Caterpillars, Webworms (Cotton-easter, Fall)
1 bottle in 10 oz. of water. Apply as a thorough foliar spray.

ROSES AND FLOWERS

Wet the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage.

Azaleas, Camellias,
Chrysanthemums,
Carnations, Gladioli,
Lilac, Rhododendron,
Roses

Aphids, Euonymus Scale Crawlers, Flea Beetles, Dipterous Leafminers, Leafhoppers, Mealybugs, Mites (Carnation Bud, Carnation Shoot, Cyclamen, European Red, Twospotted Spider), Thrips, Whiteflies.
1 bottle in 1 qt. (32 oz)

OUTSIDE CONTAINER - BLISTER CARD
BACK PANEL

STORAGE AND DISPOSAL

Do not reuse empty container bottle. Wrap container and put in trash collection.

LIMITED WARRANTY: Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions.

EPA REG. NO.: 270-238

EPA EST. NO.: 731-MA-1

Manufactured For:

Farnam Companies, Inc.

Phoenix, Arizona 85013

(c) 1989 Farnam Companies, Inc. Printed in U.S.A. 8009