10919

## UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

Ms. Cheri Purcell Regulatory Specialist Farnam Companies, Inc. 301 W. Osborn Rd. Phoenix, AZ 85013

APR 1 2 2011

Dear Ms. Purcell:

Subject:

Label Amendment to Correct Application Directions on Label Stamped March 21, 2011

Bromadiolone Bait Bar

EPA Registration No. 270-373 Submission Date: April 4, 2011

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) as amended, is acceptable. A stamped copy is enclosed for your records. Please submit one (1) final printed copy before releasing the product for shipment. If you have any questions regarding this label, please contact Jennifer Gaines at (703) 305-5967 or via e-mail at <a href="mailto:gaines.jennifer@epa.gov">gaines.jennifer@epa.gov</a>.

Sincerely yours,

Venus Eagle

Product Manager (01)

Insecticide-Rodenticide Branch Registration Division (7505P)

### ACCEPTED

APR 1 2 2011

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide Registered under EPA Reg. No. 200-373

# AGRICULTURAL USE LABEL (GREATER THAN OR EQUAL TO 8 LBS. OF BAIT)

BROMADIOLONE BAIT BAR
[ABN: JUST ONE BITE\* II BAR]
[ABN: JUST ONE BITE\* II BAIT CHUNKS]

\*NORWAY RATS AND HOUSE MICE MAY CONSUME A LETHAL DOSE IN A SINGLE NIGHT'S FEEDING WITH FIRST DEAD RODENTS APPEARING FOUR OR FIVE DAYS AFTER FEEDING BEGINS For Use In and Around Agricultural Buildings Only

FOR INDOOR AND OUTDOOR RAT AND HOUSE MOUSE CONTROL KILLS NORWAY RATS, ROOF RATS, HOUSE MICE AND WARFARIN RESISTANT NORWAY RATS

KILLS HOUSE MICE

KILLS RATS & HOUSE MICE

KILLS HOUSE MICE AND RATS

RAT AND HOUSE MOUSE CONTROL CHUNKS

RAT AND HOUSE MOUSE CONTROL BAR

- With grains and seeds
- Palatable formulation
- Rat & House Mouse Killer
- (Farnam®) Rat and House Mouse Control
- (Bait chunks) (Bar pieces) fit in burrows, runways (and) (&) hard-to-reach areas where rats and house mice live and breed
- Can be used in bait stations
- (Features) nibble ridges (to promote rat and house mouse chewing)
- Multi-edge shape gives rats and house mice gnawing edges
- 2 oz Bait chunks for bait stations
- 2 oz Bar pieces for bait stations
- Bars break into 2 oz pieces
- Bars break into 2 oz mini-bars for bait stations
- The hole through the middle allows for mounting in bait stations
- Helps Provide (effective) control
- Helps control rats and house mice
- (Bait Chunks) (Bars) (Bait Bars)
- Individually wrapped 1 pound Bars (helps to keep product fresh.)
- (Contains) (8) (9) (10) (11) (12) (13) (14) (15) one pound (1 lb) Bars
- (Contains) (64) (72) (80) (88) (96) (104) (112) (120) 2 oz Bait Chunks

- (Economical) Value (Pack) (Size) contains (8, 9, 10, 11, 12, 13, 14, 15) pounds of product!
- Great value! Contains (8, 9, 10, 11, 12, 13, 14, 15) pounds of product!

#### **ACTIVE INGREDIENT:**

Bromadiolone	0.005%
OTHER INGREDIENTS*:	99.995%
TOTAL:	

<sup>\*</sup>Contains denatorium benzoate

#### KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional precautionary statements

EPA Reg. No. 270-373

EPA Est. No. 75806-AZ-1

#### **NET CONTENTS:**

8 lbs (64, 2 oz chunks) or 9 lbs (72, 2 oz chunks) or 10 lbs (80, 2 oz chunks) or 11 lbs (88, 2 oz chunks) or 12 lbs (96, 2 oz chunks) or 13 lbs (104, 2 oz chunks) or 14 lbs (112, 2 oz chunks) or 15 lbs, (120, 2 oz chunks) or 8 lbs (8, 16 oz bars) or 9 lbs (9, 16 oz bars) or 10 lbs (10, 16 oz bars) or 11 lbs (11, 16 oz bars) 12 lbs, (12, 16 oz bars) or 13 lbs (13, 16 oz bars) or 14 lbs (14, 16 oz bars) or 15 lbs (15, 16 oz bars)

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Keep away from children, domestic animals and pets. Causes moderate eye irritation. Do not get in eyes or on clothing. All handlers (including applicators) must wear long-sleeved shirt and long pants, shoes plus socks, and gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.

#### User\_Safety\_Requirements\_

Follow manufacturer's instruction for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait

<sup>\*</sup>This product has been treated with S-methoprene to protect the bait from infestation by insects.

and before eating, drinking, chewing gum, using tobacco, or using the toilet and change into clean clothing.

FIRST AID  Have label with you when obtaining treatment advice.		
IF SWALLOWED:	<ul> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> </ul>	
IF IN EYES:	<ul> <li>Do not give anything by mouth to an unconscious person.</li> <li>Hold eye open and rinse slowly and gently with water for</li> </ul>	
	<ul> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>	

#### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-234-2269 for emergency medical treatment information.

#### **NOTE TO PHYSICIAN**

If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin  $K_1$  intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

#### TREATMENT FOR PET POISONING

If animal eats bait, call veterinarian or 1-800-234-2269 [optional] at once.

#### NOTE TO VETERINARIAN

Anticoagulant Bromadiolone: For animals ingesting bait and/or showing poisoning signs (bleeding or alleviated prothrombin times), give Vitamin  $K_1$ . If needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.

#### **ENVIRONMENTAL HAZARDS**

This product is extremely toxic to mammals and birds. Dogs, cats, and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. This pesticide is toxic to fish. Do not apply this product directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may also be hazardous to aquatic organisms in water adjacent

to treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its label.

READ THIS LABEL: READ THIS ENTIRE LABEL AND USE STRICTLY IN ACCORDANCE WITH USE DIRECTIONS, USE RESTRICTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. USE ONLY FOR THE SITES, PESTS, AND APPLICATION METHODS DESCRIBED ON THIS LABEL.

It is illegal to sell these place packs individually.

**IMPORTANT:** Do not expose children, pets or other nontarget animals to rodenticides. To help prevent accidents:

- 1. Store product not in use in a location out of reach of children and pets.
- 2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears or other potentially destructive animals, or in areas prone to vandalism.
- 3. Dispose of product container, unused, spoiled and unconsumed bait as specified on this label.

**Note:** Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used if children, pets, nontarget mammals, or birds may access the bait.

Do not apply this product in a way that will contact workers or other persons. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

USE RESTRICTIONS: For control only of Norway Rats, Roof Rats and House Mice in and around agricultural buildings only. Do not apply further than 50 feet from agricultural buildings. Do not use in homes or other human residences. Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs,

food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food. Do not broadcast bait.

Do not sell this product in individual containers holding less than 8 pounds of bait.

#### SELECTION OF TREATMENT AREAS

Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rats and/or mice or their signs have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

#### APPLICATION DIRECTIONS

#### APPLICATION DIRECTIONS - BAR

Each [name] Bait Bar weighs one pound and is scored to be broken into 2-oz pieces.

RATS: While wearing gloves, apply two to eight 2-oz pieces of bar (or one entire BAR) at each placement location in infested areas. For burrow applications, insert one to two 2-oz pieces of bar at least six inches into each active burrow. Check treated burrows each morning and evening to make sure that baits have not been pushed out of burrows. For aboveground treatment, space bait placements 15 to 30 feet apart. Maintain an uninterrupted supply of fresh bait for 10 days or until fresh signs of rat activity cease to appear.

MICE: While wearing gloves, break bar into 2 oz. pieces and apply one 2-oz bar piece per placement in infested areas at 8 to 12 foot intervals in infested areas. Maintain an uninterrupted supply of fresh bait for 15 days or until fresh signs of mouse activity cease to appear.

**FOLLOW-UP**: Replace contaminated or spoiled bait immediately. Wearing gloves, collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, eliminate sources of rodent food, water and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

#### APPLICATION DIRECTIONS – BAIT CHUNKS

Each [name] weighs 2-oz.

**RATS:** While wearing gloves, apply two to eight 2-oz blocks at each placement location in infested areas. For burrow applications, insert one to two 2-oz pieces of block at least six inches into each active burrow. Check treated burrows each morning and

evening to make sure that baits have not been pushed out of burrows. For aboveground treatment, space placements 15 to 30 feet apart in infested areas. Maintain an uninterrupted supply of fresh bait for 10 days or until fresh signs of rat activity cease to appear.

MICE: While wearing gloves, apply one 2-oz chunk per placement in infested areas. Space placements at 8 to 12 foot intervals. Maintain an uninterrupted supply of fresh bait for 15 days or until fresh signs of mouse activity cease to appear.

**FOLLOW-UP:** Replace contaminated or spoiled bait immediately. Wearing gloves, collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

**PESTICIDE STORAGE:** Store only in original container, in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

**PESTICIDE DISPOSAL:** Dispose of wastes resulting from the use of this product in trash or at an approved waste disposal facility.

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. [Plastic:] Triple rinse (or equivalent), then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or by incineration. In most states, burning is not allowed. [Paper:] Dispose of empty container by placing in trash, at an approved waste disposal facility or by incineration.

#### CONDITIONS OF SALE

To the extent consistent with applicable law FARNAM COMPANIES, INC., warrants this product in its unopened package conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions as specified on this label. There are no other warranties, expressed or implied, concerning the use of this product other than as indicated on the label. To the extent consistent with applicable law, this warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or conditions not reasonably foreseeable to seller, and buyer assumes all risk of any such use.

Manufactured for:
FARNAM COMPANIES, INC.
P.O.Box 34820 • Phoenix, AZ 85067-4820
www.farnam.com
Call with comments or questions: 1-800-234-2269
© [DATE] Farnam Companies, Inc.

#### NEW LABEL

INDUSTRIAL / COMMERCIAL USE LABEL (GREATER THAN OR EQUAL TO 16 LBS. OF BAIT)

BROMADIOLONE BAIT BAR
[ABN: JUST ONE BITE\* II BAR]
[ABN: JUST ONE BITE\* II BAIT CHUNKS]

\*NORWAY RATS AND HOUSE MICE MAY CONSUME A LETHAL DOSE IN A SINGLE NIGHT'S FEEDING WITH FIRST DEAD RATS AND MICE APPEARING FOUR OR FIVE DAYS AFTER FEEDING BEGINS.

For Use In and Around Industrial and Commercial Buildings Only

For indoor and outdoor rat and house mouse control

KILLS NORWAY RATS, ROOF RATS, HOUSE MICE AND WARFARIN RESISTANT NORWAY RATS

NORWAY RATS AND HOUSE MICE MAY CONSUME A LETHAL DOSE IN A SINGLE NIGHT'S FEEDING WITH FIRST DEAD RATS AND MICE APPEARING FOUR OR FIVE DAYS AFTER FEEDING BEGINS.

# KILLS HOUSE MICE KILLS RATS & HOUSE MICE KILLS HOUSE MICE AND RATS RAT AND HOUSE MOUSE CONTROL CHUNKS RAT AND HOUSE MOUSE CONTROL BAR

- With grains and seeds
- Palatable formulation
- Rat & House Mouse Killer
- (Farnam®) Rat and House Mouse Control
- Can be used in bait stations
- Features nibble ridges rats and house mice love to chew
- Features nibble ridges to promote rat and house mouse chewing
- Multi-edge shape gives rats and house mice gnawing edges
- 2 oz. Bait Chunks for bait stations
- 2 oz. Bar pieces for bait stations
- Bars break into 2 oz. pieces
- 2 oz. Mini-Bars for bait stations
- The hole through the middle allows for mounting in bait stations
- Helps Provides (effective) control
- Helps Control rats and house mice

- Helps provide Effective control of rats and house mice
- (Bait Chunks) (Bars)
- Individually wrapped 1 pound Bars (helps to keep product fresh.)
- (Economical) Value (Pack) (Size) contains (16, 17, 18, 19, 20, 25) pounds of product!
- Great value! Contains (16, 17, 18, 19, 20, 25) pounds of product!

#### **ACTIVE INGREDIENT:**

Bromadiolone	0.005%
OTHER INGREDIENTS*:	99.995%
TOTAL:	

<sup>\*</sup>Contains denatonium benzoate

## KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional precautionary statements

EPA Reg. No. 270-373

EPA Est. No. 75806-AZ-1

#### **NET CONTENTS:**

16 lbs (128, 2 oz chunks) or 17 lbs (136, 2 oz lbs) or 18 lbs (144, 2 oz chunks) or 19 lbs (152, 2 oz chunks) or 20 lbs (160, 2 oz chunks) or 21 lbs (168, 2 oz chunks) or 22 lbs (176, 2 oz chunks) or 23 lbs (184, 2 oz chunks) or 24 lbs (192, 2 oz chunks) or 25 lbs (200, 2 oz chunks) or 16 lbs (16, 16 oz bars) or 17 lbs (17, 16 oz bars) or 18 lbs (18, 16 oz bars) or 19 lbs (19, 16 oz bars) or 20 lbs, (20, 16 oz bars) or 21 lbs (21, 16 oz bars) or 22 lbs (22, 16 oz bars) or 23 lbs (23, 16 oz bars) or 24 lbs (24, 16 oz bars) or 25 lbs (25, 16 oz bars)

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Keep away from children, domestic animals and pets. Causes moderate eye irritation. Do not get in eyes or on clothing. All handlers (including applicators) must long-sleeved shirt and pants, shoes plus socks, and gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.

<sup>\*</sup>This product has been treated with S-methoprene to protect the bait from infestation by insects.

#### **User Safety Requirements**

Follow manufacturer's instruction for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco, or using the toilet and change into clean clothing.

FIRST AID		
Have label with you when obtaining treatment advice.		
IF SWALLOWED:	<ul> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> </ul>	
	• Do not give anything by mouth to an unconscious person.	
IF IN EYES:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes.	
	• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.	
	• Call a poison control center or doctor for treatment advice.	

#### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-234-2269 for emergency medical treatment information.

See [side][back] panel for additional precautionary statements.

#### NOTE TO PHYSICIAN

If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin  $K_1$  intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

#### TREATMENT FOR PET POISONING

If animal eats bait, call veterinarian or 1-877-xxx-xxxx [optional] at once.

#### NOTE TO VETERINARIAN

Anticoagulant Bromadiolone: For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K<sub>1</sub>. If needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood-transfusions-may-be-needed.

#### **ENVIRONMENTAL HAZARDS**

This product is extremely toxic to mammals and birds. Dogs, cats, and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. This pesticide is toxic to fish. Do not apply this product directly to

water or to areas where surface water is present or to intertidal areas below the main high water mark. Runoff also may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash water or rinsate.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its label.

READ THIS LABEL: READ THIS ENTIRE LABEL AND USE STRICTLY IN ACCORDANCE WITH USE DIRECTIONS, USE RESTRICTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. USE ONLY FOR THE SITES, PESTS, AND APPLICATION METHODS DESCRIBED ON THIS LABEL.

It is illegal to sell these place packs individually.

**IMPORTANT:** Do not expose children, pets or other nontarget animals to rodenticides. To help prevent accidents:

- 1. Store product not in use in a location out of reach of children and pets.
- 2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears or other potentially destructive animals, or in areas prone to vandalism.
- 3. Dispose of product container, unused, spoiled and unconsumed bait as specified on this label.

**Note:** Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used if children, pets, non-target mammals, or birds may access the bait.

Do not apply this product in a way that will contact workers or other persons. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

USE RESTRICTIONS: For control only of Norway Rats, Roof Rats and House Mice in and around industrial, commercial, and public buildings. This product may

also be used inside transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. Do not apply further than 50 feet from buildings. Do not use in sewers. Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food. Do not broadcast bait. When used in USDA inspected facilities, this product must be applied in tamper-resistant bait stations.

Do not sell this product in individual containers holding less than 16 pounds of bait.

#### SELECTION OF TREATMENT AREAS

Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rats and/or mice or their signs have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

#### APPLICATION DIRECTIONS

#### APPLICATION DIRECTIONS – BAR

Each [name] Bait Bar weighs one pound and is scored to be broken into 2-oz pieces.

RATS: While wearing gloves, apply two to eight 2-oz pieces of bar (or one entire BAR) at each placement location in infested areas. For burrow applications, insert one to two 2-oz pieces of bar at least six inches into each active burrow. Check treated burrows each morning and evening to make sure that baits have not been pushed out of burrows. For aboveground treatment, space bait placements 15 to 30 feet apart. Maintain an uninterrupted supply of fresh bait for 10 days or until fresh signs of rat activity cease to appear.

MICE: While wearing gloves, break bar into 2 oz pieces and apply one 2-oz piece per placement in infested areas. Space placements at intervals of 8-12 feet. Maintain an uninterrupted supply of fresh bait for 15 days or until fresh signs of mouse activity cease to appear.

**FOLLOW-UP**: Replace contaminated or spoiled bait immediately. Wearing gloves, collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, eliminate sources of rodent food, water and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

#### APPLICATION DIRECTIONS - BAIT CHUNKS

Each [name] weighs 2-oz.

**RATS:** While wearing gloves, apply two to eight 2-oz blocksat each placement location in infested areas. For burrow applications, insert one to two 2-oz pieces of block at least six inches into each active burrow. Check treated burrows each morning and evening to make sure that baits have not been pushed out of burrows. For aboveground treatment, space placements 15 to 30 feet apart in infested areas. Maintain an uninterrupted supply of fresh bait for 10 days or until fresh signs of rat activity cease to appear.

MICE: While wearing gloves, apply one 2-oz chunk per placement in infested areas. Space placements at 8 to 12 foot intervals. Maintain an uninterrupted supply of fresh bait for 15 days or until fresh signs of mouse activity cease to appear.

**FOLLOW-UP:** Replace contaminated or spoiled bait immediately. Wearing gloves, collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

**PESTICIDE STORAGE:** Store only in original container, in a cool, dry place inaccessible to children and pets.

**PESTICIDE DISPOSAL:** Dispose of wastes resulting from the use of this product in trash or at an approved waste disposal facility.

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. [Plastic:] Triple rinse (or equivalent), then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or by incineration. In most states, burning is not allowed. [Paper:] Dispose of empty container by placing in trash, at an approved waste disposal facility or by incineration.

#### **KEEP AWAY FROM FOODSTUFFS**

#### CONDITIONS OF SALE

To the extent consistent with applicable law FARNAM COMPANIES, INC., warrants this product in its unopened package conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions as specified on this label. There are no other warranties, expressed or implied, concerning the use of this product other than as indicated on the label. To the extent consistent with applicable law, this warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or conditions not reasonably foreseeable to seller, and buyer assumes all risk of any such use.

Manufactured for:

FARNAM COMPANIES, INC.

P.O.Box 34820 • Phoenix, AZ 85067-4820

www.farnam.com

Call with comments or questions: 1-800-234-2269

© [DATE] Farnam Companies, Inc.

#### **NEW LABEL**

[Front Label]

AGRICULTURAL USE INDIVIDUAL POUCH LABEL BROMADIOLONE BAIT BAR

# [ABN: JUST ONE BITE\* II BAR] [ABN: JUST ONE BITE\* II BAIT CHUNKS]

\*Norway rats and house mice may consume a lethal dose in a single night's feeding with first dead rodents appearing four or five days after feeding begins

For Use In and Around Agricultural Buildings Only

For indoor and outdoor rat and house mouse control

Kills Norway rats, roof rats, house mice and warfarin resistant Norway rats

#### **ACTIVE INGREDIENT:**

Bromadiolone	0.005%
OTHER INGREDIENTS*:	99.995%
TOTAL:	100.000%

<sup>\*</sup>Contains denatorium benzoate

EPA Reg. No. 270-373

EPA Est. No. 75806-AZ-001

#### KEEP OUT OF REACH OF CHILDREN

**CAUTION** 

Net Contents: 16 ozs. (453.6 grams) Manufactured for:

Farnam Companies, Inc.

P.O. Box 34820 • Phoenix, AZ 85067-4820

www.farnam.com

Call with comments or questions:

1-800-234-2269

<sup>\*</sup>This product has been treated with S-methoprene to protect the bait from infestation by insects.

[Back Label)

# PRECAUTIONARY STATEMENTS Hazards To Humans And Domestic Animals

**CAUTION:** Harmful if swallowed. Keep away from children, domestic animals and pets. Causes moderate eye irritation. Do not get in eyes or on clothing. All handlers (including applicators) must wear long-sleeved shirt and long pants, shoes plus socks, and gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.

#### **User Safety Requirements**

Follow manufacturer's instruction for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco, or using the toilet and change into clean clothing.

FIRST AID  Have label with you when obtaining treatment advice		
IF SWALLOWED:	<ul> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li> </ul>	
IF IN EYES:	<ul> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>	

#### NOTE TO PHYSICIAN

If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin  $K_1$  intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

#### TREATMENT FOR PET POISONING

If animal eats bait, call veterinarian or 1-877-xxx-xxxx [optional] at once.

#### NOTE TO VETERINARIAN

Anticoagulant Bromadiolone: For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin  $K_1$ . If needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.

**ENVIRONMENTAL HAZARDS:** This product is extremely toxic to mammals and birds. Dogs, cats and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. This pesticide is toxic to fish. Do not apply this product directly to water or areas where surface water is present or to intertidal areas below the mean high water mark. Runoff also may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash water or rinsate.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL AND USE STRICTLY IN ACCORDANCE WITH USE DIRECTIONS, USE RESTRICTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. REFER TO THE OUTER LABEL FOR COMPLETE DIRECTIONS FOR USE.

It is illegal to sell these bait bars individually.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

**PESTICIDE STORAGE:** Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

**PESTICIDE DISPOSAL:** Dispose of wastes resulting from the use of this product in trash or at an approved waste disposal facility.

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. [Plastic:] Triple rinse (or equivalent), then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or by incineration. In most states, burning is not allowed. [Paper:] Dispose of empty container by placing in trash, at an approved waste disposal facility or by incineration.

#### **CONDITIONS OF SALE**

To the extent consistent with applicable law, FARNAM COMPANIES, INC. warrants this product in its unopened package conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions as specified in the label. There are no other

warranties, expressed or implied, concerning the use of this product other than as indicated on the label. To the extent consistent with applicable law, this warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or conditions not reasonably foreseeable to seller, and buyer assumes all risk of any such use.