


275-18

4-13-73

1 of 3

Main Panel
1/2 lb. carton
1/10/73



Active Ingredients:

Bacillus thuringiensis, Berliner, 16,000

International Units of Potency per mg.

(at least 25 billion
viable spores per gm.)*..... 3.2%

Inert Ingredients..... 96.8%

*Equivalent to 7.26 billion International Units
of potency per pound of this product.

CAUTION: KEEP OUT OF REACH OF
CHILDREN. DO NOT
RE-USE CONTAINER. DESTROY WHEN EMPTY.

E.P.A. Registration No. 275-18-AD-50546
Lot No.




AGRICULTURAL AND VETERINARY
PRODUCTS DIVISION
ABBOTT LABORATORIES, NORTH CHICAGO, ILL. 60064

Net contents: one-half pound

ACCEPTED

APR 13 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 275-18 SUBJECT
TO ATTACHED COMMENTS.



Side Panel
1/2 lb. carton
Revised 3/26/73

DIPEL for Food Crops

Crop	Pest	Pound/Acre
Broccoli, Cabbage, Cauliflower, Collards, Kale, Lettuce, Mustard and Turnip Greens, Spinach	Cabbage Looper	1/2 to 1
	Imported Cabbageworm	1/4 to 1/2
	Diamondback Moth	1/4 to 1/2
Beans, Cucumbers, Melons, Potatoes	Cabbage Looper	1/2 to 1
Celery	Celery Looper and Cabbage Looper	1/2 to 1
Tomatoes	Cabbage Looper	1/2 to 1
	Tomato Hornworm	1/4 to 1/2
	Tomato Fruitworm	1*

*Apply on Regular Spray Program every 5 - 7 days.


DIPEL is exempt from tolerance restriction on all listed crops and may be applied up to the time of harvest.

DIPEL for Other Crops

Crop	Pest	Rate
Tobacco	Cabbage Looper	1/2 to 1 lb./acre
	Budworm	1/2 to 1 lb./acre
	Hornworm	1/4 to 1/2 lb./acre
Chrysanthemums	Cabbage Looper	1/2 to 1 lb./acre or 100 gal.

DIPEL for Greenhouse Crops

Crop	Pest	Lb./100 Gallons Spray
Cabbage, Cucumbers, Lettuce, Tomatoes	Cabbage Looper	1/2 to 1
Chrysanthemums	Cabbage Looper	1/2 to 1





General Directions for Use

DIPEL may be used in conventional ground spray equipment with suitable quantities of water to provide thorough coverage. Fill the spray tank half full of water. Slowly sprinkle the required amount of DIPEL into the tank while agitating. Then, agitate thoroughly while bringing the total volume of water to the required level. A suitable spreader-sticker may be combined with DIPEL.

Apply DIPEL when pests are first noticed and repeat at intervals sufficient to maintain control. Thorough coverage on both sides of leaves is essential since the insect must ingest DIPEL. After ingestion, the larvae stop feeding and death follows. Do not allow the spray mixture to stand in the tank for more than 12 hours.

For Smaller Spray Volumes

<u>If Rate Is:</u>	<u>Use This Amount per Gallon</u>
1/4 lb./acre or 100 gals.	1/2 tsp.
1/2 lb./acre or 100 gals.	1 tsp.
1 lb./acre or 100 gals.	2 tsps.

