

275-18

4-27-76

1 of 2

APR 27 1976

ABBOTT LABORATORIES
Agricultural and Veterinary Product Div.
North Chicago, ILL 60054

Gentlemen:

Subject : DIPEL WP
EPA Reg. No. 275-18
Your application of April 8, 1976

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable, and a stamped copy is enclosed for your records.

Sincerely

Gerald Harrison
Product Manager 17
Insecticide Rodenticide Branch
Registration Division WH-567

WH-567:GHarrison:pl: WSME, rm 229:X69425

4-26-76



2 of 2

DIPEL® Biological Insecticide Wettable Powder

ACCEPTED
APR 28 1976
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 275-18

Supplemental Directions for Use of DIPEL® WP and Chlordimeform (Fundal® or Galecron®) Tank Mix* on Cotton

For the control of Tobacco Budworm or Cotton Bollworm apply 1/8 to 1/4 pound DIPEL WP (product) in combination with 1/8 to 1/4 pound chlordimeform (active ingredient) per acre.

The lower rate of DIPEL WP (1/8 lb acre) is recommended for light to moderate infestation levels in early season. Increase to the higher rates and use shorter spray intervals under heavier pressures. If an extremely heavy worm population occurs due to an interruption of spray schedule or to high egg deposition, it may be necessary to apply an insecticide with rapid knockdown and good larvicidal activity as a temporary replacement for DIPEL/chlordimeform. Once populations are reduced to light to moderate levels, DIPEL/chlordimeform sprays may be resumed.

Apply DIPEL/chlordimeform in 5 to 10 gallons of spray mix per acre with ground equipment and in 2 to 5 gallons for aerial application. Begin treatments when eggs first appear and maintain a 3 to 5 day spray schedule to achieve timely applications against repeated infestations.

Before mixing and applying a tank mix of DIPEL/chlordimeform, READ the container labels for both products for environmental and usage cautions.

To mix DIPEL and chlordimeform, slowly add required amount of DIPEL WP to the spray tank about half-filled with water. Agitate mix thoroughly until DIPEL is evenly dispersed. Add required amount of chlordimeform and maintain agitation while bringing total spray volume to the desired level.

PRECAUTIONS

Reapply DIPEL plus chlordimeform as soon as possible if rainfall occurs within 12 hours after application.

Do not apply diluted sprays which have been prepared for more than 24 hours.

EPA Reg. No. 275-18-AA-50546

Agricultural and Veterinary Products Division
Abbott Laboratories, North Chicago, Illinois 60064

*The mixture of DIPEL with chlordimeform for inhibiting the growth of lepidopterous larvae on plants is covered by Abbott U.S. patent 3,937,813.

Fundal® trademark of NOR-AM for chlordimeform

Galecron® trademark of CIBA-GEIGY for chlordimeform

