

275-18


5-10-77

10/1



# DIPEL® Biological Insecticide Wettable Powder

Supplemental Directions for Use of  
DIPEL® WP on Cotton



### APPLICATION RATE

Thorough Coverage is Essential for Good Control

CROP	PEST	APPLICATION RATE*
Cotton	Tobacco Budworm	1/2 - 1 lb/acre
	Cotton Bollworm	1/2 - 1 lb/acre

\*Dipel (Bacillus thuringiensis) wettable powder is a biological insecticide that may be used to suppress very small worms (1st and 2nd instar larvae) under a pest management program that includes close scouting. Apply just as eggs begin to hatch and reapply every 3-5 days as long as eggs and/or small larvae are present or until larvae have been reduced by an acceptable amount. If heavy worm populations occur, it may be necessary to apply an insecticide with rapid knockdown and good larvicidal activity as a replacement.

Thorough coverage of the plant terminals is essential for adequate suppression. Apply as a 5-15 gallons/acre total volume spray with ground equipment and 2-5 gallons/acre total volume spray with aerial equipment.

SEE CONTAINER LABEL FOR GENERAL DIRECTIONS FOR USE

EPA Reg. No. 275-18

Agricultural and Veterinary Products Division  
Abbott Laboratories, North Chicago, Illinois 60064

