

United States Environmental Protection Agency  
Washington, D.C. 20460

NOV 1 1993

Office of  
Prevention, Pesticides and  
Toxic Substances

Marvin H. Alphin  
ABBOTT LABORATORIES  
DR. JOSEPH C. WHITE DIRECTOR, REGULATORY  
14TH AND SHERIDAN RD.  
NORTH CHICAGO IL 60064

Subject: Label Amendment Submission of 06/23/93  
in Compliance with WPS Labeling Requirements  
EPA Reg No. 275-37  
DIPEL 2X BIOLOGICAL INSECTICIDE

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is accepted subject to the comments listed below.

Based on your certification, the Agency has accepted only those changes to your labeling which are necessary to comply with PR Notices 93-7 and 93-11, which reflect the WPS labeling requirements of 40 CFR part 156, subpart K. Any other labeling changes submitted in connection with this amendment application and not directly related to compliance with PR Notice 93-7 or 93-11 have neither been reviewed nor accepted by the Agency. If you wish to make any such changes, you must submit a separate amendment application proposing them. If your product registration is currently suspended, acceptance of this labeling amendment does not affect the suspension in any way.

A copy of your proposed labeling stamped "Accepted with comments" is attached. Make any required changes described in the attached and send three copies of final labeling as soon as it is available to:

Document Processing Desk (FIN-LABEL)  
Office of Pesticide Programs (H-7504C)  
U.S. Environmental Protection Agency  
401 M Street SW  
Washington, DC 20460-0001


Hand or courier deliveries of final labels may be made to:

Document Processing Desk (FIN-LABEL)  
Room 266A Crystal Mall 2  
1921 Jefferson Davis Highway  
Arlington, VA 22202

Please correct the typographical errors circled on the draft before printing final labeling.

In your final labeling the "Agricultural Use Requirements" text" must be contained in a clearly separate box. This box may be set apart by a line, by another graphical device, by a different color background, or in any other way that clearly distinguishes it from surrounding text.

Sincerely,



James Tompkins, Deputy Chief  
Registration Support Branch  
Registration Division (H7505W)

3 9 22

DIPEL 2X  
BIOLOGICAL INSECTICIDE

WETTABLE POWDER  
BIOLOGICAL INSECTICIDE

Active Ingredient:

*Bacillus thuringiensis*, subsp. *kurstaki*.....6.4%w/w

Inert Ingredients:.....93.6%w/w

Total: 100.0% w/w

Potency: 32,000 International Units per mg of product  
or 14.52 billion International Units per pound  
of product.

Potency units should not be used to adjust use rates beyond those  
specified in the Directions For Use section.

KEEP OUT OF REACH OF CHILDREN  
CAUTION

EPA Registrations No. 275-37  
EPA Est. No. 33762-IA-1

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water. Get medical attention if irritation  
persists.

IF ON SKIN: Wash skin with plenty of soap and water. Get medical attention if  
irritation persists.

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid  
contact with skin, eyes, or clothing. Wash thoroughly with soap and water after  
handling.

DiPel 2X.epa  
Page 1 of 20  
06-18-93

ACCEPTED  
with COMMENTS  
by EPA Letter Dated

NOV 1 1993

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

275-37

**Personal Protective Equipment**

Some materials that are chemical-resistant to this product are listed below.

Applicators and other handlers must wear:

- \* Long-sleeved shirt and long pants
- \* Waterproof gloves
- \* Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations**

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**BEST AVAILABLE COPY**

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- \*Coveralls,
- \*waterproof gloves
- \*shoes
- \*socks.

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

**RE-ENTRY STATEMENT**

~~Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive intervals for various crops treated~~

**BEST AVAILABLE COPY**

~~with this product, consult your State Department of Agriculture for further information.~~

~~Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must include the following information:~~

~~Inform workers of the area of fields that must not be entered without protective clothing until sprays have dried. In case of accidental exposure, wash with plenty of water. If there is any irritation in eyes or on skin after washing, get medical attention.~~

~~When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:~~

~~"Area treated with DiPel 2X on (date of application). Do not enter without appropriate protective clothing until sprays have dried. In case of accidental exposure, wash with plenty of water. If there is any irritation in eyes or on skin after washing, get medical attention."~~

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**Storage:** Reclose containers of unused DiPel 2X. Store in a dry place.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**APPLICATION DIRECTIONS:**

**Days to harvest:** There are no restrictions on applying DiPel 2X up to the time of harvest.

**Sites:** DiPel 2X may be used for any labeled pest in both field and greenhouse uses.

**BEST AVAILABLE COPY**

DiPel 2X is a highly selective insecticide for use against listed caterpillars (larvae) of lepidopterous insects. Close scouting and early attention to infestations is highly recommended. Larvae must eat deposits of DiPel 2X to be affected. Always follow these directions:

- Treat when larvae are young (early instars) before the crop is extensively damaged.
- Larvae must be actively feeding on treated, exposed plant parts.
- Thorough spray coverage is needed to provide a uniform deposit of DiPel 2X at the site of larval feeding. For some crops directed drop nozzles by ground machine are required.
- Under heavy pest population pressure, use the higher label rates, shorten the spray interval, and/or raise gallonage to improve spray coverage.
- Repeat applications at an interval sufficient to maintain control, usually 3 to 14 days depending on plant growth rate, moth activity, rainfall after treating, and other factors. If attempting to control a pest with a single spray, make the treatment when egg hatch is essentially complete, but before extensive crop damage occurs.
- A spreader-sticker which has been approved for use on growing and harvested crops should be added for hard-to-wet crops such as cole crops, or to improve weather-fastness of the spray deposits.
- DiPel 2X is a non-restricted use pesticide and does not require a restricted use permit for purchase and use.

After eating a lethal dose of DiPel 2X, larvae stop feeding within the hour, and will die within several days. Dying larvae move slowly, discolor, then shrivel, blacken and die.

DiPel 2X may be applied in conventional ground or aerial equipment with quantities of water sufficient to provide thorough coverage of infested plant parts. The amount of water needed per acre will depend on crop size, weather, spray equipment, and local experience. Unless otherwise indicated, use at least 2 gallons of water per acre by air; except in the Western U.S., where 5 to 10 gallons is the usual minimum. Add water to the spray or mixing tank at the level that provides maximum agitation. With the agitator running, slowly sprinkle in the DiPel 2X. Continue agitation. Then add

8 7 22

other spray materials (if any). Add the balance of the water and agitate until mixed. Maintain the suspension while loading and spraying. Do not mix more DiPel 2X than can be used in a 12-hour period.

**For Smaller Spray Volumes:**

<u>If Rate is</u>	<u>Use This Amount Per Gallon</u>
1/4 lb./acre or 100 gals.	1/2 tsp.
1/2 lb./acre or 100 gals.	1 tsp.
1 lb./acre or 100 gals.	2 tsps.
2 lb./acre or 100 gals.	4 tsps.



## **CHEMIGATION USE DIRECTIONS**

Chemigation directions apply only to the state of Florida and to the following crop categories: Flowers, bedding plants, ornamentals, greenhouse/shade house and outdoor nursery crops. Refer to these label sections under **General Instruction** for application rate information when chemigation is used.

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation systems. Do not connect an irrigation system (including greenhouse systems) used for pesticide applications to a public water system.

### **Spray Preparation**

First prepare a suspension of DiPel 2X in a mix tank. Fill tank with 1/2 to 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Add the required amount of DiPel 2X, and then the remaining volume of water. Then set the sprinkler to deliver a minimum of 0.1 to 0.3 inch of water per acre. Start sprinkler and uniformly inject the suspension of DiPel 2X into the irrigation water line so as to deliver the desired rate per acre. The suspension of DiPel 2X should be injected with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing. Any questions on calibration should be directed to your State Extension Service Specialists, to equipment manufacturers or other experts.

**NOTE:** When treatment with DiPel 2X has been completed, further field irrigation over the treated area should be avoided for 24 to 48 hours to prevent washing the material off the crop.

## **GENERAL PRECAUTIONS FOR APPLICATIONS THROUGH SPRINKLER IRRIGATION SYSTEMS**

Maintain continuous agitation in the mix tank during mixing and application to insure a uniform suspension.

Greater accuracy in calibration and distribution will be achieved by injecting a larger volume for a more dilute solution per unit time.

Crop injury, lack of effectiveness, or illegal residues in the crop can result from nonuniform distribution of treated water. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise. Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

**CHEMIGATION USE DIRECTIONS (continued)**

Do not apply when wind speed favors drift, when system connections or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled and drained.

**GENERAL INSTRUCTIONS**

**APPLICATION RATES**

<b>Crop Group (Typical Crops)</b>	<b>PESTS</b>	<b>POUNDS/ACRE</b>
<b>Root and Tuber</b> such as Carrots, Potatoes, Beets and Sugarbeets <b>Bulb</b> such as Onions (green and bulb) and Garlic <b>Leafy and Cole Crops</b> such as Lettuce (head and leaf), Kale, Celery, Spinach, Broccoli, Cabbage, Mustard Greens, Brussels Sprouts, Cauliflower, Collards, Chinese Cabbage, Endive, Kohlrabi and Parsley	Loopers	1/2 - 1
	Omnivorous Leafroller	1/2 - 1
	Hornworms	1/2 - 1
	Imported Cabbageworm	1/2 - 1
	Diamondback Moth	1/2 - 1
	Green Cloverworm	1/2 - 1
	Webworm	1/2 - 1
	Saltmarsh Caterpillar	1/2 - 1
	Armyworms*	1/2 - 2
	Cutworms	1/2 - 1
<b>Fruiting Vegetables</b> such as Tomatoes, Peppers and Eggplant	Loopers	1/2 - 1
	Hornworm	1/2 - 1
	Tomato Fruitworm	1/2 - 1
	Variiegated Cutworm	1/2 - 1
	Saltmarsh Caterpillar	1/2 - 1
	Armyworms*	1/2 - 2
	Pinworm	1 - 2
<b>Curcubit Vegetables</b> such as Melons, Cucumbers and Squash	Loopers	1/2 - 1
	Melonworms	1/2 - 1
	Rindworm complex	1/2 - 1
	Armyworms*	1/2 - 2

**APPLICATION RATES (continued)**

\*DiPel 2X may be used to control small armyworms (1st and 2nd instar) when populations are light and full coverage sprays are applied. Repeat treatment as necessary. If mature worms or heavy populations are present a contact insecticide should be used to enhance control.

<b>Crop Group (Typical Crops)</b>	<b>Pests</b>	<b>Pounds/Acre</b>
<b>Legume Vegetables such as Beans, Peas, Lentils and Soybeans</b>	Loopers	1/2 - 1
	Soybean Looper	1/2 - 1
	Green Cloverworm	1/2 - 1
	Velvetbean Caterpillar	1/2 - 1
	Armyworms*	1/2 - 2
	Podworms*	1/2 - 1
<b>OTHER CROPS:</b>		
Tobacco	Tobacco Budworm	1/2 - 1
	Hornworms	1/4 - 1/2
	Loopers	1/2 - 1
Safflower	Loopers	1/2 - 1
	Saltmarsh Caterpillar	1/2 - 1
	Armyworms*	1/2 - 2
Sunflowers	Loopers	1/2 - 1
	Head Moth	1/2 - 1
Peanuts	Loopers	1/2 - 1
	Velvetbean Caterpillar	1/2 - 1
	Green Cloverworm	1/2 - 1
	Podworms*	1/2 - 1

\*DiPel 2X may be used to control small armyworms and/or podworms (1st and 2nd instar) when populations are light and full coverage sprays are applied. Repeat treatment as necessary. If mature worms or heavy populations are present a contact insecticide should be used to enhance control.

**OTHER CROPS (CONTINUED)**

<b>CROP</b>	<b>PEST</b>	<b>POUND/ACRE</b>
Alfalfa (Hay and Seed) Hay and Other Forage Crops	Loopers	1/2 - 1
	Alfalfa Caterpillar	1/2 - 1
	European Skipper (Essex Skipper)	1/2 - 1
	Armyworms*	1/2 - 2
Cotton	Tobacco Budworm**	1/2 - 2
	Cotton Bollworm**	1/2 - 2
	Loopers	1/2 - 1
	Saltmarsh Caterpillar	1/2 - 1
	Armyworms*	1/2 - 2

\*\* Use to control light to moderate populations of newly hatched worms in integrated pest management conditions. Repeat treatments at 4 to 5-day intervals as long as necessary and results are acceptable. Use in combination with ovicidal rates of labeled Heliothis ovicides.

<b>CROP</b>	<b>PEST</b>	<b>POUND/ACRE</b>
Avocado	Amorbia Moth	1/2 - 2
	Omnivorous Leafroller	1/2 - 2
	Omnivorous Looper	1/2 - 2
	Orange Tortrix	1/2 - 2
	Spanworm	1/2 - 2
Malanga	Saltmarsh Caterpillar	1/2 - 1
	Armyworms*	1/2 - 2
Water Cress	Loopers	1/2 - 1
	Armyworms*	1/2 - 2
	Diamondback Moth	1/2 - 1

\*DiPel 2X may be used to control small armyworms (1st and 2nd instar) when populations are light and full coverage sprays are applied. Repeat treatment as necessary. If mature worms or heavy populations are present a contact insecticide should be used to enhance control.

**OTHER CROPS (continued)**

<b>CROP</b>	<b>PEST</b>	<b>POUND/ACRE</b>
Kiwi Fruit	Omnivorous Leafroller	1/2 - 2
Hops	Loopers Armyworms*	1/2 - 1 1/2 - 2
Bananas	Banana Skipper	1/2 - 1
Asparagus	Armyworms*	1/2 - 2
Corn (Sweet and Field and Sorghum)	Armyworms* Headworms	1/2 - 2 1/2 - 1
Rape	Looper Armyworms* Heliothis	1/2 - 1 1/2 - 2 1/2 - 2
Herbs, Spices and Mints such as Basil, Chives, Dill, Leeks and Peppermint	Looper Saltmarsh Caterpillar Armyworms*	1/2 - 1 1/2 - 1 1/2 - 2

\*DiPel 2X may be used to control small armyworms (1st and 2nd instar) when populations are light and full coverage sprays are applied. Repeat treatments as necessary. If mature worms or heavy populations are present a contact insecticide should be used to enhance control.

<b>CROP</b>	<b>PEST</b>	<b>POUND/ACRE</b>
Pineapple	Gummosos-Batrachedra comosae (Hodges) Thecla-Thecla basilides (Geyr)	1/4 - 1/2
Turf	Sod Webworm	1 - 2

**OTHER CROPS (continued)**

CROP	PEST	POUND/ACRE
Flowers, Bedding Plants and Ornamentals** (Note: Aerial application should be applied in a minimum of 5 gallons per acre)	Loopers Tobacco Budworm Omnivorous Looper Omnivorous Leafroller Diamondback Moth Armyworms* Ello Moth (Hornworm) Io Moth Oleander Moth Azalea Caterpillar	1/4 - 1/2 1/4 - 1/2 1/4 - 1/2 1/4 - 1/2 1/4 - 1/2 1/2 - 2 1/4 - 1/2 1/4 - 1/2 1/4 - 1/2 1/4 - 1/2
Greenhouse/Shadehouse and Outdoor Nursery Crops** such as Leafy, Herbs, Brassica and Fruiting groups	Loopers Heliothis	1/2 - 1 1/2 - 2

\*DiPel 2X may be used to control small armyworms (1st and 2nd instar) when populations are light and full coverage sprays are applied. Repeat treatment as necessary. If mature worms or heavy populations are present a contact insecticide should be used to enhance control.

\*\*Chemigation applications only in the state of Florida for flowers, bedding plants, ornamentals, greenhouse/shadehouse and outdoor nursery crops.

## DiPel 2X for Small Fruits

CROP	PEST	POUND/AC RE
Small Fruit and Berries such as Grapes, Strawberries, Blackberries and Cranberries	Grapeleaf Skeletonizer (ground only)	1/2 - 1
	Grape Leafroller	1/2 - 1
	Achema Sphinx Moth (Hornworm)	1/2 - 1
	Saltmarsh Caterpillar (ground only)	1/2 - 1
	Omnivorous Leafroller (ground only)	1/2 - 1
	Loopers	1/2 - 1
	Orange Tortrix	1/2 - 1
	Oblique Banded Leafroller	1/2 - 1
	Armyworms*	1/2 - 2
	Tobacco Budworm	1/2 - 2
Grape Berry Moth	1/2 - 1	
Small Grains	Loopers	1/2 - 1
	Armyworms*	1/2 - 2

\*DiPel 2X may be used to control small armyworms (1st and 2nd instar) when populations are light and full coverage sprays are applied. Repeat treatment as necessary. If mature worms or heavy populations are present a contact insecticide should be used to enhance control.



**DiPel 2X for Small Fruits (continued)**

**DiPel 2X for Tree Fruits, Nuts, Citrus and Tropical Fruit**

<b>CROP</b>	<b>PEST</b>	<b>POUND/ACRE</b>
<b>Stone Fruit</b> such as Cherries, Plum, Peach, Prune and Nectarine <b>Pone Fruit</b> such as Apples and Pears <b>Tree Nuts</b> such as Almonds, Pecan, Walnut and Filbert <b>Pomegranates</b>	Redhumped Caterpillar	1/2 - 2
	Tent Caterpillars	1/2 - 2
	Omnivorous Leafroller	1/2 - 2
	Fall Webworm	1/2 - 2
	Walnut Caterpillar	1/2 - 2
	Cankerworms	1/2 - 2
	Gypsy Moth	1/2 - 2
	Variegated Leafroller	1/2 - 2
	Redbanded Leafroller	1/2 - 2
	Tufted Apple Budworm	1/2 - 2
	Fruittree Leafroller	1/2 - 2
	Fall Webworm	1/2 - 2
	Wainut Caterpillar	1/2 - 2
	Filbert Leafroller	1/2 - 2
	Obliquebanded Leafroller	1/2 - 2
Codling Moth	1/2 - 2	
Cutworms	1/2 - 2	
Citrus	Fruittree Leafroller	1/2 - 2
	Orangedog	1/4 - 1
	Citrus Cutworm*	1/2 - 2

\*Apply to light to moderate populations of newly-hatched worms

<b>CROP</b>	<b>PEST</b>	<b>POUND/ACRE</b>
Tropical Fruits	Hornworm	1/2 - 2
	Leafrollers	1/2 - 2
	Omnivorous Looper	1/2 - 2
	Loopers	1/2 - 2

**APPLICATION RATE (continued)**

**DiPel 2X for Trees and Forests**

<b>CROP</b>	<b>PEST</b>	<b>LBS/100 GALLONS* (GROUND EQUIPMENT)</b>	<b>LBS/ACRE APPLICAT</b>
Forest, Shade, Sugar Maple Trees and Ornamentals	Gypsy Moth	1/4 - 3/4	1/2 - 1 1/2
	Bagworm	1/4 - 3/4	1/2 - 1 1/2
	Redhumped Caterpillar	1/4 - 3/4	1/2 - 1 1/2
	Spring & Fall Cankerworm	1/4 - 3/4	1/2 - 1 1/2
	Fall Webworm	1/4 - 3/4	1/2 - 1 1/2
	Elm Spanworm	1/4 - 3/4	1/2 - 1 1/2
	Tent Caterpillars	1/4 - 3/4	1/2 - 1 1/2
	California Oakworm	1/4 - 3/4	1/2 - 1 1/2
	Pine Butterfly	1/4 - 3/4	1/2 - 1 1/2
	Spruce Budworms	1/4 - 3/4	1/2 - 1 1/2
	Saddle Prominent Caterpillar	1/4 - 3/4	1/2 - 1 1/2
	Douglas Fir Tussock Moth	1/4 - 3/4	1/2 - 1 1/2
	Western Tussock Moth	1/4 - 3/4	1/2 - 1 1/2
	Fruittree Leafroller	1/4 - 3/4	1/2 - 1 1/2
	Blackheaded Budworm	1/4 - 3/4	1/2 - 1 1/2
	Mimosa Webworm	1/4 - 3/4	1/2 - 1 1/2
Jack Pine Budworm	1/4 - 3/4	1/2 - 1 1/2	
Saddleback Caterpillar	1/4 - 3/4	1/2 - 1 1/2	
Greenstriped Mapleworm	1/4 - 3/4	1/2 - 1 1/2	

\*Rate for hydraulic sprayer. For mist blowers, mix the applicable amounts (lbs.) in 10 gallons of water.

**\*\*For aerial application, use in one to five gallons of water depending on type and density of trees. For best results, spray systems which deliver droplet size of LESS THAN 150 microns should be used.**

**DIPel 2X FOR STORED AGRICULTURAL COMMODITIES**

**GRAINS, SOYBEANS, SUNFLOWER SEED, CROP SEED, CONDIMENTAL SEEDS, SPICES, HERBS, BIRDSEED<sup>1</sup> AND POPCORN<sup>1</sup>**

PEST	RATE
Indian Meal Moth Almond Moth	3/8 lb. 100 bu (undiluted and diluted)*

**\*As a surface treatment, apply 1/2 lb. DiPel 2X in 5-10 gal. of water per 500 sq. ft. of grain surface area, mix into top 4 inches.\*\***

<sup>1</sup>For all States except California

For the control and prevention of these pests, apply DiPel 2X in a constantly agitated water suspension to the top four inch surface layer of grain in the bin. Use a sprinkler can or sprayer to apply the dosage into the grain stream as the last (top) four inch layer is augered into the bin. Mix 1/20 lb. DiPel 2X per gallon of water. Apply 0.6 pint of this mixture per bushel as grain is augered into storage. Or, sprinkle the dosage into the surface of the grain in the bin and mix thoroughly with a scoop or rake to the depth of four inches. More thorough coverage may be achieved by dividing the recommended dosage into three applications and mixing the grain between applications.

19  
9  
22

For the protection of bagged grain including popcorn, apply the dosage to the entire grain mass and mix thoroughly prior to bagging.

Treatments can be applied to stored grain at any time, but for best results, make application immediately after harvest before moth activity occurs. In areas where late fall harvested grain is not subject to infestation because of low temperatures, application can be delayed until late winter or early spring before moth activity begins. Control for a full storage season should normally be expected; however, repeat application if infestation recurs.

This treatment controls the moth larvae. If an infestation is present when the grain is treated, moth emergence may continue for several days. If immediate control of severe infestations is desired, grain should be fumigated prior to application of this treatment. **DiPel 2X will not control weevils or other beetles.**

Grain treated with DiPel 2X can be used at any time after treatment for any use.

**\*\*For commodities coarser than shelled corn, increase depth of treatment according to the habit of the pest.**

## **PEANUTS**

### **PEST**

### **RATE**

Indian Meal Moth,  
Almond Moth

1/4 lb./ton\*

**\*Apply this rate to the top four to eight feet of nuts when filling the warehouse.**

To prevent and control these pests, spray an even coating of DiPel 2X on the farmer stock peanuts while filling the warehouse. To make the spray solution, mix 3-3/4 lbs. DiPel 2X per 5 gallons of water. Apply to 15 tons of commodity. Do not pre-mix more

**APPLICATION RATES (Continued)**

**DIPel 2X FOR STORED AGRICULTURAL COMMODITIES (Continued)**

**GRAINS, SOYBEANS, SUNFLOWER SEED, CROP SEED, CONDIMENTAL SEEDS, SPICES, HERBS, BIRDSEED<sup>1</sup> AND POPCORN<sup>1</sup>. (Continued)**

spray solution than will be used within 12 hours. Keep the spray suspension agitated during application, and use pressures and nozzles sufficient to handle this suspension.

Before filling the warehouse, clean the facility, then spray interior of the facility with a DiPel 2X suspension at the rate of 1/2 oz. DiPel 2X per 100 gallons water. Spray enough suspension to wet all cracks and crevices.

For bagged peanuts, treat the whole mass of commodity at the rate indicated above.

**FLUE-CURED TOBACCO**

<b>PEST</b>	<b>RATE</b>
Tobacco Moth	0.2 oz./100 lbs*

\*Apply 0.2 ounce (approximately 2 1/2 teaspoonsful) of DiPel 2X in one quart of water per 100 pounds of tobacco as a fine mist spray. Avoid oversetting. Tobacco should have just enough moisture to be handled without shattering at the time of application.

**Tobacco to be Stored up to Twelve Months.**

Spray loose leaves as the tobacco is being bundled from the curing barn. For tobacco on sticks, treat both sides of leaves.

**Stored Tobacco.**

For tobacco which is to be carried over, rebundle or restack sticks, fluff up tobacco and spray loose leaves.

For tobacco that has been stored over three weeks, apply at first signs of infestation promptly open bundles, spray loose leaves, then bundle.

**Treatment of Storage Barns.**

If tobacco has been treated, or is going to be treated, treatment of the floors and walls may be made to aid in control. Sweep out the area, especially cracks and corners,

and all of the loose tobacco pieces in which the moth might breed. Make a spray mixture containing 1/2 oz. DiPel 2X per 2 1/2 gallons of water. Apply this at a rate of 1/2 gallon per 1000 sq. ft. of surface area. Be sure to spray into cracks and between floorboards.

**NOTICE TO USER**

Seller makes no warranty, express or implied, of merchantability, fitness or otherwise concerning the use of this product other than as indicated on the label. User assumes all risks of use, storage or handling not in strict accordance with accompanying directions.

(C) 1992  
Chemical and Agricultural Products Division  
Abbott Laboratories  
North Chicago, Illinois 60064