

MYCAR is a biological acaricide for suppression of susceptible mite populations on citrus and other crops. The active ingredient *H. thompsonii* is a naturally occurring, highly specific fungal pathogen of certain mite species. The fungal conidia attach to and penetrate the mite cuticle. Death results in 48 hours. The fungus grows in the host and releases new conidia onto the leaf and fruit surface where they are available for infection of other mites. In this way a residual effect is provided until the mite populations are reduced.

PRECAUTIONARY STATEMENTS

Hazard to Humans & Domestic Animals — Avoid contact with eyes. In case of contact, immediately flush eyes with plenty of water. Get medical attention if irritation persists.

Environmental Hazard — Keep out of lakes, ponds and streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE — Keep refrigerated.

DISPOSAL — Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

BENEFICIAL INSECTS MYCAR has not been found to be harmful to predacious and parasitic insects and mites in citrus groves.

Lot No

Not for sale or use after

Not registered in California.

Chemical and Agricultural Products Division
Abbott Laboratories
North Chicago, Illinois 60064



LIST 5533
01

Wettable Powder

Active Ing. *Hirsutella thompsonii* 1%
3.6 x 10⁸ CFUs (Colony Forming Units) of *Hirsutella thompsonii* per pound (This is equivalent to 0.8 x 10⁶ CFUs per gram) 1%
Inert Ingredients 99%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panel for precautionary statements and directions for use.

EPA Registration No 275-38
EPA Est. No. 275-IL-1

Net Contents: 6 x 4 pounds

MIXING INSTRUCTIONS

Add the required amount of MYCAR® to a spray tank about half filled with water. Add a spreader sticker* to the tank at the manufacturer's recommended rate. Agitate while bringing the total volume of water to the required level.

*Phac® or Nufilm®

APPLICATION:

Crop	Pest	Pounds per Acre
Citrus	Citrus Rust Mite	2-4

Apply as a full coverage spray.

Apply during post-bloom period and/or summer. A summer oil may be tank mixed. At other times of the year apply when 20-40% of the leaves and/or fruit are infested with one or more mites.

Blueberries	Blueberry Mite	2-4
-------------	----------------	-----

Apply 150 gallons per acre with a hydraulic sprayer at 150-200 psi.

Apply when populations are building, normally in June and late August.

Turf	Bermuda Turf Mite	2-4
------	-------------------	-----

Apply 200-300 gallons per acre.

Apply when populations are building, normally in late spring and early summer.

APPLICATION NOTES AND RESTRICTIONS

Do not apply MYCAR

- In combination with a fungicide or other pesticides (except summer oil)
- In air temperature less than 45°F or greater than 95°F

NOTICE TO USER

Seller makes no warranty, express or implied of merchantability, fitness or otherwise concerning the use of this product other than as indicated on the label. User assumes all risks of use, storage or handling not in strict accordance with accompanying directions.

*Phac and Nufilm are trademarks of companies other than Abbott Laboratories.

01-2854-2/81