

1020
pm 7
275-41

NOT REGISTERED IN CALIFORNIA

WM 12-4-84

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
(and Domestic Animals)**

ACTION
Avoid contact with skin, eyes or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS
Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

Beneficial Insects
Insects foraging treated areas are not harmed by this product.
This product does not interrupt the activities of beneficial and predatory arthropods in pest management programs.


Restrictions for Use
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Refer to Package Enclosure for crops, rates and general directions.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
STORAGE: Keep containers tightly closed when not in use. At temperatures less than 0°F and greater than 100°F, DWPEL 6L should be stored under cover.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Triple rinse (or equivalent).

NOTICE TO USER
Manufacturer makes no warranty, express or implied, of merchantability, fitness or otherwise concerning use of this product other than as indicated on the label. User assumes all risks of use, storage or handling not in accordance with accompanying directions.

1 No 5383 82
1 A Registration No. 275-40
1 A Est No 33782-1A-1

 Chemical and Agricultural Products Division
Abbott Laboratories
North Chicago, IL 60064

1 No
3106-2/R2

LIQUID WORM KILLER


JAN 2 1985
275-41

EMULSIFIABLE SUSPENSION
BIOLOGICAL INSECTICIDE

Active Ingredient:
Bacillus thuringiensis, var. kurstaki, 13,200 International Units of Potency per mg (48 billion International Units per gallon) 2.6%
Inert Ingredients 97.4%

KEEP OUT OF REACH OF CHILDREN
CAUTION

NET CONTENTS: BULK TANK CAR/TRUCK



WORM KILLER
EMULSIFIABLE SUSPENSION
BIOLOGICAL INSECTICIDE

Active ingredient:
Bacillus thuringiensis, var. kurstaki,
13,200 International Units of Potency per
mg (48 billion International Units per
gallon) 2.6%
Inert ingredients 97.4%

KEEP OUT OF REACH OF CHILDREN
CAUTION

E.P.A. Registration No. 275-48
E.P.A. Est. No. 33762-1A-1

CAUTION

Avoid contact with skin, eyes or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

BENEFICIAL INSECTS

Honeybees foraging treated areas are not harmed by DIPEL 6L use.

DIPEL 6L does not interrupt the activities of beneficial and predacious arthropods in pest management programs.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

See container label.

Days to harvest: There are no restrictions on applying DIPEL 6L up to the time of harvest.

DIRECTIONS FOR USE

DIPEL 6L is a highly selective insecticide for use against listed caterpillars (larvae) of lepidopterous insects. Close scouting and early attention to infestations is highly recommended. Larvae must eat deposits of DIPEL 6L to be affected. Always follow these directions:

- Treat when larvae are young (early instars) before extensive damage has occurred.
- Larvae must be actively feeding on treated, exposed plant parts.
- Thorough spray coverage is needed to provide a uniform deposit of DIPEL 6L at the site of larval feeding.
- Under heavy pest population pressure, use the higher label rates and/or raise gallonage to improve spray coverage.
- If attempting to control a pest with a single spray, make the treatment when egg hatch is essentially complete, but before extensive crop damage occurs.
- A spreader-sticker which has been approved for use on growing and harvested crops may be added to improve weather-fastness of the spray deposits.

After eating a lethal dose of DIPEL 6L, larvae stop feeding within the hour and will die within several days. Dying larvae move slowly, discolor, then shrivel, blacken and die.

DIPEL 6L may be applied in conventional ground or aerial equipment with quantities of water sufficient to provide thorough coverage of infested plant parts. The amount of water needed per acre will depend on weather, spray equipment, and local experience. Fill the mix tank or plane hopper with the desired quantity of water. Start the mechanical or hydraulic agitation to provide moderate circulation before adding DIPEL 6L. Always add the sticker to the water prior to the addition of DIPEL 6L. Add the desired volume of DIPEL 6L to the tank or plane hopper and continue circulation. Include rinse water from containers. Never add DIPEL 6L to the mix tank before introducing the desired quantity of water. Maintain the suspension while loading and spraying. Do not mix more DIPEL 6L than can be used in a 144-hour period. **CAUTION:** Rinse and flush spray equipment thoroughly following each use.

JAN 22 1985

E.P.A. Reg. No. 275-48