

Left Panel

Vectobac®-AS is a highly selective insecticide for use against mosquito and black fly larvae.

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS
AND DOMESTIC ANIMALS

As a precautionary measure in case of contact, flush eyes with plenty of water. In case of irritation, contact a physician.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a cool place.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Do not reuse container.

Chemical & Agricultural Products Division
Abbott Laboratories
North Chicago, IL 60064

VECTOBAC®-AS
Biological Mosquito
and
Black Fly Larvicide

Aqueous Suspension

Active Ingredient: Bacillus thuringiensis, Serotype, H-14, 600 International Toxic Units (ITU) per milligram (Equivalent to 2.19 billion ITU per gallon; 0.576 billion ITU per liter) 0.6%
Inert Ingredients99.4%

KEEP OUT OF REACH OF CHILDREN

CAUTION

EPA Reg. No. 275-52

EPA Establishment No.

List No.

Lot No.

Net Volume : 55 gallons

ACCEPTED
JAN 16 1984
EPA REG. NO. 275-52

Right Panel

APPLICATION DIRECTIONS

MOSQUITOES

Habitat Irrigation ditches, roadside ditches, flood water, standing ponds, catch basins, storm water retention areas, tidal water, salt marshes and rice fields.
Suggested Range Rate 0.5 pint-1 quart/acre

*Use higher rate in deep or polluted water, and when late 3rd and 4th instar larvae predominate.

Apply recommended rate in 0.5 to 100 gallons of water per acre by ground or aerial application. Uniform coverage is necessary for best results. Reapply as needed. Do not mix more Vectobac®-AS than can be used in a 24-hour period.

BLACK FLIES

Habitat Streams (non-potable)
Suggested Range Rate 1.0 to 10 mg/liter stream water** (=ppm)

**Use higher rate when stream contains high concentration of organic material.

Apply by either conventional ground or aerial spray or with metered release for 1-15 minutes upstream from infested sites to achieve the recommended concentration. Reapply as needed.

MIXING DIRECTIONS

Shake or agitate the Vectobac®-AS before use. Fill the mix tank or plane hopper with the desired quantity of water and start the agitator. Add the Vectobac®-AS with continued agitation. Maintain the suspension while loading and spraying. Do not mix more Vectobac®-AS than can be used in a 24-hour period.

NOTICE TO USER

Seller makes no warranty, express or implied, of merchantability, fitness or otherwise concerning the use of this product other than as indicated on the label. User assumes all risks of use, storage or handling not in strict accordance with accompanying directions.