

320 / 429,221
 38 /
 337 / 429,224
 18 /
 346 / 429,227
 38 /
 345 / 412,440
 38 /
 300 / 429,240
 38 /

NOV 13 1992

306 / 429,217
 38 /
 300 / 421,220
 38 /

Mr. Marvin H. Alphin
 Registration Specialist
 Abbott Laboratories, C.A.P.D.
 1401 Sheridan Road
 North Chicago, IL 60064-4000

Dear Mr. Alphin:

Subject: Alternate Formula Specified in the Vectobac AS Confidential Statement of Formula Dated 11/28/90 and Modified Labeling Adding 1) Indoor Ornamental and Plantscapes, 2) Vegetable Crops, and 3) an Increase in the Rice Chemigation Rate EPA Registration No. 275-52

The amendments referred to above, submitted in connection with registration under section 3(c)(7)(A) of Federal Insecticide, Fungicide, and Rodenticide Act are acceptable provided that you do the following.

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) and section 4 when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling.
 - a) Replace "Bacillus thuringiensis, Serotype H-14," with "Bacillus thuringiensis subspecies israelensis" in the ingredients statement.
 - b) Replace the text in the Hazards to Humans section to read "Harmful if absorbed through skin or inhaled. Avoid contact with skin, eyes, or clothing and avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse."

CONCURRENCES

SYMBOL	H-7505C						
SUF NAME	Handwritten						
DATE	11/13/92						

c) Add the following Statement of Practical Treatment block.

STATEMENT OF PRACTICAL TREATMENT

If in Eyes: Flush eyes with plenty of water. Get medical attention.

If on Skin: Wash thoroughly with plenty of soap and water. Get medical attention.

If Inhaled: Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth to mouth. Get medical attention.

d) Modify "Vectobac AS is not phytotoxic to ornamental plant species." to read "Vectobac AS is not known to be phytotoxic to ornamental plant species."

e) Replace "Follow label directions in chemigation section when applying this product through any irrigation system." to read "Do not apply this product through any irrigation system unless labeling on chemigation is followed."

f) Move "Vectobac AS can be applied by injection into drip or overhead (sprinkler) irrigation systems." to the chemigation section.

g) Modify "Apply this product only through sprinkler, including solid set or drip (trickle) irrigation systems." to read "Apply this product only through: sprinkler including solid set; flood (basin); or drip (trickle) irrigation systems."

3. Submit an eye irritation study, using the 11/28/90 Vectobac 12AS alternate formulation, that complies with Good Laboratory Practice Regulations within 6 months of this letter.

4. Submit five (5) copies of your final printed labeling before you release your product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA Section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

CONCURRENCES							
SYMBOL							
SURNAME							
DATE							

3910

A stamped label is enclosed for your records.

Sincerely,



Phil Hutton
Product Manager 18
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosure

CONCURRENCES							
SYMBOL							
SURNAME							
DATE							

4 of 10

VECTOBAC^R AS
AQUEOUS SUSPENSION
BIOLOGICAL LARVICIDE

ACCEPTED
with COMMENTS
by EPA Label Office
NOV 13 1992
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
275-52

Active Ingredient:

<u>Bacillus thuringiensis, Serotype</u>	
H-14, 600 International Toxic Units	
(ITU) per milligram (Equivalent to 2.19	
billion ITU per gallon; 0.576 billion	
ITU per liter)	0.6%
<u>Inert Ingredients</u>	<u>99.4%</u>
<u>Total.....</u>	<u>100.0%</u>

KEEP OUT OF REACH OF CHILDREN

CAUTION

EPA Reg. No. 275-52

EPA Establishment No. 33762-IA-1

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS

As a precautionary measure in case of contact, flush eyes with plenty of water. In case of irritation, get medical attention.

PHYSICAL OR CHEMICAL HAZARDS

Diluted or undiluted Vectobac AS can cause corrosion if left in prolonged contact with aluminum spray system components. Rinse spray system with plenty of clean water after use. Care should be taken to prevent contact with aluminum aircraft surfaces, structural components and control systems. In case of contact, rinse with plenty of water. Inspect aluminum aircraft components regularly for signs of corrosion.

BEST AVAILABLE COPY

59 10

DIRECTIONS FOR USE

General Classification

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply directly to treated finished drinking water reservoirs or drinking water receptacles.

CHEMIGATION

Follow label directions in chemigation section when applying this product through any irrigation system.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a cool [59° - 86°F (15° - 30°C)], dry place.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. Do not reuse container.

APPLICATION DIRECTIONS

(Fungus Gnat Control)

Fungus Gnat Habitat

Suggested Range Rate

Indoor ornamental and plantscape use.
Ornamental and nursery plantings
in greenhouse or potting soil
mixtures.

Light infestation:
16 - 32 oz/100 gallons*
applied as a soil drench

Vegetable plants such as the
following: Tomatoes, leafy and
cole crops, cucumbers, peppers
and eggplants.

Heavy Infestation:
64 - 128 oz/100 gallons**
applied as a soil drench

* 16 - 32 oz/100 gallons = 1 - 2 tsp/gallon.

** 64 - 128 oz/100 gallons = 4 - 8 tsp/gallon.

BEST AVAILABLE COPY

Apply VectoBac AS with adequate water by soil drench to sufficiently wet the soil surface above and under benches where larvae are found. Areas under benches should be treated at high rates as this is one of the primary breeding areas. Reapply as needed. In situations where all life forms (eggs, larvae, pupae, and adults) are present, such as with existing infestations, make three (3) weekly applications at the suggested rate range for heavy infestations.

Regular follow-up applications using the suggested light infestation rates, will establish a long term maintenance program.

VectoBac AS is a larvicide and will not control adult gnats, therefore, applications must be timed for a stage of development when larvae are present in the soil.

Fungus gnat larvae generally respond to VectoBac AS treatment within 24 hours following application.

VectoBac AS can be applied by injection into drip or overhead (sprinkler) irrigation systems.

VectoBac AS is not phytotoxic to ornamental plant species. However, since all ornamental plant species have not been evaluated, sensitivity to VectoBac AS should be checked on several plants prior to wide scale usage.

PRECAUTIONS

Important: VectoBac AS should not be injected in combination with fertilizers or fungicides containing copper or chlorine, as this may neutralize the active ingredients. (Chlorine levels in potable water supplies should not present a problem with VectoBac AS performance).

Do not apply soil drenches to plants under stress, or follow application with excessive amount of water.

For best results, apply drenches toward the end of irrigation period.

APPLICATION DIRECTIONS

(Mosquito and Blackfly Control)
Suggested Rate Range*

Mosquito Habitat

(Such as the following examples):
Irrigation ditches, roadside ditches,
flood water, standing ponds, woodland pools,
snow melt pools, pastures, catch basins,
storm water retention areas, tidal water,
salt marshes and rice fields.

VectoBac AS

0.5 - 2 pts/acre

BEST AVAILABLE COPY

Polluted water
(such as sewage lagoons, animal
waste lagoons).

2 pts /acre

*Use higher range in polluted water when late 3rd and early 4th instar larvae predominate, mosquito populations are high, water is heavily polluted, and/or algae are abundant.

7710

Suggested Rate Range

Blackflies Habitat

VectoBac AS

Streams

stream water** (=ppm) for
1 minute exposure time.

1 - 50 mg/liter

stream water** (=ppm) for
10 minute exposure time.

0.1 - 5 mg/liter

**Use higher rate range when stream contains high concentration of organic materials, algae, or dense aquatic vegetation.

Ground and Aerial Application

VectoBac AS may be applied in conventional ground or aerial application equipment with quantities of water sufficient to provide uniform coverage of the target area. The amount of water needed per acre will depend on the weather, spray equipment, and mosquito habitat characteristics. Do not mix more VectoBac AS than can be used in a 72-hour period.

For most ground equipment, apply in 5 - 100 gallons per acre using hand-pump, airblast, mist blower, etc., spray equipment.

For aerial application, VectoBac AS may be applied either undiluted or dilute with water. For undiluted applications, apply 0.5 to 2.0 pts/acre of VectoBac AS through fixed wing or helicopter aircraft equipped with either conventional boom and nozzle systems or rotary atomizers.

For diluted applications, fill the mix tank or plane hopper with the desired quantity of water. Start the mechanical or hydraulic agitation to provide moderate circulation before adding the VectoBac AS. VectoBac AS suspends readily in water and will stay suspended over normal application periods. Brief recirculation may be necessary if the spray mixture has sat for several hours or longer. AVOID CONTINUOUS AGITATION OF THE SPRAY MIXTURE DURING SPRAYING.

Rinse and flush spray equipment thoroughly following each use.

For blackfly aerial applications, VectoBac AS can be applied undiluted via fixed wing or helicopter aircraft equipped with either conventional nozzle systems or open pipes. Rate of application will be determined by the stream discharge and the required amount of VectoBac AS necessary to maintain a 1 - 50 ppm concentration in the stream water. VectoBac AS can also be applied diluted with similar spray equipment. Do not mix more product than can be used in a 72-hour period.

BEST AVAILABLE COPY

8710

SMALL QUANTITY DILUTION RATES
Gallons Spray Solution/Acre
(Ounces Needed per Gallons of Spray)

VectoBac AS Rate in Pints per Acre	10 Gal/A	25 Gal/A	50 Gal/A
0.5	.8	.33	.16
1.0	1.6	.66	.32
2.0	3.2	1.30	.64

CHEMIGATION

Apply this product only through sprinkler, including solid set or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service Specialist, equipment manufacturers or other experts.

Do not connect irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

RICE - FLOOD (BASIN) CHEMIGATION

Systems using gravity flow pesticide dispensing systems must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops.

VectoBac AS is metered or dripped into the rice floodwater at application stations positioned at the point of introduction (levee cut) of water into each rice field or pan. Two pints of VectoBac AS are diluted in water to a final volume in 5 gallons. The diluted

BEST AVAILABLE COPY

9910

solution is contained in a 5 gallon container and metered or dispersed into the irrigation water using a constant flow device at the rate of 80 ml per minute.

Introduction of the solution should begin when 1/3 to 1/2 of the pan or field is covered with floodwater. Delivery of the solution should continue for a period of approximately 4-1/2 hours. Floodwater depth should not exceed 10-12 inches to prevent excessive dilution of VectoBac AS which could result in reduced larval kill. Agitation is not required during the period in which the VectoBac AS solution is being dispersed.

Application of VectoBac AS into rice floodwater is not permitted using a pressurized water and pesticide injection system.

Greenhouse - Drip (Trickle) and Sprinkler Chemigation

Public water system means a system for the provision to the public of piped water for human consumption if such system has a least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of the pesticide introduction. As an option to the RPZ the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlock controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a water pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials compatible with pesticides and capable of being fitted with a system interlock.

The system must contain a functional check valve, vacuum relief

BEST AVAILABLE COPY

10 y

valve and low pressure drain, appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

If VectoBac AS needs to be diluted prior to injection, material may be mixed separately and then put into the injector. Agitation may be necessary if materials are kept more than one day. VectoBac AS may be applied continuously.

Where supply tanks are used for continuous application, fill the supply tank with the desired quantity of water. Start mechanical or hydraulic agitation to provide moderate circulation before adding VectoBac AS. VectoBac AS suspends readily in water and will stay suspended over normal application periods. Avoid continuous agitation of spray mixture during spraying. Brief recirculation may be necessary if the spray mixture has sat for several hours.

NOTICE TO USER

Seller makes no warranty, express or implied, of merchantability, fitness or otherwise concerning the use of this product other than as indicated on the label. User assumes all risk of use, storage or handling not in strict accordance with accompanying directions.

Chemical and Agricultural Products Division
Abbott Laboratories
North Chicago, Illinois 60064

(c) 1992

PA-18

275-52

1 of 10

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

330 / 429,221
38

337 / 429,224
18

340 / 429,227
88

345 / 429,226
38

325 / 412,440
38

300 / 429,240
38

NOV 13 1992

306 / 429,217
38

300 / 421,220
38

Mr. Marvin H. Alphin
 Registration Specialist
 Abbott Laboratories, C.A.P.D.
 1401 Sheridan Road
 North Chicago, IL 60064-4000

Dear Mr. Alphin:

Subject: Alternate Formula Specified in the Vectobac AS Confidential Statement of Formula Dated 11/28/90 and Modified Labeling Adding 1) Indoor Ornamental and Plantscapes, 2) Vegetable Crops, and 3) an Increase in the Rice Chemigation Rate
 EPA Registration No. 275-52

The amendments referred to above, submitted in connection with registration under section 3(c)(7)(A) of Federal Insecticide, Fungicide, and Rodenticide Act are acceptable provided that you do the following.

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) and section 4 when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling.
 - a) Replace "Bacillus thuringiensis, Serotype H-14," with "Bacillus thuringiensis subspecies israelensis" in the ingredients statement.
 - b) Replace the text in the Hazards to Humans section to read "Harmful if absorbed through skin or inhaled. Avoid contact with skin, eyes, or clothing and avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse."

CONCURRENCES

SYMBOL	47505C						
SURNAME	[Signature]						
DATE	11/13/92						

20710

c) Add the following Statement of Practical Treatment block.

STATEMENT OF PRACTICAL TREATMENT

If in Eyes: Flush eyes with plenty of water. Get medical attention.

If on Skin: Wash thoroughly with plenty of soap and water. Get medical attention.

If Inhaled: Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth to mouth. Get medical attention.

d) Modify "Vectobac AS is not phytotoxic to ornamental plant species." to read "Vectobac AS is not known to be phytotoxic to ornamental plant species."

e) Replace "Follow label directions in chemigation section when applying this product through any irrigation system." to read "Do not apply this product through any irrigation system unless labeling on chemigation is followed."

f) Move "Vectobac AS can be applied by injection into drip or overhead (sprinkler) irrigation systems." to the chemigation section.

g) Modify "Apply this product only through sprinkler, including solid set or drip (trickle) irrigation systems." to read "Apply this product only through: sprinkler including solid set; flood (basin); or drip (trickle) irrigation systems."

3. Submit an eye irritation study, using the 11/28/90 Vectobac 12AS alternate formulation, that complies with Good Laboratory Practice Regulations within 6 months of this letter.

4. Submit five (5) copies of your final printed labeling before you release your product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA Section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

CONCURRENCES							
SYMBOL							
SURNAME							
DATE							

3 of 10

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

3

A stamped label is enclosed for your records.

Sincerely,



Phil Hutton
Product Manager 18
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosure

CONCURRENCES							
SYMBOL							
SURNAME							
DATE							

402

VECTOBAC^R AS
AQUEOUS SUSPENSION
BIOLOGICAL LARVICIDE

ACCEPTED
with COMMENTS
by EPA Laboratory
NOV 13 1992
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
275-52

Active Ingredient:

<u>Bacillus thuringiensis, Serotype</u> H-14, 600 International Toxic Units (ITU) per milligram (Equivalent to 2.19 billion ITU per gallon; 0.576 billion ITU per liter)	0.6%
<u>Inert Ingredients</u>	99.4%
<u>Total.....</u>	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

EPA Reg. No. 275-52

EPA Establishment No. 33762-IA-1

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS

As a precautionary measure in case of contact, flush eyes with plenty of water. In case of irritation, get medical attention.

PHYSICAL OR CHEMICAL HAZARDS

Diluted or undiluted Vectobac AS can cause corrosion if left in prolonged contact with aluminum spray system components. Rinse spray system with plenty of clean water after use. Care should be taken to prevent contact with aluminum aircraft surfaces, structural components and control systems. In case of contact, rinse with plenty of water. Inspect aluminum aircraft components regularly for signs of corrosion.

DIRECTIONS FOR USE

General Classification

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply directly to treated finished drinking water reservoirs or drinking water receptacles.

CHEMIGATION

Follow label directions in chemigation section when applying this product through any irrigation system.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a cool [59° - 86°F (15° - 30°C)], dry place.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. Do not reuse container.

APPLICATION DIRECTIONS

(Fungus Gnat Control)

Fungus Gnat Habitat

Suggested Range Rate

Indoor ornamental and plantscape use.
Ornamental and nursery plantings
in greenhouse or potting soil
mixtures.

Light infestation:
16 - 32 oz/100 gallons*
applied as a soil drench

Vegetable plants such as the
following: Tomatoes, leafy and
cole crops, cucumbers, peppers
and eggplants.

Heavy Infestation:
64 -128 oz/100 gallons**
applied as a soil drench

* 16 - 32 oz/100 gallons = 1 - 2 tsp/gallon.

** 64 - 128 oz/100 gallons = 4 - 8 tsp/gallon.

Apply VectoBac AS with adequate water by soil drench to sufficiently wet the soil surface above and under benches where larvae are found. Areas under benches should be treated at high rates as this is one of the primary breeding areas. Reapply as needed. In situations where all life forms (eggs, larvae, pupae, and adults) are present, such as with existing infestations, make three (3) weekly applications at the suggested rate range for heavy infestations.

Regular follow-up applications using the suggested light infestation rates, will establish a long term maintenance program.

VectoBac AS is a larvicide and will not control adult gnats, therefore, applications must be timed for a stage of development when larvae are present in the soil.

Fungus gnat larvae generally respond to VectoBac AS treatment within 24 hours following application.

VectoBac AS can be applied by injection into drip or overhead (sprinkler) irrigation systems.

VectoBac AS is not phytotoxic to ornamental plant species. However, since all ornamental plant species have not been evaluated, sensitivity to VectoBac AS should be checked on several plants prior to wide scale usage.

PRECAUTIONS

Important: VectoBac AS should not be injected in combination with fertilizers or fungicides containing copper or chlorine, as this may neutralize the active ingredients. (Chlorine levels in potable water supplies should not present a problem with VectoBac AS performance).

Do not apply soil drenches to plants under stress, or follow application with excessive amount of water.

For best results, apply drenches toward the end of irrigation period.

APPLICATION DIRECTIONS

(Mosquito and Blackfly Control)
Suggested Rate Range*

Mosquito Habitat

(Such as the following examples):
Irrigation ditches, roadside ditches,
flood water, standing ponds, woodland pools,
snow melt pools, pastures, catch basins,
storm water retention areas, tidal water,
salt marshes and rice fields.

VectoBac AS

0.5 - 2 pts/acre

Polluted water

(such as sewage lagoons, animal
waste lagoons).

2 pts /acre

*Use higher range in polluted water when late 3rd and early 4th instar larvae predominate, mosquito populations are high, water is heavily polluted, and/or algae are abundant.

70810

Suggested Rate Range

Blackflies Habitat

VectoBac AS

Streams

stream water** (=ppm) for
1 minute exposure time.

1 - 50 mg/liter

stream water** (=ppm) for
10 minute exposure time.

0.1 - 5 mg/liter

**Use higher rate range when stream contains high concentration of organic materials, algae, or dense aquatic vegetation.

Ground and Aerial Application

VectoBac AS may be applied in conventional ground or aerial application equipment with quantities of water sufficient to provide uniform coverage of the target area. The amount of water needed per acre will depend on the weather, spray equipment, and mosquito habitat characteristics. Do not mix more VectoBac AS than can be used in a 72-hour period.

For most ground equipment, apply in 5 - 100 gallons per acre using hand-pump, airblast, mist blower, etc., spray equipment.

For aerial application, VectoBac AS may be applied either undiluted or dilute with water. For undiluted applications, apply 0.5 to 2.0 pts/acre of VectoBac AS through fixed wing or helicopter aircraft equipped with either conventional boom and nozzle systems or rotary atomizers.

For diluted applications, fill the mix tank or plane hopper with the desired quantity of water. Start the mechanical or hydraulic agitation to provide moderate circulation before adding the VectoBac AS. VectoBac AS suspends readily in water and will stay suspended over normal application periods. Brief recirculation may be necessary if the spray mixture has sat for several hours or longer. AVOID CONTINUOUS AGITATION OF THE SPRAY MIXTURE DURING SPRAYING.

Rinse and flush spray equipment thoroughly following each use.

For blackfly aerial applications, VectoBac AS can be applied undiluted via fixed wing or helicopter aircraft equipped with either conventional nozzle systems or open pipes. Rate of application will be determined by the stream discharge and the required amount of VectoBac AS necessary to maintain a 1 - 50 ppm concentration in the stream water. VectoBac AS can also be applied diluted with similar spray equipment. Do not mix more product than can be used in a 72-hour period.

85710

SMALL QUANTITY DILUTION RATES
Gallons Spray Solution/Acre
(Ounces Needed per Gallons of Spray)

VectoBac AS Rate in Pints per Acre	10 Gal/A	25 Gal/A	50 Gal/A
0.5	.8	.33	.16
1.0	1.6	.66	.32
2.0	3.2	1.30	.64

CHEMIGATION

Apply this product only through sprinkler, including solid set or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service Specialist, equipment manufacturers or other experts.

Do not connect irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

RICE - FLOOD (BASIN) CHEMIGATION

Systems using gravity flow pesticide dispensing systems must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops.

VectoBac AS is metered or dripped into the rice floodwater at application stations positioned at the point of introduction (levee cut) of water into each rice field or pan. Two pints of VectoBac AS are diluted in water to a final volume in 5 gallons. The diluted

solution is contained in a 5 gallon container and metered or dispersed into the irrigation water using a constant flow device at the rate of 80 ml per minute.

Introduction of the solution should begin when 1/3 to 1/2 of the pan or field is covered with floodwater. Delivery of the solution should continue for a period of approximately 4-1/2 hours. Floodwater depth should not exceed 10-12 inches to prevent excessive dilution of VectoBac AS which could result in reduced larval kill. Agitation is not required during the period in which the VectoBac AS solution is being dispersed.

Application of VectoBac AS into rice floodwater is not permitted using a pressurized water and pesticide injection system.

Greenhouse - Drip (Trickle) and Sprinkler Chemigation

Public water system means a system for the provision to the public of piped water for human consumption if such system has a least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of the pesticide introduction. As an option to the RPZ the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a water pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials compatible with pesticides and capable of being fitted with a system interlock.

The system must contain a functional check valve, vacuum relief

valve and low pressure drain, appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

If VectoBac AS needs to be diluted prior to injection, material may be mixed separately and then put into the injector. Agitation may be necessary if materials are kept more than one day. VectoBac AS may be applied continuously.

Where supply tanks are used for continuous application, fill the supply tank with the desired quantity of water. Start mechanical or hydraulic agitation to provide moderate circulation before adding VectoBac AS. VectoBac AS suspends readily in water and will stay suspended over normal application periods. Avoid continuous agitation of spray mixture during spraying. Brief recirculation may be necessary if the spray mixture has sat for several hours.

NOTICE TO USER

Seller makes no warranty, express or implied, of merchantability, fitness or otherwise concerning the use of this product other than as indicated on the label. User assumes all risk of use, storage or handling not in strict accordance with accompanying directions.

Chemical and Agricultural Products Division
Abbott Laboratories
North Chicago, Illinois 60064

(c) 1992