

Storage and Disposal: Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate water, food, or feed by storage or disposal

Storage: Store in a cool, dry place

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag in application equipment, then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

General Information: Dipel[®] 10G is a granular formulation of Bacillus thuringiensis, variety kurstaki, designed specifically as a selective insecticide for control of European Corn Borer infesting seed corn, field corn, popcorn and sweet corn. For practical determinations of active ingredient, Dipel 10G can be expressed as a 10% formula of Dipel WP. The active ingredient in Dipel 10G is a stomach poison and therefore must be eaten by larvae to be effective. Climatic extremes such as heavy rain or strong wind shortly after treatment may act to remove Dipel 10G from the feeding zone making re-treatment necessary.

Directions for Use: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Application (Ground or Aerial Equipment):

<u>Crop</u>	<u>Pest</u>	<u>Rate</u>
Seed Corn, Field Corn, Popcorn, and Sweet Corn	European corn Borer (<u>Ostrinia nubilalis</u>), First and Second Generation	10 lbs/acre

First Generation

Apply Dipel[®] 10G directly to corn whorls. Treat seed corn when 25 percent of plants show shot hole feeding in the whorls and field corn when 35 percent of whorls show feeding. Larvae should be small and present at this time. Satisfactory control cannot be expected if treatment is delayed to higher injury levels or after larvae have begun stalk boring.

Second Generation

Apply Dipel 10G when a field count shows 100 egg masses per 100 plants and the first hatch is occurring. Under conditions of intensive population pressure, a second application may be necessary.

NOTICE

Local conditions may affect the use of DIPEL 10G. Consult State Agricultural Extension or Experiment Station specialists for specific recommendations related to local crop protection problems. Abbott makes no warranty, express or implied, including the warranties of commercially acceptable quality and/or fitness for any particular purpose concerning this material, other than those which are contained on this label.

DIPEL[®] 10G (Granular)

(Bacillus thuringiensis, var. kurstaki)

BIOLOGICAL INSECTICIDE

For control of European Corn Borer
Larvae on Seed Corn,
Field Corn, Popcorn, and Sweetcorn

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Active Ingredient: Bacillus thuringiensis,
var. kurstaki, 1600 International Units of
Potency per mg. * 0.32%
Inert ingredients 99.68%

*This product contains 0.726 Billion IU/lb.

EPA Reg. No. 275-55
EPA Est. No. 34704-NB-1

Net Weight

Lot No.

275-55



Chemical and Agricultural Products Division
Abbott Laboratories
North Chicago, Illinois 60064

Com. No 95M-81

Abbott 275-55
Draft Label
9/7/84
Page 1 of 2